

Original research article

## A retrospective observational clinic-pathological study of post-dated pregnancy

Dr. Reena Kumari<sup>1</sup>, Dr. Kusum Kumari<sup>2</sup>

<sup>1</sup>Associate Professor, Department of obstetrics and gynaecology, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India.

<sup>2</sup>Senior Resident, Department of obstetrics and gynaecology, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India.

Corresponding Author: Dr. Reena Kumari

E-mail: drreenakumari7@gmail.com

### Abstract

**Aim:** To find out incidence of post-dated pregnancies, associated maternal complications and perinatal morbidity and mortality.

**Material and Methods:** This was a retrospective observational study conducted in the Department of obstetrics and gynaecology, Anugrah Narayan Magadh Medical College, Gaya, Bihar, India for 15 months. Total 160 patients in the antenatal ward and labour room were selected for the study and they were divided into two groups, Control group with Gestational age 37-40 weeks and Study group with Gestational age >40 weeks. The maternal outcome was noted in terms of need for cesarean section, postpartum haemorrhage and sepsis. Foetal outcome was noted in terms of intrapartum asphyxia, intrauterine foetal death, admission to neonatal intensive care unit etc.

**Results:** The maximum number of patients belonged to the age group of 25-30 years both in cases (56.25%) and control group (53.75%) and the maximum number of patients in the study group (75%) belonged to the gestational age of 40-41 weeks while all the controls belonged to 37-40 weeks gestational age. In this study 63.75% of the patients in study group were primigravida and in the control group 53.75% were primigravida. The study group the percentage of LSCS was 28.75% which was higher than in the control group where it was 12.5%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 3.75%). Among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium stained liquor (11.25%) followed by cephalopelvic disproportion (6.25%). In the control group, most common indication was non progress of labour (5%) followed by acute foetal distress (3.33%) and non reactive CTG (3.33%). The maternal complications like LSCS, PPH and sepsis all were higher in the study group as compared to the control group. Among the foetal outcomes, 13.75% of infants in the study group had asphyxia as compared to only 6.25% in the control group. 15% infants of the study group had to be admitted to the NICU as compared to 11.25% in the control group. 1.25% was the percentage of intrauterine deaths in the study group as compared to none in the control group.

**Conclusion:** Considering this, policy of early intervention should be undertaken in post-dated pregnancy to avoid maternal and perinatal complications.

**Keywords:** Maternal complications, Post datism, Perinatal morbidity

### Introduction

Post dated pregnancy is defined as one which has crossed expected date of delivery. Pregnancy more than 42 weeks or more than 294 days is called post term pregnancy.

Fernandos Arias defined prolonged pregnancy as those pregnancies advancing beyond the expected date of delivery (EDD).<sup>1</sup> Prolongation of pregnancy complicates up to 10% of all pregnancies and carries increased risk to mother and fetus.<sup>2,3</sup> Post term perinatal mortality is greater than that of term pregnancy in almost all studies reviewed.<sup>4</sup> The growth and survival of most post dated infants suggests that the placenta uncommonly deteriorates with increasing length of gestation; thus the changes seen in fetuses afflicted with post maturity syndrome may not be explained by placental findings alone. Vorherr described critical reductions of fetal oxygen supply after 43rd week of gestation by cord blood oxygen content determinations.<sup>4</sup> The combination of continued fetal growth and arrested placental growth may lead to situation of decreasing placental nutrient reserve, compromised fetal circulation and eventually fetal distress. However, a recent electron microscopy study of placental changes in prolonged pregnancy suggests that the uteroplacental ischemia and not placental aging may be more important in genesis of post maturity syndrome.<sup>5</sup> Prolonged pregnancies are associated with an increased incidence of macrosomia. Macrosomia infants account for about 1% of term deliveries and 3-10% of post term deliveries.<sup>6</sup> Post maturity infants particularly with macrosomia and post maturity are at increased risk of hypoglycemia. They also have increased chance of polycythemia.<sup>7</sup> The maternal risks of post-dated pregnancy are often underappreciated. These include an increase in labor dystocia (9-12% vs 2-7% at term), an increase in severe perineal injury (3rd and 4 th degree perineal lacerations) related to macrosomia (3.3% vs 2.6% at term) and operative vaginal delivery, and a doubling in the rate of cesarean delivery (14% vs 7% at term).<sup>8-11</sup> The latter is associated with higher risks of complications such as endometritis, hemorrhage, and thromboembolic disease.<sup>10,12</sup> As there is fetal and maternal risk associated with post dated pregnancy, need of induction is more with post-dated pregnancy. There is general consideration that perinatal mortality and morbidity controversy is centered on adequacy of detecting different methods for the fetus at risk, the time when testing should be done, the method of monitoring, optimum time for delivery and mode of delivery. The availability of biophysical profile and electronic fetal monitoring can affect the outcome of a given pregnancy. One recent systematic review showed that a policy of labor induction for women with post-dated pregnancy compared with expectant management is associated with fewer perinatal deaths and fewer Caesarean sections.<sup>13</sup> The aim of this study was to find out incidence of post-dated pregnancies, associated maternal complications and perinatal morbidity and mortality.

### **Material and methods**

This was a retrospective observational study conducted in the Department of obstetrics and gynaecology, Anugrah Narayan Magadh Medical college, Gaya, Bihar, India for 15 months. Total 160 patients in the antenatal ward and labour room were selected for the study and they were divided into two groups, Control group with Gestational age 37-40 weeks and Study group with Gestational age >40 weeks

### **Inclusion criteria**

- Singleton pregnancy
- Cephalic presentation
- Absence of any other maternal complication

### **Exclusion criteria**

- Previous cesarean section
- Gestational hypertension
- Gestational diabetes
- Malpresentation

- Abruptio
- Placenta previa

### Methodology

All the data regarding the age, parity gestational age, any maternal complications like oligohydromnios, intrauterine growth restriction etc was collected. The maternal outcome was noted in terms of need for cesarean section, postpartum haemorrhage and sepsis. Foetal outcome was noted in terms of intrapartum asphyxia, intrauterine foetal death, admission to neonatal intensive care unit etc.

### Results:

Table 1 shows that maximum number of patients belonged to the age group of 25-30 years both in cases (56.25%) and control group (53.75%).

**Table 1: Age wise distribution of cases and controls**

Age (Years)	Number of Cases (%)	Number of Controls (%)
Below -25	23(28.75%)	24(30%)
25-30	45(56.25%)	43(53.75%)
Above 30	12(15%)	13(16.25%)
Total	80	80
Mean $\pm$ SD	26.7 $\pm$ 4.11	27.1 $\pm$ 4.07

**Table 2: Distribution of cases and controls by gestational age**

Period of gestation	Number of Cases (%)	Number of Controls
37-40 weeks	0	80(100%)
40-41 weeks	60 (75%)	0
41-42 weeks	20 (25%)	0
Total	80	80

Table 2 shows that the maximum number of patients in the study group (75%) belonged to the gestational age of 40-41 weeks while all the controls belonged to 37-40 weeks gestational age.

**Table 3: Distribution of cases and controls according to parity**

Parity	Number of cases (%)	Number of controls (%)
Primigravida	51 (63.75%)	43 (53.75%)
Multigravida	29 (36.25%)	37 (46.25%)
Total	80	80

As shown in table 3, 63.75% of the patients in study group were primigravida and in the control group 53.75% were primigravida.

**Table 4: Distribution of cases and controls according to the type of delivery**

Type of delivery	Number of cases (%)	Number of controls (%)
NVD	49 (61.25%)	67(83.75%)
Instrumental delivery	8 (10%)	3(3.75%)
LSCS	23 (28.75%)	10 (12.5%)
Total	80	80

Table 4 shows that in the study group the percentage of LSCS was 28.75% which was higher than in the control group where it was 12.5%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 3.75%)

**Table 5: Distribution of cases and controls according to the indication of LSCS**

Indication of LSCS	Number of Cases (%)	Number of Controls (%)
Acute foetal distress/MSL	9(11.25%)	3(3.75%)
Failed induction	3(3.75%)	0
Non progress of labour	3(3.75%)	4(5%)
Non reactive CTG	3(3.75%)	3(3.75%)
CPD	5(6.25%)	0
Total	23	10

As shown in table 5, among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium stained liquor (11.25%) followed by cephalopelvic disproportion (6.25%). In the control group, most common indication was non progress of labour (5%) followed by acute foetal distress (3.33%) and non reactive CTG (3.33%).

**Table 6: Distribution of cases and controls according to maternal complications**

Maternal complication	Number of cases	Number of controls
LSCS	23 (28.75%)	10 (12.5%)
PPH	13 (16.25%)	5 (6.25%)
Sepsis	12 (15%)	4 (5%)
Total	48	19

Table 6 shows the maternal complications like LSCS, PPH and sepsis all were higher in the study group as compared to the control group.

**Table 7: Distribution of cases and controls according to the foetal outcome**

Foetal outcome	Number of Cases (%)	Number of Controls (%)
No asphyxia	56 (70%)	66 (82.5%)
Fetal asphyxia (APGAR score < 6/10)	11 (13.75%)	5 (6.25%)
Admission to NICU	12 (15%)	9 (11.25%)
IUD	1 (1.25%)	0
Total	80	80

Among the foetal outcomes, as is depicted in table 7, 13.75% of infants in the study group had asphyxia as compared to only 6.25% in the control group. 15% infants of the study group had to be admitted to the NICU as compared to 11.25% in the control group. 1.25% was the percentage of intrauterine deaths in the study group as compared to none in the control group.

### Discussion

The present study was conducted to find out the incidence of maternal complications, perinatal mortality and morbidity in post-dated pregnancies. Total cases were 160 which were enrolled based on inclusion and exclusion criteria. In this study the maximum number of patients belonged to the age group of 25-30 years both in cases (56.25%) and control group (53.75%). Beischer in his study found that majority of post-dated patients belonged to the age group of 25-30 years, while Bancroft et al found that majority of patients belonged to 21-30 years.<sup>14,15</sup> Reddy UM et al found in their study that women who are of advance maternal age are at higher risk of still birth throughout gestation, the peak risk period is 37 to 41 weeks.<sup>16</sup>

In the study, the mean age being  $26.7 \pm 4.11$  years in the study group. Similar studies by Mahapatro<sup>18</sup> and Eden et al<sup>17</sup> have shown the mean age to be  $24.19 \pm 3.30$  and 25.8 years respectively. 63.75% of the patients in study group were primigravida which is similar to Mahapatro<sup>18</sup> and Alexander et al's<sup>19</sup> study.

Among the mode of delivery, in the study group the percentage of LSCS was 28.75% which was higher than in the control group where it was 12.5%. Incidence of instrumental delivery was also higher in the study group as compared to control group (10% as compared to 3.75%). In a similar study by Mahapatro<sup>18</sup> the rate of LSCS was found to be 28.9% and that of instrumental delivery was 5.72%. In study by Singhal et al.<sup>20</sup> the rate of LSCS was found to be 14.7% and that of instrumental delivery was 8.6%. Davinder et al.<sup>21</sup> study showed the rate of instrumental delivery as 10.35%. In this study among the indications for LSCS, the most common indication among the study group was acute foetal distress which includes meconium stained liquor ((11.25%) followed by cephalopelvic disproportion (6.25%). In the control group, most common indication was non progress of labour (5%) followed by acute foetal distress (3.33%) and non reactive CTG (3.33%). Bhriegu R et al<sup>22</sup> in their study also found that Meconium stained liquor with fetal distress was the most common indication for LSCS (23.5%) and in Mahapatro's<sup>18</sup> study, again fetal distress was found to be the most common indication for LSCS (65.5%). In our study, the maternal complications like LSCS, PPH and sepsis all were higher in the study group as compared to the control group. Among the foetal outcomes, 13.75% of infants in the study group had asphyxia as compared to only 6.25% in the control group. 15% infants of the study group had to be admitted to the NICU as compared to 11.25% in the control group. 1.25% was the percentage of intrauterine deaths in the study group as compared to none in the control group. Bhriegu R et al<sup>22</sup> in their study, also found increased incidence of obstetric complications such as rate of LSCS, perineal tear, atonic postpartum haemorrhage, and perinatal complications such as fetal distress and meconium aspiration syndrome. Similar studies by Singhal et al.<sup>20</sup> and Alexander et al<sup>19</sup> also revealed increased incidence of maternal and perinatal complications like increased LSCS rate, low Apgar scores and admission into NICU.

### Conclusion

Considering this, policy of early intervention should be undertaken in post-dated pregnancy to avoid maternal and perinatal complications.

### References

1. Fernando Arias. Practical guide to high risk pregnancy and delivery 3 Edition; India Harcourt private limited; 2008:255-270.
2. Olesen AW, Westergaard JG, Olsen J. Perinatal and maternal complications related to post term delivery: a national register-based study, Am. J Obstet Gynecol. 1978-1993;189:222-227.
3. Norwitz ER, Snegovskikh VV, Caughey AB Prolonged pregnancy: when should we intervene? Clin Obstet Gynecol. 2007;50:547-57.
4. Vorherr H. Placental insufficiency in relation to postterm pregnancy and fetal postmaturity: Evaluation of fetoplacental function Management of post term gravida. Am J Obstet Gynecol. 1975;123:67.
5. Jones JP, Fox H. Ultrastructure of the placenta in prolonged pregnancy. J Pathol. 1978;126(3):173-9.
6. Spellacy WN, Miller S, Winegar A, Peterson PQ. Macrosomia maternal characteristics and infant complications. 1985;66(2):158-61.
7. Ratnam SS, Arulkumaran S. Post term infant. Obstetrics and Gynecology, 2 nd Edition, India Orient Longman. 2003;2:48-53.

8. Rand L, Robinson JN, Economy KE. Post-term induction of labor revisited. *Obstet Gynecol.* 2000;96(5 Pt 1):779-83.
9. Campbell MK, Ostbye T, Irgens LM. Post-term birth: risk factors and outcomes in a 10-year cohort of Norwegian births. *J Obstet. Gynecol.* 1997;89(4):543- 8.
10. Alexander JM, McIntire DD, Leveno KJ. Forty weeks and beyond: pregnancy outcomes by week of gestation. *Obstet Gynecol.* 2000;96(2):291-4.
11. Treger M, Hallak M, Silberstein T. Post-term pregnancy: should induction of labor be considered before 42 weeks? *J Maternal Fetal Neonatal Med.* 2002;11(1):50-3.
12. Eden RD, Seifert LS, Winegar A. Perinatal characteristics of uncomplicated postdate pregnancies. *Obstet. Gynecol.* 1987;69:296-9.
13. Gulmezoglu AM, Crowther CA, Middleton P, et al; Induction of labour for improving birth outcomes for women at or beyond term. *Cochrane Database Syst Rev.* 2012;6:CD004945
14. Beischer NA, Evans JH, Townsend L. Studies in prolonged pregnancy. I: Incidence of prolonged pregnancy. *Am J Obstet Gynecol.* 1969;103:476.
15. Bancroft-Livingston G, Neill DW. Studies in prolonged pregnancy, Cord blood oxygen levels at delivery. *J Obstet Gynaecol.* 1957;64:498-503.
16. Reddy UM, KO CW, Willinger M. Maternal age and risk of stillbirth throughout pregnancy in the United states. *Am J Obstet Gynecol.* 2006;195(3):764-70.
17. Eden RD, Gergely RZ, Schifrin BS, Wade ME. Comparison of antepartum testing schemes for the management of the postdate pregnancy. *Am J Obstet Gynecol* 1982;144:683-92.
18. Mahapatro A. Fetomaternal outcome in pregnancy beyond 40 weeks. *Int J Pharm Bio Sci* 2015;6:53-8
19. Alexander JM, McIntire DD, Leveno KJ. Forty weeks and beyond: Pregnancy outcomes by week of gestation. *Obstet Gynecol* 2000;96:291-4.
20. Singhal P. Fetomaternal outcome following postdate pregnancy. A prospective study. *J Obstet Gynecol India.* 2001;51:89-93.
21. Kaur D, Saini AS, Kaur J. Maternal and fetal outcome in postdated pregnancies. *J Obstet Gynecol India* 1997;47:331-4.
22. Bhriegu R, Agrawal M, Hariharan C. Assessment of maternal and perinatal outcome in postdated pregnancy. *J Datta Meghe Inst Med Sci Univ* 2017;12:35-40.

Received: 20-08-2020 || Revised: 12-09-2020 || Accepted: 27-10-2020