

Original research article

Role of IgM antibody capture ELISA for the early diagnosis of dengue viral infection- Prospective observational study**Dr. Krishna Kumar Mani¹, Dr. Abhimanyu Kumar Singh², Dr. Kumar Amresh³, Dr. Ranjan Kumar⁴****¹Associate Professor, Department of Microbiology Jawaharlal Nehru Medical College and Hospital, Bhagalpur, Bihar, India.****²Tutor, Department of Microbiology Jawaharlal Nehru Medical College and Hospital, Bhagalpur, Bihar, India.****³Assistant Professor, Department of Microbiology Jawaharlal Nehru Medical College and Hospital, Bhagalpur, Bihar, India.****⁴Tutor, Department of Microbiology Jawaharlal Nehru Medical College and Hospital, Bhagalpur, Bihar, India.****Corresponding Author: Dr. Abhimanyu Kumar Singh****Email id: drabhi.2004@gmail.com****Abstract**

Background: The most challenging problem associated with patient management in Dengue infection is early diagnosis. Secondary infection with dengue virus is the most accepted risk factor for the development of dengue haemorrhagic fever. Serologic diagnosis of dengue virus infection using ELISA of both IgM and IgG distinguishes primary and secondary infections.

Aim: to determine the seroprevalence of dengue viral infection using IgM antibody capture ELISA for the early diagnosis in Bihar region.

Material and methods: This was a prospective observational study conducted in the Department of Microbiology Jawaharlal Nehru medical college and Hospital, Bhagalpur, Bihar, India for 1 year. A total of 570 serum samples from suspected dengue cases attending OPD or admitted in the hospital were tested for the confirmation of Dengue. We have received blood samples in our microbiology laboratory, the blood samples were allowed to clot at room temperature and then we centrifuged the samples and serum samples were separated. From the serum samples we have done NS1 Ag and IgM Ab testing by ELISA. **Results:** Out Of 570, 110 samples were positive for dengue. Seroprevalence of Dengue was 19.30%. Out of 110 dengue patients 72 (65.46 %) were male patients and 38 (27.14 %) were female patients. Out of 110 dengue patients, 80(72.73%) patients were from urban area and 30(27.27%) from rural area. In our study dengue infection was observed more (30.90%) in the age group 20 to 30 years followed by 10 to 20 years (23.64%) and 30 to 40 years (21.82%). All dengue positive patients in our study had fever of 2 to 7 days. The most common presenting symptoms of dengue were fever with body ache (46.36%), headache (37.27%), nausea (33.64%) and vomiting (22.73%). Out of 110 dengue cases fever with rash was observed in 8 cases (7.27%).

Conclusion: The present results revealed that the study region is epidemic for dengue viral infection and there is an urgent need for the constant monitoring to control further spreading of the infection in the community, hence serological test have important role in the early diagnosis.

Keywords: IgM ELISA, Dengue viral infection, Aedes aegypti.

Introduction

Dengue is a mosquito-borne viral disease that has quickly spread in all regions of World Health Organization (WHO) in recent years. Female mosquito mainly of the species *Aedes aegypti* and to a lesser point, *A. Albopictus*, transmit dengue virus.¹ Dengue infection is a major health problem in our country. Worldwide the occurrence of dengue has increased in the recent years. The WHO approximates that presently about two-fifth of the population is at risk for this viral infection. In recent years dengue in Asian and Latin American countries are common, where it has become the leading cause of hospitalization among both adults and children in these regions, India is one of the recognized countries in South-East Asian region frequently reporting occurrence of dengue fever/dengue hemorrhagic fever (DF/DHF) outbreaks and transforming into surrounding regions and nearby niches for dengue infection.² The Infection with any one serotype confers an individual life-long immunity to that same serotype but it has cross reactivity to the other serotype. Secondary infection with another serotype or multiple infections with different serotypes leads to severe form of dengue dengue hemorrhagic fever (DHF) and dengue shock syndrome (DSS) due to this cross-reactivity.³ The incidence of dengue has increased over last 50 years with 2.5 billion people living in areas where dengue is endemic.⁴ It affects 100 million people each year with 500,000 cases of DHF and DSS with around 30,000 deaths mostly among children.¹ It is known that early and specific diagnosis of DHF and DSS followed by supportive therapy reduces mortality and morbidity.⁵ Viral Isolation by cell culture and subsequent detection by immunofluorescence, though the gold standard tests for identification of dengue infection are not within the reach of peripheral and even most tertiary care laboratories.⁶ For a long time, detection of dengue specific IgM/IgG has been the main stay of diagnosis of dengue infection. Antibody detection is an indirect method of diagnosis and therefore is prone to false positive as well as false negative results.⁷ NS1 antigen is detectable from day 1 of fever both in primary and secondary infections. NS1 is shown to be highly specific viral marker making it extremely reliable parameter for diagnosis of dengue infection from day 1 of fever.⁸

A small percentage of persons who have previously been infected by one dengue serotype develop bleeding and endothelial leak upon infection with another dengue serotype. This syndrome is termed severe dengue (reclassified in 2009 by the WHO, previously referred to as dengue hemorrhagic fever and dengue shock syndrome). Severe dengue has also been termed dengue vasculopathy. Vascular leakage in these patients results in hemo concentration and serous effusions and can lead to circulatory collapse. This, in conjunction with severe hemorrhagic complications, can lead to a shock syndrome, which poses a greater fatality risk than bleeding per se.⁹ Dengue is endemic to the Indian sub-continent. Dengue is associated with explosive urban epidemics and has become a major public health problem in India.¹⁰

Material and methods

This was a prospective observational study conducted in the Department of Microbiology Department of Microbiology Jawaharlal Nehru medical college and Hospital, Bhagalpur, Bihar, India for 1 year.

Methodology

A total of 570 serum samples from suspected dengue cases attending OPD or admitted in the hospital were tested for the confirmation of Dengue. All the age group patients were include in this study. A suspected case of dengue was considered a patient with signs and symptoms like headache, retro-orbital pain, myalgia, arthralgia, rash and haemorrhagic manifestation, etc.

Serum samples from these patients were tested for Dengue NS1 antigen using dengue NS1 antigen capture ELISA (PanBio Diagnostics) and dengue IgM antibody by dengue IgM capture ELISA (PanBio Diagnostics) for the confirmation of dengue cases. ELISA tests were

performed as per the manufacturer's instructions. We have received blood samples in our microbiology laboratory, the blood samples were allowed to clot at room temperature and then we centrifuged the samples and serum samples were separated. From the serum samples we have done NS1 Ag and IgM Ab testing by ELISA.

Results:

Table 1: Seroprevalence of Dengue

Total no of patients	Dengue positive patients	%
570	110	19.30

Table 1: Out Of 570, 110 samples were positive for dengue. Seroprevalence of Dengue was 19.30%.

Table 2: Demographic profile of patients

Gender	N=110	%
Male	72	65.46%
Female	38	27.14%
Age years		
Below 10	10	9.10%
10-20	26	23.64%
20-30	34	30.90%
30-40	24	21.82%
40-50	9	8.18%
Above 50	7	6.3%
Area		
Urban	80	72.73%
Rural	30	27.27%

Out of 110 dengue patients 72 (65.46 %) were male patients and 38 (27.14 %) were female patients. Out of 110 dengue patients, 80(72.73%) patients were from urban area and 30(27.27%) from rural area. In our study dengue infection was observed more (30.90%) in the age group 20 to 30 years followed by 10 to 20 years (23.64%) and 30 to 40 years (21.82%).

Table 3: Clinical profile of dengue patients

Clinical presentation	No of Patients	%
Fever + myalgia	13	11.82%
Fever + rash	8	7.27%
Fever + headache	41	37.27%
Fever+ nausea	37	33.64%
Fever + vomiting	25	22.73%
Fever + arthralgia	17	15.45%
Fever + bodyache	51	46.36%
Fever + itching	14	12.73%

All dengue positive patients in our study had fever of 2 to 7 days. The most common presenting symptoms of dengue were fever with body ache (46.36%), headache (37.27%), nausea (33.64%) and vomiting (22.73%). Out of 110 dengue cases fever with rash was observed in 8 cases (7.27%).Table 3.

Table 4: Serology results of rapid dengue tests

Test results	No. of patients	%
NS1/NS1+IgM/IgM Positive	97	88.18%

IgG Positive	7	6.36%
IgG + IgM Positive	6	5.45%
Total	110	100%

Out of 110 dengue cases, NS1/NS1+IgM/IgM were positive for 97(88.18%) patients, suggesting primary infection. IgM and IgG positive was seen in 7(6.36%) patients, suggesting late primary or early secondary infection. IgG was positive in 6(5.45%) cases, suggesting secondary or past infection. Out of all dengue cases thrombocytopenia ($<1,00,000/\text{mm}^3$) was observed in 41 cases. In 6 patients platelet count was $<20,000/\text{mm}^3$

Discussion

Serological diagnosis of dengue virus infection using a commercial capture ELISA of both IgM and IgG distinguishes primary and secondary infections is preferred. ELISA is a simple, reliable and cost effective method in the diagnosis of dengue infection. Total 570 blood samples of the patients suspected of having dengue infection were tested in the laboratory by rapid immuno chromatography tests for NS1 Ag, IgG and IgM. Out of these 110 samples were positive for dengue. Seroprevalence of Dengue was 19.30%. 11.92% prevalence was reported by P. Jyoti and B Metri.¹¹ 18.99% prevalence was observed over period of 2008 to 2011 by Smita Sood in Rajasthan.¹² Low prevalence 3.55% was reported by Mahesh kumar et al.¹³ A study from central; India reported 31.3% prevalence rate.¹⁴

Out of 110 dengue patients 72 (65.46 %) were male patients and 38 (27.14 %) were female patients. Similar result was observed by Mahesh kumar et al, in their study out of total positive dengue cases, 62.63% were males and 37.37% females.¹³ Many studies have observed higher prevalence of dengue infection among males than females.^{11,12,15,16} S. Fayaz Ahammad et al reported 46.6% male & 53.4 female dengue patients.¹⁶ Study by Kale A V et al reported 63.33% were males & 36.66% were females.¹⁵

In our study, out of 110 dengue patients, 80(72.73%) patients were from urban area and 30(27.27%) from rural area. similar results was by S. Fayaz Ahammad et al. (2016), 109 cases (75%) were from rural area where as 25 cases (25%) were from urban area.¹⁶ According to their report the rural broaden of dengue infection is comparatively a recent phenomenon which is supposed to be linked with the shortage of water in rural areas, designing of schemes for water supply to the rural areas and development of newer water transport system in the rural places.

In our study dengue infection was observed more (30.90%) in the age group 20 to 30 years followed by 10 to 20 years (23.64%) and 30 to 40 years (21.82%). Mahesh Kumar et al in their study observed maximum dengue cases in age group 10-20 years (31.58%) and 21 to 30 yrs. (15.78%).¹³ Kale et al, observed commonest age group affected was (34%) was between 11-15 years.¹⁵ Some Indian studies have reported that dengue infection is more common in children.^{17,18}

All dengue positive patients in our study had fever of 2 to 7 days. The most common presenting symptoms of dengue were fever with body ache (46.36%), headache (37.27%), nausea (33.64%) and vomiting (22.73%). Out of 110 dengue cases fever with rash was observed in 8 cases (7.27%). Similar clinical presentation was observed by Mahesh Kumar et al, fever was present in almost all cases (n=380) followed by, headache (n=274), joint pain (n=2432), myalgia (n=144), retro-orbital pain (n=141), backache (n=95), skin rash (n=80).¹³

Out of 110 dengue cases, NS1/NS1+IgM/IgM were positive for 97(88.18%) patients, suggesting primary infection. IgM and IgG positive was seen in 7(6.36%) patients, suggesting late primary or early secondary infection. IgG was positive in 6(5.45%) cases, suggesting secondary or past infection. Mahesh kumar et al reported that, Out of the 380 dengue positive cases, 136(35.79%) were NS-1 positive, 117(30.79%) were IgM positive, 38(10%) were IgG

positive, 71(18.68%) were IgG/IgM positive, 14(3.68%) were IgG NS- 1/IgMNS-1 positive and 4(1.05%) were IgGIgMNS-1 positive.¹³

Though among methods used for diagnosis of dengue the virus isolation, molecular methods are more specific tests, facilities are not available in all institutes. Serological tests are most commonly used in most of the laboratories. Dengue virus specific IgM antibodies tend to appear as early as 3 days after infection and remains in circulation for 30 to 60 days. IgG antibodies arise at about 7 days, they reach a peak at 2-3 weeks and persists for life long.¹⁸ NS1 detection has been a promising test to diagnose dengue in its early febrile stage. The NS1 protein was found to be highly conserved in all dengue serotypes, circulating in high levels during the first few days of illness. It correlates with the development of Dengue Fever. There is no cross reaction of the dengue NS1 protein with those of other related *flavi viruses*.^{19,20} Out of all dengue cases thrombocytopenia (<1, 00,000/mm³) was observed in 41 cases. In 6 patients platelet count was < 20,000/mm³. One of the WHO diagnostic criteria for DHF is Thrombocytopenia: <1 lakh/mm³. P Jyoti and Basawaraj reported thrombocytopenia in 51.5% patient.¹¹ Kale A V et al observed thrombocytopenia in 56% patients, platelet count <40,000 in 33.33% cases.¹⁵ Platelet count less than 1, 00,000/ml was noticed in 220 cases (68.75%), report published by R D Kulkarni et al.²¹

Conclusion

The present results revealed that the study region is epidemic for dengue viral infection and there is an urgent need for the constant monitoring to control further spreading of the infection in the community, hence serological test have important role in the early diagnosis. Therefore IgM ELISA is recommended in all the suspected dengue patients so as to instigate essential treatment and assessment of morbidity and mortality rate during an outbreak.

Reference:

1. World Health Organization. Dengue and severe dengue. Fact Sheet No. 117, February 2015; WHO, Geneva
2. Ashwini K, Chythra RR, Vinay P, Seema S, Chanaveerappa B, Charmaine MS et al. Clinical manifestations and trend of dengue cases admitted in a tertiary care hospital, Udupi District, Karnataka. Indian J Community Med 2010; 35(3):386–90.
3. World Health Organization. Dengue: Comprehensive guidelines for prevention and control of dengue and dengue hemorrhagic fever, revised and expanded edition; Chapter 4, clinical manifestations and diagnosis.2011:24.
4. Smith AW, Chen LH, Massad E, Wilson ME. Threat of dengue to blood safety in dengue – endemic countries. Emerg Infect Dis.2009;15(1):8-11.
5. Peters CJ. Infections caused by arthropod – and Rodent borne viruses. In : Fauci AS, editor. Harrison's Principles of Internal Medicine 17th ed. New York: McGraw Hill Medical Publishing Division.2008:1226- 39.
6. Chakravarti A, Kumaria R, Batra VV, Varma V. Improved detection of dengue virus serotypes from serum samples – Evaluation of single-tube multiplex RT-PCR with cell culture. Dengue Bulletin.2006;30:133-40.
7. Peeling RW, Artsob H, Pelegrino JL, Buchy P, Cardoso MJ, Devi S. Evaluation of diagnostic test: Dengue. Nat Rev Microbiol. 2010;8:S30–7.
8. Datta S, Wattal C. Dengue NS1 antigen detection: A useful tool in early diagnosis of dengue virus infection. Indian J Med Microbiol. 2010;28(2):107-10.
9. Statler J, Mammen M, Lyons A, Sun W. Sonographic findings of healthy volunteers infected with dengue virus. J Clin Ultrasound. 2008;36(7):413-7.

10. Mahesh Kumar, Sharma R, PariharG, SharmaM. Seroprevalence of Dengue in Central Rajasthan: A Study at a Tertiary Care Hospital. *Int. J. Curr. Microbiol. App. Sci.*2015;4(9): 933-940
11. Parameswarappa Jyothi, Basavraj C Metri. Correlation of serological markers and platelet count in the diagnosis of Dengue virus infection. *Adv Biomed Res.*2015;4:26
12. Smita Sood A Hospital Based Serosurveillance Study of Dengue Infection in Jaipur (Rajasthan), India *Journal of Clinical and Diagnostic Research.* 2013 Sept, Vol-7(9): 1917-1920
13. Mahesh Kumar, Sharma R, PariharG, SharmaM. Seroprevalence of Dengue in Central Rajasthan: A Study at a Tertiary Care Hospital. *Int. J. Curr. Microbiol. App. Sci* (2015) 4(9): 933-940
14. Ukey PM, Bondade SA, Paunipagar PV, Powar RM, Akulwar SL. Study of Seroprevalence of Dengue Fever in Central India. *Ind J Community Med.* 2010; 35(4): 517-19
15. Kale AV et.al, clinical profile and outcome of dengue fever from a tertiary care centre at Aurangabad Maharashtra India: an observational study, *IOSR journal of dental and medical sciences*, volume 13, issue 9 ver. Vii sep. 2014; 14-19
16. S. Ahammad F et al. Clinico- demographic profile of dengue fever in a southindian tertiary care teaching hospital. *World Journal of Pharmacy and Pharmaceutical Sciences* Vol 5, Issue 1, 2016. 1602-1609
17. Garg A, Garg J, Rao YK, Upadhyay GC, Sakhuja S. Prevalence of dengue among clinically suspected febrile episodes at a teaching hospital in North India. *J Infect Dis and Immun.* 2011; 3(5):85-89.
18. Vijaykumar TS, Chandy S, Satish N, Abraham M, Abraham P, et al. Is dengue an emerging as a major public health problem? *Ind J Med Res.* 2005;121:100-07
19. Shrivastava A, Dash PK, Tripathi NK, Sahni AK, Gopalan N, Lakshmana Rao PV. Evaluation of a commercial Dengue NS1 enzyme-linked immunosorbent assay for early diagnosis of dengue infection. *Indian J Med Microbiol*, 2011; 29(1):51-5.
20. Datta S, Wattal C. Dengue NS-1 antigen detection: A useful tool in early diagnosis of dengue virus infection. *Indian J Med Microbio* 2010; 28:107- 190.
21. Kulkarni R D, Patil S S, Ajantha G S, Upadhya A K, Kalabhavi A S, Shubhada R M, Shetty P C, Jain P A. Association of platelet count and serological markers of dengue infection- importance of NS1 antigen. *Indian J Med Microbiol* 2011;29:359-62

Received :19-08-2020 Revised: 22-09-2020. Accepted:25-10-2020