

Original research article

A Cross-sectional Investigation of Measles Antibody Titres and Nutritional Condition in Children**Dr. Dheeraj Kumar¹, Dr. Sanjeev Kumar²****¹Assistant Professor, Department of Pediatrics, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India****²Assistant Professor & HOD, Department of Pediatrics, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India****Corresponding Author: Dr. Sanjeev Kumar****Abstract**

Aim: The aim of the present study to determine the association of measles antibody titres with nutritional status in paediatric population.

Methods: This was a cross-sectional study conducted in the Department of Pediatrics, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India for 1 year, on 1-10 year old children. Total 300 patients were include in this study for finding out the seroprevalence and anti-measles antibody levels, and studying their association with age, gender, as well as nutritional status of these children. Blood samples were tested for presence of measles specific IgG antibodies.

Results: The relationship between age and vaccination status was not found to be statistically significant ($p=0.22$). No statistically significant difference was observed in the baseline characteristics of vaccinated and unvaccinated group except for mean weight for age Z score which was significantly lower in the unvaccinated group ($p = 0.023$). According to weight for age, in well-nourished children seropositivity was 67.68% with GMT 889 mIU/ml, moderately malnourished (wasting) 68% with GMT 687 mIU/ml and in severely malnourished (severe wasting) 60.90% with GMT 548 mIU/ml. It was not found statistically significant with p value 0.05 for seropositivity but significant with p value 0.0001 for antibody levels. According to Height for age in well- nourished children seropositivity was 54.32% with GMT 685 mIU/ml, moderately malnourished (stunting) seropositivity was 71.36% with GMT 387 mIU/ml and in severely malnourished (severe stunting) seropositivity 46.67% with GMT 361 mIU/ml. P value noted was 0.005 which is significant. In weight for length/height both seropositivity and GMT were found insignificant in well-nourished, moderately malnourished as well as severely malnourished subject (P value: 0.47)

Conclusion: Nutritional status of children has an association with measles antibody titres as well GMT of measles specific IgG antibody, with those with better nutritional status having higher measles antibody titres.

Keywords: Measles, Vaccine, Antibody, Vaccination, Malnourished, Anthropometry

Introduction

Before the introduction of the mass immunization programme in 1978, measles was endemic with a 2-year epidemic cycle in Taiwan, China.¹ The most recent outbreak affecting the whole island occurred in 1988–89.² After that outbreak, a goal for measles elimination by 2000 was established in 1991. Towards achieving this, an MMR (measles, mumps and rubella combined vaccine) campaign targeting primary and secondary schoolchildren (6–15 years old) was conducted in 1991–94 and reached 90% of the target population.³ Meanwhile, a 2-dose policy with 1 dose of measles vaccine at 9 months of age and 1 dose of MMR at 15 months of age has been implemented since 1991. Based on the vaccine coverage survey conducted among

children aged 13–24 months in 1993–94, the mean coverage was 84% (range: 77–96%) for one dose of measles vaccine and 69% (range: 60–83%) for one dose of MMR vaccine.^{3 (3)} Recent health statistics showed that the MMR coverage in 2-year-old children in 1995 was about 80–85%.⁴ In addition, immunization requirements have been implemented by screening vaccination records and immunizing unvaccinated individuals among newcomers (6 years of age) to primary school since 1991 ('mop-up'). This policy has been extended to kindergarten since 1994. Under the strategy of 'mop-up', the overall vaccine coverage with at least 1 dose of MMR vaccine in primary schoolchildren was 96% in 1995.⁴ In addition to improving measles vaccine coverage, since 1992 measles surveillance has also been strengthened by incorporating active case investigation and laboratory diagnosis. Except for a small-scale outbreak with 33 confirmed cases in the northern part of the island in 1994, only sporadic confirmed cases were identified in 1992 (7 cases) and 1993 (2 cases). No confirmed cases were detected in 1995 and 1996. However, 7 confirmed cases, including 2 imported cases, were identified in 1997, and a small-scale outbreak with 9 confirmed cases occurred in the southern part of the island in 1998.³⁻⁶ Seroepidemiological studies conducted in Taiwan, China, in 1993–97 showed that measles IgG seroprevalence were 85–92% in preschool and kindergarten children (2–5 years old) and reached 92–98% in schoolchildren (6–15 years old) and young adults over 16 years old.⁶⁻⁹

In India, Measles contributes to 2.3 percent of all deaths and one tenth of all deaths in the preschool children.¹⁰ A review of community based studies of published measles outbreaks, investigations found a median case fatality ratio of 3.7%, range 0 to 23.9%, primarily affecting the remote tribal populations have been reported and case fatality rates of 5-30 percent have been observed. Despite the declining trends in measles cases from a reported 162560 in 1989 to 29339 in 2011¹¹, measles remains a major cause of morbidity & mortality in the children in India. Measles vaccine coverage in India is 74% (Urban-78% and rural-72%), especially 52.8% in Uttar Pradesh.¹² This is lower than the 95% coverage required to eliminate measles¹³, and much of the World Health Organization (WHO) African Region, including the Democratic Republic of the Congo (DRC), has even lower coverage than this worldwide average.¹⁴ Measles is endemic in the DRC.¹⁴ Currently, DRC gives one routine dose of measles vaccine to children nine months of age, and in outbreak settings, to children as young as six months. Although the WHO states that all countries should include a second routine dose of MV, regardless of national routine coverage level of the first dose⁴, this recommendation has not been implemented in the DRC. Because coverage achieved through healthcare is low in the DRC, attempts are made to reach missed children through Supplementary Immunization Activities (SIA), which Doshi et al. found to be associated with decreased measles incidence.¹⁵ The ability of an infant to seroconvert is agedependent due to level and decay of maternal antibodies and immunological development; regional differences in seroprevalence have been observed. Expectant mothers in endemic areas may be more likely to have had natural measles infection, resulting in higher measles antibody levels, and so pass on higher levels of measles antibody transplacentally to their infants, resulting in longer lasting protection than would occur in expectant mothers with vaccine-induced antibody.^{7,8} Children in measles-endemic regions are also at risk of exposure to measles at an earlier age, and this must be considered when determining ideal age of vaccination.⁴ Determining drivers of seropositivity and low vaccine effectiveness (VE) is complex and can depend on immunization program logistical capacity¹⁶, vaccine potency¹⁷ and host factors, particularly immune system robustness as a function of development (age) and nutritional status.^{18,19} Measles has occurred even in well-vaccinated populations, raising questions of why adequate protection is not achieved in such groups.²⁰ While vaccination induces humoral and cellular immune responses similar to those caused by

natural disease, the resulting antibody levels are lower among those with vaccine - induced versus natural immunity. As measles continues to be inadequately controlled in DRC.^{21,22}

Material and methods

This was a cross-sectional study conducted in the Department of Pediatrics, Vardhman Institute of Medical Sciences, Pawapuri, Nalanda, Bihar, India for 1 year, on 1-10 year old children, after taking the approval of the protocol review committee and institutional ethics committee. The technique, risks, benefits, results and associated complications of the procedure were discussed with all patients. Total 300 patients were include in this study for finding out the seroprevalence and anti-measles antibody levels, and studying their association with age, gender, as well as nutritional status of these children. Blood samples were tested for presence of measles specific IgG antibodies. Children in the age group of 1 to 10 years were included in this study. Children were received blood or blood components within last 3 months, Children received corticosteroid therapy or other immunosuppressive therapy, HIV positive children, Transplant recipients (bone marrow/ solid organ) and Received of gamma globulins within last 2 months were excluded from this study.

The techniques of measurement described in Cogill's (2003)²³ Anthropometric Indicators Measurement Guide were followed to make the following measurements. Weight was measured using a portable electronic weighing scale with a weighing capacity from 1 kg to 150 kg in 100g divisions, accuracy +/- 100g. Height: was measured in centimetres to a precision of 0.1cm by a wall mounted tape measuring up to 2 meters. An infantometer was used to measure the length for children less than 2 years of age. The following indices & their z scores were calculated: Body Mass Index (BMI) = $Weight (Kg) / Height (m)^2$. Weight for age: for children less than 10 years of age by W.H.O standard growth chart and z score was calculated. Height for age: for all children based on W.H.O standard growth chart and z score was calculated. Weight for height: for children less than 5 years based on W.H.O standard growth chart and zscore was calculated. Nutritional status of children was classified on the basis of the WHO Growth Standards, 2006 for 0-60 months; and the WHO Reference, 2007 for 5-19 years.

Results

Children 5-18 Years: Overweight: >+1SD (equivalent to BMI 25 kg/m² at 18 years) Obesity: >+2SD (equivalent to BMI 30 kg/m² at 18 years). Thinness: <- 2SD. Severe thinness: <-3SD. Children 0-5 years: Moderate wasting: weight-for length/ height Z -score -2 to -3 Severe wasting (severe acute malnutrition): weight-for-length/ height Z -score <-3. Overweight: BMI-for-age or weight-for-length/ height Z -score > 2. Obesity: BMI-for-age or weight for-length/ height Z -score>3. Moderate stunting: length/ height for age Z -score -2 to -3. Severe stunting: length/ height for age Z -score < -3. Blood samples were collected and serums were separated by centrifugation and stored at -22 degree Celsius till the time of assay. Measles specific IgG antibodies were detected by using a commercial IgG ELISA kit (Measles Virus IgG ELISA, IBL International GMBH) in accordance with the manufacturer's instructions

Table 1: Vaccination status of children against measles

Age group (years)	Vaccinated N (%)	Unvaccinated N (%)	Total	P value
1-10	200(66.67)	100(33.33)	300	0.22

Majority (66.67%) of the total subjects had been vaccinated against measles. A similar trend was observed in each of the age groups. However, the relationship between age and vaccination status was not found to be statistically significant (p=0.22)

Table 2: Baseline characteristics of measles vaccinated and unvaccinated children

Characteristics	Vaccinated mean±SD	Unvaccinated mean±SD	P value
Age (years)	7.5±2.8	6.5±2.9	0.28
Weight (kg)	17.2±5.8	16.6±5.9	0.049
Height (cm)	108.4±20.5	107.8±19.1	0.23
BMI (kg/m ²)	15.7±2.7	14.7±2.3	0.25
Weight for age Z score(1-10 years)	-1.7±1.1	-2.4±1.2	0.023
Height for age Z score	-1.5±0.8	-1.9±1.1	0.061
Weight for height Z score(1-5 years)	-1.7±1.6	-1.6±1.5	0.49
BMI Z Score	-1.7±3.2	-1.8±2.1	0.75

No statistically significant difference was observed in the baseline characteristics of vaccinated and unvaccinated group except for mean weight for age Z score which was significantly lower in the unvaccinated group ($p = 0.023$).

Table 3: Nutritional status of subjects

Parameter of Nutritional status	Total N (%)	Vaccinated N (%)	Unvaccinated N (%)	
Weight for age Z Score (age ≤10yrs)	<-3	54(18)	34(17)	20(20)
	-2 to -3	64(21.33)	40(20)	24(24)
	>-2	182(60.67)	126(63)	56(56)
	Total	300	200	100
Weight for Height Z Score (age ≤5yrs)	<-3	23(14.37)	16(16)	10(16.67)
	-2 to -3	32(20)	21(21)	13(21.67)
	>-2	105(65.63)	63(63)	37(61.67)
	Total	160	100	60
Height for Age Z Score	<-3	15(5)	10(5)	5(5)
	-2 to -3	81(27)	50(25)	31(31)
	>-2	206(68.67)	140(70)	64(64)
	Total	300	200	100
BMI for age Z score (age > 5yrs;	<-3	28(20)	19(19)	11(27.50)
	-2 to -3	17(12.14)	12(12)	6(15)
	>-2 to 1	93(66.43)	68(68)	23(57.50)
	>1	2(1.43)	1(1)	0
Total	140	100	40	

% of the total subjects ≤5 years old had severe wasting (severe acute malnutrition), while 21.33% had moderate wasting. Severe and moderate stunting was observed in 5% and 24% of the total subjects. In children > 5 years, 20% had severe thinness, 12.14% had thinness and only 2 (1.43%) case was overweight.

16% of the vaccinated subjects ≤5 years old had severe wasting (severe acute malnutrition), while 21% had moderate wasting. Severe and moderate stunting was observed in 8.70% and 15.63% of the total subjects. In children > 5 years, 19% had severe thinness, 12% had thinness and only 1(1%) case was overweight.

Amongst unvaccinated subjects, 16.67% children ≤ 5 years old had severe wasting (severe acute malnutrition), while 21.67% had moderate wasting. Severe and moderate stunting was observed in 5.88% and 30% of the total subjects. In children > 5 years, 27.50% had severe thinness, 15% had thinness and no case was overweight.

Table 4: Relationship of measles antibody status with nutritional status of total subjects

Parameter of nutritional status		Antibody status			Total	P value
		Positive N (%)	Negative N (%)	Equivocal N (%)		
Weight for Age z score	<-3	24(44.44)	24(44.44)	6(11.11)	54	0.055
	-2 to-3	39(60.94)	19(29.69)	6(9.37)	64	
	>-2	245(67.68)	67(18.50)	50(13.81)	182	
Height for Age Z score	<-3	7(46.67)	6(40)	2(13.33)	15	0.019
	-2 to-3	44(54.32)	31(38.27)	6(7.40)	81	
	>-2	147(71.36)	50(24.27)	9(4.37)	206	
Weight for Height z score	<-3	12(52.17)	9(39.13)	2(8.70)	23	0.55
	-2 to-3	17(53.13)	10(31.25)	5(15.63)	32	
	>-2	54(51.43)	44(41.90)	7(6.67)	105	
BMI Z score (age \geq 5yrs)	<-3	16(57.14)	9(32.14)	3(10.71)	28	0.001
	-2 to-3	10(58.82)	6(35.29)	1(5.88)	17	
	>-2 to 1	80(86.02)	9(9.67)	3(3.22)	93	
	>1	2(100)	0	0	2	

Table 5: Nutritional status wise geometric mean titer (GMT) of measles specific igg antibody of total children

Parameter of nutritional status		GMT (mIU/mL)	P value
W/A z score	> -2SD	889	0.001
	<-2SD to-3SD	687	
	<-3SD	548	
H/A z score	-2SD	1832	0.005
	-2SD to-3SD	685	
	<-3SD	387	
W/H z score	> -2SD	625	0.47
	-2SD to-3SD	587	
	<-3SD	549	

Discussion

Measles vaccination triggers both a cellular and a humoral immune response. Following the activation of T-lymphocytes, B-cells produce measles-specific antibodies. The specific level of immunoglobulin is an indicator of the immune response. Protein energy malnutrition can be a contributory factor for immunodeficiency, thereby diminishing the immunological response to the vaccine. Though many studies reported normal antibody response to measles vaccination in malnourished children, however, specific antibody response was suppressed in severe cases of PEM.

Our study also gives conflicting results. We have taken four parameters to study nutritional status: weight for age, height for age, weight for height and BMI. A highly statistically significant relationship ($p=0.001$) was observed between BMI Z scores (in subjects aged ≥ 5 yrs) and seropositivity, with higher seropositivity being noted in children with higher BMI z scores.

Similarly, a statistically significant relationship ($p=0.019$) was observed between height for age Z scores and seropositivity, with higher seropositivity being noted in children with higher height for age z scores. However, no significant relationship was observed between seropositivity and either weight for height z scores or weight for age z scores. According to weight for age, in well-nourished children seropositivity was 67.68% with GMT 889 mIU/ml, moderately malnourished (wasting) 68% with GMT 687 mIU/ml and in severely malnourished (severe wasting) 60.90% with GMT 548 mIU/ml. It was not found statistically significant with p value 0.05 for seropositivity but significant with p value 0.0001 for antibody levels. According to Height for age in well- nourished children seropositivity was 54.32% with GMT 685 mIU/ml, moderately malnourished (stunting) seropositivity was 71.36% with GMT 387 mIU/ml and in severely malnourished (severe stunting) seropositivity 46.67% with GMT 361 mIU/ml. P value noted was 0.005 which is significant. In weight for length/height both seropositivity and GMT were found insignificant in well-nourished, moderately malnourished as well as severely malnourished subject (P value: 0.47)

In a study in Nigeria by Ifekwunigwe et al.²⁴, the geometric mean titer in subjects whose nutritional status was normal (>90% of median weight for age), mildly (75 to 90%), moderately (60 to 75%), or severely (<60%) malnourished were 7.5, 8.8, 7.9, and 7.9, respectively. So, malnutrition did not affect the children ability to develop adequate immune response to measles. In another study by Dao et al²⁵, seroconversion was not associated with anthropometric indices. McMurray et al²⁶ found that the children's nutritional status had no effect after vaccination. All the children have equal immunological response with respect to nutritional status. Mean hemagglutination-inhibition titres are slightly reduced in all nutritional groups 14 months after vaccination. Smedman et al²⁷, Halsey et al²⁸, Ekunwe et al²⁹ found good antibody response in children which were not severely malnourished. Similarly Lyamuya et al³⁰ found there were no significant differences in measles antibody levels with regard to variations in nutritional status. Our study is not only showing antibody response in moderately nourished children but also in severely nourished children. Some studies reported seroconversion rates at least as high in malnourished as in well-nourished children because it is cell mediated immunity that is suppressed not the humoral immunity.^{31,32}

Delayed antibody response to measles vaccine was seen in malnourished children.³³ Similar to our study, there was one study which demonstrated that stunting is associated with low antibody response.³⁴ In the same study, apart from severe stunting, severe wasting was also associated with lower antibody response, an observation which was not observed in our study. Idris et al³⁵ found decreased antibody titre in children with Kwashiorkar. Hafez et al³⁶ found decrease humoral response to measles vaccine.

So, it was seen that malnourished children in the community can be safely and effectively vaccinated against measles. But some studies showing good antibody response and some showing poor antibody response. The mechanisms behind the immunological response are still inadequately understood. More researches are needed in this field to come to any conclusion.

Conclusion

Nutritional status of children has an association with measles antibody titres as well GMT of measles specific IgG antibody, with those with better nutritional status having higher measles antibody titres.

Reference

1. Lee MS et al. Epidemiology of measles in Taiwan: dynamics of transmission and timeliness of reporting during an epidemic in 1988–9. *Epidemiology and Infection*, 1995, 114: 345–359.

2. Lee MS et al. Seroepidemiology and evaluation of passive surveillance during 1988–1989 measles outbreak in Taiwan. *International Journal of Epidemiology*, 1992, 21: 1165–1174.
3. Department of Health (Taiwan, China). Proceedings of the national meeting for measles, poliomyelitis, congenital rubella and neonatal tetanus elimination in Taiwan. Taipei, April 1995 (in Chinese).
4. Department of Health (Taiwan, China). Proceedings of the national annual meeting for infectious diseases control. Taipei, 1996 (in Chinese).
5. Department of Health (Taiwan, China). Cases of notifiable and reportable diseases, Taiwan-Fukien area. *Epidemiology Bulletin*, 1999, 15: 11.
6. Lee MS et al. Post mass-immunization measles outbreak in Taoyuan County, Taiwan: dynamics of transmission, vaccine effectiveness and herd immunity. *International Journal of Infectious Disease*, 1999, 3: 64–69.
7. Liu CC, Lei HY, Chiang YP. Seroepidemiology of measles in southern Taiwan: two years after implementation of the measles elimination program. *Journal of the Formosa Medical Association*, 1996, 95: 37–40.
8. Chiu HH et al. Seroepidemiological study of measles after the 1992 nation-wide MMR revaccination program in Taiwan. *Journal of Medical Virology*, 1997, 51: 32–35.
9. Lee MS et al. An investigation of measles elimination in Taiwan by 2000. Paper presented at: The 8th International Congress for Infectious Diseases. Boston, USA, May 1998: 166.
10. Narain JP, Banerjee KB. Measles in India: epidemiology and control. *The Indian Journal of Pediatrics*. 1989 Jul 1;56(4):463-72.
11. Measles [Internet]. World Health Organization. 2016 [cited 11 September 2014]. Available from: <http://www.who.int/mediacentre/factsheets/fs286/en/>. [Internet]. 2016 [cited 11 August 2013]. Available from: [http://www.unicef.org/india/Uttar_Pradesh_Fact_Sheet.p df](http://www.unicef.org/india/Uttar_Pradesh_Fact_Sheet.pdf).
12. World Health Organization. Measles vaccines: WHO position paper – April 2017. *Wkly Epidemiol Rec* 2017;17(92):205–28.
13. Centers for Disease Control and Prevention. Progress toward measles elimination-African Region, 2013–2016. *Morb Mortal Wkly Rep (MMWR)* 2017;66(17):436–43.
14. Doshi R et al. The effect of immunization on measles incidence in the Democratic Republic of Congo: results from a model of surveillance data. *Vaccine* 2015;33(48):6786–92.
15. Cutts F, Grabowsky M, Markowitz L. The effect of dose and strain of live attenuated measles vaccines on serological responses in young infants. *Biologicals* 1995;23(1):95–106.
16. Leuridan E et al. Early waning of maternal measles antibodies in era of measles elimination: longitudinal study. *Bmj* 2010;340:c1626.
17. Doshi R et al. Field evaluation of measles vaccine effectiveness among children in the Democratic Republic of Congo. *Vaccine* 2015;33(29):3407–14.
18. Adu FD et al. Low seroconversion rates to measles vaccine among children in Nigeria. *Bull World Health Organ* 1992;70(4):457–60.
19. Carney JM et al. Cell-mediated immune defects and infection: a study of malnourished hospitalized children. *Am J Dis Children* 1980;134(9):824–7.
20. Neumann CG et al. Immunologic responses in malnourished children. *Am J Clin Nutr* 1975;28(2):89–104.
21. Mancini S et al. Description of a large measles epidemic in Democratic Republic of Congo, 2010–2013. *Confl Health* 2014;8(9)
22. Gerard S, Kyrousis E, Zachariah R. Measles in the Democratic Republic of Congo: an urgent wake-up call to adapt vaccination implementation strategies. *Public Health Action* 2014;4(1):6–8.

23. Cogill B. Anthropometric indicators measurement guide. Revised 2003.
24. Ifekwunigwe AE, Grasset N, Glass R, Foster S. Immune responses to measles and smallpox vaccinations in malnourished children. *The American journal of clinical nutrition*. 1980 Mar 1; 33(3):621-4.
25. Dao H, Delisle H, Fournier P. Anthropometric status, serum prealbumin level and immune response to measles vaccination in Mali children. *Journal of tropical pediatrics*. 1992 Aug 1; 38(4):179-84.
26. McMurray DN, Rey H, Casazza LJ, Watson RR. Effect of moderate malnutrition on concentrations of immunoglobulins and enzymes in tears and saliva of young Colombian children. *The American journal of clinical nutrition*. 1977 Dec 1;30(12):1944-8.
27. Smedman L, Silva MC, Gunnlaugsson G, Norrby E, Zetterstrom R. Augmented antibody response to live attenuated measles vaccine in children with *Plasmodium falciparum* parasitaemia. *Annals of tropical paediatrics*. 1986 Jun; 6(2):149-53.
28. Halsey NA, Boulos R, Mode F, Andre J, Bowman L, Yaeger RG, Toureau S, Rohde J, Boulos C. Response to measles vaccine in Haitian infants 6 to 12 months old: influence of maternal antibodies, malnutrition, and concurrent illnesses. *New England journal of medicine*. 1985 Aug 29;313(9):544-9.
29. Ekunwe EO. Malnutrition and seroconversion following measles immunization. *Journal of tropical pediatrics*. 1985 Dec 1;31(6):290-1.
30. Lyamuya EF, Matee MI, Aaby P, Scheutz F. Serum levels of measles IgG antibody activity in children under 5 years in Dar-es-Salaam, Tanzania. *Annals of Tropical Paediatrics: International Child Health*. 1999 Jun 1;19(2):175-83.
31. Bhaskaram P, Madhusudhan J, Radhakrishna KV, Reddy V. Immune response in malnourished children with measles. *Journal of tropical pediatrics*. 1986 Jun 1;32(3):123-6.
32. Kimati VP, Loretu K, Munube GM, Kimboi F. The problem of measles virus response with reference to vaccine viability, age, protein energy malnutrition and malaria in the tropics. *Journal of tropical pediatrics*. 1981;27(4):205-9.
33. Powell GM. Response to live attenuated measles vaccine in children with severe kwashiorkor. *Annals of tropical paediatrics*. 1982 Sep;2(3):143-5.
34. Waibale P, Bowlin SJ, Mortimer EA, Whalen C. The effect of human immunodeficiency virus-1 infection and stunting on measles immunoglobulin-G levels in children vaccinated against measles in Uganda. *International journal of epidemiology*. 1999 Apr 1;28(2):341-6.
35. Idris S, El Seed AM. Measles vaccination in severely malnourished Sudanese children. *Annals of tropical paediatrics*. 1983 Jun;3(2):63-7.
36. Hafez M, Aref GH, Mehareb SW, Kassem AS, El-Tahhan H, Rizk Z, Mahfouz R, Saad K. Antibody production and complement system in protein energy malnutrition. *The Journal of tropical medicine and hygiene*. 1977 Feb;80(2):36-9.

Received: 10-05-2021 // Revised: 30-05-2021 // Accepted: 15-06-2021