

ORIGINAL RESEARCH

A Comparative Study of Serum Creatinine, Serum Uric Acid and Blood Urea in Normal Pregnant and Pregnancy Induced Hypertensive Subject

Pawan Kumar Saini¹, MD Shahbaz Alam², Piyush Kumar³, Subhra Srivastava⁴

¹Assistant professor, Department of Surgery, VIMS, Gajraula, Uttar Pradesh, India.

²Associate Professor, Department of Anaesthesia, Teerthankar Mahaveer Medical College and Research Centre, Moradabad, Uttar Pradesh, India

³Assistant Professor, Department of Surgery, Teerthankar Mahaveer Medical College and Research Centre, Moradabad, Uttar Pradesh, India

⁴Assistant Professor, Department of Obstetrics and Gynaecology, Teerthankar Mahaveer Medical College and Research Centre, Moradabad, Uttar Pradesh, India

ABSTRACT

Background: World Health Organization reported that there were 14.1 million new cancer cases, 8.2 million cancer deaths and 32.6 million people living with cancer within 5 years of diagnosis worldwide.

Materials and Methods: Data were collected on the bases of findings of history and clinical examination. Fine needle aspiration cytology (FNAC) was done in patients with palpable lump in breast and suspicious lesions. Suspected cases were sent for histo pathological examination.

Results: Most of the cancer patients were found in the 51–55-year age group. In this study, cyto-pathological features were occur, Fibroadenoma (54%), fibroadenosis(8%), Fibrocystic disease(4%), seb. Cyst breast (2%), gynaecomastia(0%), breast abscess(12%), mastitis(0%), ductal carcinoma(18%), paget disease of nipple(2%).

Conclusion: Benign breast lesions are more common among female population than malignant lesion, the frequency of breast cancer is increasing rapidly across the global.

Keywords: Fibroadenoma, Breast Lumps, Paget Disease, Mastitis.

Corresponding Author:Dr. Subhra Srivastava, Assistant Professor, Department of Obstetrics and Gynaecology, Teerthankar Mahaveer Medical College and Research Centre, Moradabad, Uttar Pradesh, India

INTRODUCTION

Breast cancer is the most commonly diagnosed cancer in women worldwide. In developing countries, it is the leading cause of death in women. The general approach to evaluate breast cancer is clinical examination, imaging techniques usually mammography, ultrasonography, or both, and histopathological assessment. Increased public awareness and improved screening are very important for earlier diagnosis and led to improved survival rates for women diagnosed with breast cancer.

Breast cancer is one of the most prevalent cancers in women. Due to increase in life expectancy, urbanization and adoption of western lifestyle, the cases of breast cancer is increasing in the developing country. World Health Organization reported that there were 14.1 million new cancer cases, 8.2 million cancer deaths and 32.6 million people living with cancer within 5 years of diagnosis worldwide.^[1] A number of females visit the clinic with the common presentation of breast lump.^[2]

The major symptoms of breast lump are swellings, protuberance, bulge or bump in the breast which feels different from the breast tissue in the same area of the other breast. A breast lump

can be either benign or malignant. Though, most of the breast lumps are benign. As breast cancer is commonly presenting with a lump, so diagnosis of benign from malignant lesions is important. Worldwide, awareness is created about the significant role of breast lump in breast cancer. Fibro adenoma, fibrocystic disease, abscess, tumor and fat necrosis or malignant breast lump are the most common breast lumps. Most of the cases are related to burst lump which are presented to surgery departments. Breast cancer patients are always in a state of fear until they have undergone specialist assessment, investigations and eventual reassurance. A number of patients referred to a surgery department is said to have benign disease.^[3-5] The most common of the benign breast diseases is fibroadenoma is.^[6-9] The main aim of this present study is to audit the cyto-pathologic features of patients with breast lump hospital in India.

MATERIALS & METHODS

Study Population: Fifty cases included in this study which were attended with breast symptomatology.

Data collection: Data were collected on the bases of findings of history and clinical examination. Fine needle aspiration cytology (FNAC) was done in patients with palpable lump in breast and suspicious lesions. Suspected cases were sent for histo pathological examination.

Data analysis: Data were analyzed by using Microsoft excel.

RESULTS




Figure 1: Distribution of patient according to stage

The present study consisted of 18% cases of malignant out of 50 cases. The reason behind such finding is due to secondary hospital data and breast malignant cases are referred to medical college hospital from large surrounding rural, suburban and urban population. Out of 50 cases, 35 belonged to benign lesions and 9 malignant and 6 were of inflammatory pathology. The ratio of benign to malignant breast disease was calculated as 4:1. Fibroadenoma 27 was the most common benign lesion. Most of the cancer patients were found in the 51–55-year age group. In this study, cyto-pathological features were occur,

Fibroadenoma (54%), fibroadenosis (8%), Fibrocystic disease (4%), Seb. Cyst breast (2%), gynaecomastia(0%), breast abscess(12%), mastitis(0%), ductal carcinoma(18%), paget disease of nipple(2%).




Figure 2: Distribution of patient according to age




Figure 3: Distribution of patient according to Cytopathology

DISCUSSION

The youngest and the oldest patient included in the study were 20 years old and 60 years old respectively. Most of the breast cancer patients were found 51–55-year age group cases.

Benign conditions of breast cases are significantly higher than malignant conditions. M. Kumar et al,^[10] observed that in Indian rural population the benign breast diseases are 5 to 10 times more common than breast cancers. Similar results were found in Aisha Memon et al,^[11] study. In the present study, cases of benign breast lesions are 4 times more than cancerous lesions. It has been revealed by M. Kumar et al,^[10] that incidences of benign breast diseases are common in developing countries but due to lack of awareness women disregard the breast lump. The delay in both benign and malignant lesions is associated with illiteracy, social taboo, unawareness of patients. Such delay in malignant lesions leads to poor prognosis. In one of their study Aisha Memon et al,^[11] found 58.8% benign cases in 500 cases. While in an another study Adesunkanmi et al,^[12] in Nigeria revealed that 87.2% patients had benign breast lumps. The present study found 82% cases of benign lesions (including 13% inflammatory cases). Therefore, our findings were also supported by the findings of Adesunkanmi et al,^[12] Vissa Shanthi et al,^[13] studies. Pradhan et al,^[14] conducted a study in Nepal and observed that upto 15.5% cases were malignant. In another study Mayun et al,^[15] reported that malignant lesions were diagnosed approximately 40%. This study found 9 (18%) cases malignant and 1(2%) case of paget's disease. Mortality and incidence are relatively lower in developing countries and other parts of globe in comparison to western population.^[16]

CONCLUSION

Although benign breast lesions are more common among female population than malignant lesion, the frequency of breast cancer is increasing rapidly across the global. It is important to screen females at a younger age to detect early breast cancer. Internationally mass awareness should be created regarding detection of early breast cancer and to foster knowledge about the medical and socio economic implications of a common public health issue.

REFERENCES

1. WHO-Globocan 2012: Estimated cancer incidence, mortality and prevalence.
2. Chaudry I. qureshi s, Rasul, Aquela B. Pattern of benign breast disease. JSP, volume 8 (3), July Spetember; 5-7.
3. Osime OC Ohanaka EC Analysis of five-year breast biopsies carried out in the University of Benin Teaching Hospital Benin City. Niger PostgradMed J. 2008 ;15(3):160-163.
4. Ohene-Yeboah MO. An audit of excised breast lumps in Ghanaian women. West Afr J Med. 2005;24(3):252-255.
5. Clegg-Lampsey JN, Aduful HK, Yarney J, Adu- Aryee NA, Vanderpuye V, Kyereh M, et al. Profile of breast disease at a self-referral clinic in Ghana.NepalMedColl J. 2004;6(2):129-132.
6. Oluwole SF, Freeman HP. Analysis of benign breast lesions in blacks. Am J Surg. 1979 Jun;137(6):786-789.
7. Sönmez K, Türkyilmaz Z, Karabulut R, Demirođullari B, Ozen IO, Moraliöđlu S, Surgical breast lesions in adolescent patients and a review of the literature. Acta Chir Belg. 2006 Jul-Aug;106(4):400-404.
8. Adeniji KA, Adelusola KA, Odesanmi WO. Benign disease of the breast in Ile-Ife: a 10 year experience and literature review. Cent Afr J Med. 1997;43(5):140-143.
9. Ihekwa FN. Benign breast disease in Nigerian women: a study of 657 patients. J R Coll Surg Edinb. 1994;39(5):280-283.
10. M Kumar, K Ray, S Harode, DD Wagh.The Pattern of Benign Breast Diseases in Rural Hospital in India, East and Central African Journal of Surgery 2010;15:2.

11. Aisha Memon, Shahida Parveen, A.K. Sangrarasi, Arshad M. Malik, Aziz Laghari, K. Altaf Hussain Talpur. Changing Pattern of Benign Breast Lumps in Young Females. *World Journal of Medical Sciences* 2007;2:21-24.
12. Adesunkanmi, A.R., E.A. Agbakwuru. BBD at Wesley guild hospital Ilesha Nigeria. *West Afr. J. Med* 2001;20:146-51.
13. VissaShanthi, Kashmir Ali, Nandam Mohan Rao, Baddukonda Appala Rama Krishna, Kuppali Venkata Muralimohan. Clinicopathological study of breast lesions in females with assessment of correlation between tumor grade and prognostic factors. *J Biosci Tech* 2011; 2:367-378.
14. Pradhan M, Dhakal HP. Study of breast lumps of 2,246 cases by fine needle aspiration. *J Nepal Med Assoc* 2008;47:205-9.
15. Mayun AA, Pindiga UH, Babayo UD. Pattern of histopathological diagnosis of breast lesions in Gombe, Nigeria. *Niger J Med* 2008;17:159-62.
16. Khan S, et al. Prospective study of pattern of breast diseases at Nepalgunj medical college (NGMC). *Nepal Kathmandu Univ Med J* 2003;1:95-100.