

Assessment Knowledge regarding Vitamin D deficiency among female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021

Saud AbdulhaiGari¹, Wesaam Mohammad Jamal Ghafuri², Abeer Mohammed Noor Hosawi³, Mohammed Abed Gurunfula⁴, Makkiyyah Sultan Mohammed Aljayzani⁵, EffatMagbel Al Mawalad⁶, Rasha Naser Albeshri⁷, Jamilah HamdHmad Algurashi⁵, Azzah Zamil Alzaid Alsharif⁸, Roqaiyah Mohammad Bakur Hawsawi⁹, Nasr Mohmmad Maroof¹⁰, Jarallah Hassan Algamdy¹¹

¹General dental practitioner, Kudi-Alhjera primary health care, Makkah, Saudi Arabia.

²Manager of infection control administration, public health-general directorate of health affairs at Makkah region

³Specialist Nursing, Health Affairs in Riyadh, Saudi Arabia.

⁴Epidemiology technician, Al Eskan primary health care, Makkah, Saudi Arabia.

⁵Nursing Technician, BathaQuresh, Makkah, Saudi Arabia.

⁶Nursing Technician, Alhndawiay, Makkah, Saudi Arabia.

⁷Nursing Technician, Hadda Primary Health Care Centre, Makkah, Saudi Arabia.

⁸Nursing Technician, Shmeisi Medical Complex, Makkah, Saudi Arabia.

⁹Nursing Aid, AlthagahHospital, Jeddah, Saudi Arabia.

¹⁰Nursing technician, Hindawiya health center, Makkah, Saudi Arabia.

¹¹Nursing Technician, AL Rashdia Primary Health Care Center, Makkah, Saudi Arabia.

Abstract

Background :

Vitamin D deficiency is prevailing in Saudi Arabia. Recent national data indicated an inverse association between vitamin D status and coronary heart disease (CHD), which increases concerns about vitamin D deficiency as a serious public health problem. Therefore, the current study aimed to Assessment Knowledge regarding Vitamin D deficiency among female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021. Saudi Arabia has one of the highest reported incidence of vitamin D deficiency in studies conducted worldwide. However, there has been very limited exploration of vitamin D related knowledge, attitudes and practices among female adult in Saudi Arabia. However, the case of vitamin deficiency is reported to increase in many female Saudi Arabia worldwide, especially in Saudi Arabia. Mothers, having adequate knowledge about vitamin D and its supplementation, may lead to possible lowering of the cases of vitamin D deficiency and its complications. **Ami of the study:** To assessment Knowledge regarding Vitamin D deficiency among female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021 **Methods:** This was an analytical cross-sectional study conducted during 2021, among female adult attending the Primary Health clinic at Makkah AL-Mokarramah random sampling technique was applied and an Arabic self-administered questionnaire used for data collection. **Result:** shows that most of the participants (37.0%) were in the age 25-30 years also regarding level of education the majority of participant are Secondary education were (34.0%). regarding Marital status the majority of participant are married were (58.0%). Regarding the Economic level the majority of participant Average were (45.0%). Regarding the Chronic

disease the majority of participant No.were(72.0%). **Conclusion:**The low levels of knowledge about vitamin D and low consumption of vitamin supplementation, including vitamin D, calcium, multivitamin, and calcium supplements with vitamin D, may have contributed to the higher prevalence of vitamin D deficiency among the female adult. Further studies using a qualitative approach are crucial to explore the underlying reasons for low knowledge about vitamin D and behaviors related to vitamin D including the intake of vitamin supplementation that may contribute to the high burden of vitamin D deficiency in the country.

Keywords:Assessment, Knowledge, Vitamin D,deficiency, adult, attending,Primary Health clinic, Makkah

Introduction

Recent evidence has indicated that vitamin D deficiency and insufficiency are becoming global epidemics. Studies conducted in Western countries have shown that vitamin D deficiency was present in 20% -25% of the total population . In the Middle East region, approximately 60%–65% of the population was affected . Vitamin D deficiency also has a significant presence in Saudi Arabia, even though there is plentiful sunlight throughout the year. The majority of studies that have measured vitamin D levels in Saudi Arabia have indicated a high prevalence of vitamin D deficiency among different population groups . A recent national survey showed that almost 40% of males and 60% of females in Saudi Arabia had vitamin D deficiency. Vitamin D deficiency is highly prevalent in sunny countries such as Saudi Arabia. Previous studies within the country indicated a high rate of vitamin D deficiency among both males and females [1,2]

A recent national survey reported high rates of vitamin D deficiency: 40% in males and 60% in females in Saudi Arabia [3]. Vitamin D deficiency has historically been known to increase the risk of somemusculoskeletal disorders, such as osteoporosis [4]. Furthermore, several studies have shown anThere is an increasing bodyofevidence that suggests vitamin D deficiency is becoming a global epidemic, with South Asians being one of the most affected population groups [5]. Exposure to sunlight is the main source of vitamin D. Some of the best dietary sources of vitamin D include oily fish and egg yolks, as well as vitamin D dietary supplements [6]. Despite receiving plentiful sunlight throughout the year in their countries, South Asians remain one of the most vulnerable populations to suffer from low vitamin D levels [7]. A number of studies that have measured vitamin D levels among South Asian population groups have indicated a high prevalence of its deficiency among healthy young adults, women and infants [8]. Aside from the classical role of vitamin D in bone health and the regulation of calcium and bone homeostasis, several large observational studies worldwide have shown an association between vitamin D deficiency and the risk of coronary heart disease (CHD) and associated risk factors such as hypertension and diabetes [9]. Furthermore, recent meta-analyses of observational studies also reported significant associations of vitamin D deficiency with cardiovascular disease (CVD) mortality [10], and the increased risk of CVD [11]. Thus, the existing literature of observational studies indicated an association between vitamin D deficiency and the risk of CHD. Nevertheless, to date only a few

randomized controlled trials (RCTs) have been conducted to examine the effect of vitamin D supplementation on reducing the risk of CHD [12]

Several studies have focused on the biological factors that cause vitamin D deficiency around the world; however, few studies have examined the effects of cultural and lifestyle behaviors, knowledge and beliefs that influence vitamin D status [13]. Recently, we conducted a study using a quantitative approach to examine knowledge and attitudes about, and behaviors toward, vitamin D in subjects with and without CHD in Saudi Arabia [14]. The results showed that low levels of knowledge about vitamin D and low consumption of vitamin supplements, including multivitamins and vitamin D supplements, were associated with vitamin D deficiency [15]

Literature Review

Tariq et al. BMC Public Health (2020) Despite this female adult having access to the internet and health information, knowledge about vitamin D was quite poor. Poor knowledge could be due to conflicting information available about vitamin D and lack of clear localised messages about sun exposure. [16]

Similar limitations to the knowledge of vitamin D have been reported in studies across Bangladesh, the Middle East, Iran, Lebanon and also among immigrants from South Asian countries living across Europe. Similar to our study, most studies have found their participants not able to identify sources of vitamin D in food and sunlight as critical enabler in vitamin D production [17, 18]

However, results are in contrast to those reported in the UK where the participants did demonstrate good level of knowledge about vitamin D [19]. This perhaps is indicative of the increased public awareness and food fortification practices in the developed world. Multi-prong strategies are required to appraise public knowledge and awareness about vitamin D.

Study findings demonstrated that vitamin D deficiency was significantly more prevalent in the female /+adult than the controls. Previous studies have reported similar results [20, 21]

[22] Reported that People should receive information, in local languages, that reflects the actual state of knowledge regarding vitamin D and its association with health, along with clear information on vitamin D sources [22]. These national public health messages should be relayed across various platforms (i.e., social media, health centres, universities, schools), especially through the media, to increase awareness among all sub-groups of the population [23]. The traditional knowledge, attitudes, and practice (KAP) survey theory suggests a direct linear relationship between knowledge, attitudes, and behaviors, which is, according to several studies, very simple and not true. This is because people's behaviors have a multifactorial nature and depend on many factors such as socio-cultural and environmental factors, not just knowledge and attitudes [24]. Further it is important to recognise awareness regarding the causes and prevention of vitamin D deficiency as an avenue to be explored by the public health offices, which should begin by conducting conclusive studies to determine general public knowledge regarding this pertinent issue [22]

A study among older adults in Netherlands has reported similar results as the higher levels of knowledge about vitamin D was associated with higher vitamin D serum levels [27]

Food Fortification Programme (FFP) in KSA is exploring the food fortification of staple foods like milk, vegetable oil and wheat with vitamin D [25] This is in alignment with evidence that suggests universal vitamin D food fortification can improve serum 25-hydroxyvitamin D levels [22]. However, to encourage public adaptation, it is important to address any public concerns about food fortification, which have been reported to exist especially among South Asians [12].

Rational:

Vitamin D deficiency is widespread in Saudi Arabia However, there has been very limited exploration of vitamin D related knowledge, attitudes and practices among female adult in Saudi Arabia. Vitamin D is known to have essential roles in the human body. However, the case of vitamin deficiency is reported to increase in many female Saudi Arabia worldwide, especially in Saudi Arabia. Mothers, having adequate knowledge about vitamin D and its supplementation, may lead to possible lowering of the cases of vitamin D deficiency and its complications. The researcher has an interest in vitamin D in female adult PHCCs are the appropriate location for study the when the researcher was working in the PHCC, she noticed that a lot of mothers who came for vaccinating their children have no idea or proper information about the vitamin D and a lot of mother's consultations about vitamin D

Ami of the study: To assessment Knowledge regarding Vitamin D deficiency among female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021

Objectives:

To assessment Knowledge regarding Vitamin D deficiency among female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021

Methodology

Study Design: Cross-sectional analytical study.

Study Population: female adult attending the Primary Health clinic at Makkah, Saudi Arabia, 2021

Eligibility Criteria:

Inclusion Criteria

- Female gender.
- All nationalities.
- Mothers attended PHCCs in Makkah Al-Mokarramah.

Exclusion Criteria

- Illiterates mothers.
- Mothers can't speak Arabic language.

Study Area:

Makkah Al-Mokarramah is the holy city for all Muslims. It is the place where the Prophet Muhammad (peace be upon him), was born, and where the message of God was first revealed to him and where the presence of Almasjid Alharam

(the holy mosque) which contains Alkaa'ba representing the Qibla, the guidance for all Muslims in their prayers.

Makkah Al-Mokarramah is located in the western region in an area called Makkah region and contains 85 PHCC which divided under seven PHCC sectors (13 PHCC under Aladel sector, 11 PHCC under Alkakah sector, 13 PHCC under Asharaie sector, 14 PHCC under Azzaher sector, 10 PHCC under Kheles sector, 11 PHCC under Alkamel sector and 13 PHCC under Aljamoom sector).

There are several services in the PHCC in Makkah Al-Mokarramah including family medicine clinics, general clinics, chronic disease clinics, antenatal clinics, dental clinics, well baby clinics and vaccination clinics, as well as pharmacy, radiology and laboratory services. Family medicine clinics run by certified specialists and consultants while the other remaining clinics run by the general practitioners.

Total population in Makkah Al-Mokarramah city in the last statistics at 2020 is 2,578,722 according to General Authority of Statistic (GAS) in the Kingdom of Saudi Arabia (KSA).

Sample Size:

According to the GAS of KSA, the total population in Makkah Al-Mokarramah city in 2020 is 2,578,722. According to statistics of Ministry of Health (MOH), the percentage of the children who are their age is 0-5 years is 8.4%, then accordingly the total population of this study calculated and it was 194,143.

The sample size was calculated by Raosoft online website, based on the following assumptions: the total population was 194,143, the prevalence of knowledge about weaning was 50%, confidence Interval was 95%, and error margin was 5%. Accordingly, the estimated sample size was (300). It was $384 + 38.4$ (10%) for non-responders = 422.4 and by approximation it became (300).

Sampling Technique:

All areas of Makkah Al-Mokarramah city and its environs is divided under seven sectors of PHCCs were legible residents registered by Directorate General of Health Affairs in Makkah Al-Mokarramah region. From each sector, one PHCC (Stratified) was randomly selected to represent its equivalent portion (1/7) in the study sample, then simple randomization was adopted on numbers using online random number generator, to select one PHCC from the list of all names specific to each sector. Systematic random sampling technique was used to invite 55, 67, 65, 65, 50, 65 and 55 mothers from Aliskan PHCC, Azzaher PHCC, Asharaie PHCC, Alawali PHCC, Um Algerm PHCC, HadaAlsham PHCC, and Alkamel PHCC, respectively to participate in the study.

The total eligible mothers attended the randomly selected PHCCs By dividing the total eligible mothers per day to the required sample ($30/10 = 3$), therefore, every third eligible mother was invited to participate in the study. The second eligible mother was selected by simple random as a starting point, and then every third mother (2nd, 5th, 8th, 11thetc.) were invited. In the case of

refusing to participate in the study the next eligible mother was invited, and then the selection was continued in the same interval of every third eligible mother.

DataCollection Tool:

An Arabic self-administered questionnaire was used for collecting the data. The questionnaire modified and translated from a validated english interview schedule. The questionnaire collected data on written consent for participation, socio-demographic details, previous information about the weaning and its source, the knowledge about weaning (e.g. age of introduction of solid food, amount, and types of solid food), as well as the weaning practice (e.g. when the solid food began, which amount and types of food used).

DataCollectionTechnique:

After getting all needed permissions, the researcher assigned one week for each randomly selected PHCC and the required sample divided into five portions to be collected through five days. The researcher took place in the reception of the PHCC beside the receptionist, and when the patients came to take a paper for getting a service in the PHCC, the mothers with the eligible criteria was invited to participate in the study by asking three questions (do you have a 5-years child or younger, can you read and write and verbal consent for participation).

At first, the second eligible mother was selected by simple random as a starting point, and then every third mother (2nd, 5th, 8th, 11thetc.) were invited. In the case of refusing to participate in the study the next eligible mother was invited, and then the selection was continued in the same interval of every third eligible mother.

An arabic, self-administered questionnaire was distributed to the invited mothers with a pen to fill the questionnaire while they were sitting in the waiting area before getting the service. The mothers took 5-10 minutes to fulfill the questionnaire, and after they finished, they placed the questionnaire in a small basket presented in a table in the waiting area as requested by the researcher. When the researcher noticed that, she came to collect the filled questionnaire and thank each mother for her participation and gave her a brochure about proper weaning process and its importance and risks of inappropriate weaning with a candy for her and her child if accompanied by her.

Data Entry and Analysis:

All collected data verified by hand, then coded before its entry to a personal computer. Statistical Package for the Social Sciences (SPSS) version 24 used for data entry and analysis. Data were presented using descriptive statistics in the form of frequencies and percentages for qualitative categorical variables, and means and standard deviations for quantitative variables. Analytic statistics using the association between two variables were applied. P-value less than 0.05 was considered statistically significant.

Pilot Study:

The pilot study conducted on 10% of the sample size with similar characteristics as the study's sample, but who were not included in the final

sample using the similar questionnaire. The sample of the pilot chosen by non-probability convenient sampling technique in Al-Eskan PHCC which included under Al-Kaakiah PHCC sector in Makkah Al-Mokarramah. It aims to test the feasibility of the study, the clarity of the questionnaire and estimating the time needed to fill it. Important suggestions, omissions, and corrections from the results of the pilot study were incorporated in the final questionnaire and therefore enhanced its reliability.

Ethical Considerations:

- Approval from the research committee was obtained.
- Approval from Makkah joint program of family and community medicine was obtained.
- Approval from General Administration of Research and Studies, MOH, KSA.
- Approval from the Directorate General of Health Affairs of Makkah Al-Mokarramah, public health, and PHCC affairs sector was obtained.
- Written consent from each participant in the study was obtained for data collection.
- All information kept confidential.
- The researcher acknowledged the supervisor, advisors, helpers, facilitators, participants and family members.
- Brochure about weaning practice was provided for each participant in the study.

Budget:

The study was self-funded.

Results

Table 1 Distribution of demographic data (age, gender, social, Smoking, HYN duration, Occupation) in our study (n=200)

	N	%
Age		
<25	56	28
25-30	74	37
30-35	38	19
>35	32	16
Level of education		
Primary	24	12
Intermediate	56	28
Secondary	66	33
High education	54	27
Marital status		
Married	170	85

Unmarried	30	15
Economic level		
Low	74	37
Average	90	45
High	36	18
Chronic disease		
Yes	56	28
No	144	72

Table 1 shows that most of the participants (37.0%) were in the age 25-30 years also regarding level of education the majority of participant are Secondary education were(34.0%). regarding Marital status the majority of participant are married were(58.0%). Regarding the Economic level the majority of participant Averagewere(45.0%). Regarding the Chronic diseasethe majority of participant No.were(72.0%).

Table 2. Distribution of the Knowledge towards Vitamin D of the studied population

	N	%
Knowledge on sources of vitamin D		
Fruits & vegetables	84	42
Water	12	6
Mushroom	22	11
Fatty fish	32	16
Vitamin D supplements	28	14
Sun	68	34
Air	4	2
Selected cereals	34	17
Milk/Diary	40	20
Nuts	44	22
Cod-liver oil	18	9
Chicken	8	4
Egg	38	19
Beef	6	3
Don't know	74	37
Exposed to sun		
Yes	178	89
No	22	11
Sun exposure time		
Below 30 min	74	37
30 min above	40	20
More than 1 hours	68	34
Not exposed	18	9
Wearing full sleeved shirt /pant		
Yes	24	12
No	176	88
Covering of arms while working		
Yes	18	9
No	182	91
Wearing cap/umbrella		
Yes	40	20
No	160	80

Use of cosmetic cream contain SPF more than 8		
Yes	66	33
No	134	67
Use of sunscreen lotion		
Yes	18	9
No	182	91
Frequency of use cosmetic cream		
Once a day	48	24
Twice a day	40	20
More than 2	6	3
Not using	194	97
Duration of use		
1-5 years	38	19
6-10 years	22	11
Above 10 years	36	18
No	164	82

Table 2 shows Knowledge towards Vitamin D regarding the Knowledge towards Vitamin D the most of participant Fruits & vegetables were (42.0%) followed by Don't know were (37.0%), regarding Exposed to sun the majority of participant answer Yes were (89.0%) ,but regarding Sun exposure time the most of participant Below 30 min while regarding Wearing full sleeved shirt /pant the most of participant answer No were (88.0%), regarding the Covering of arms while the most of participant answer No were(91.0%). Regarding the Wearing cap/umbrella the most of participant answer No were (80.0%), regarding the Use of cosmetic cream contain SPF more than 8 the most of participant answer No were (67.0%), regarding the Use of sunscreen lotion the most of participant answer No were (91.0%), regarding the Frequency of use cosmetic cream the most of participant Not using were (97.0%), regarding the Duration of use the most of participant No were (82.0%)

Table 3. Distribution of the Knowledge towards Vitamin D of the studied population

	N	%
Knowledge on time needed in sun to get adequate vitamin D		
Less than 30 min	18	9
More than 30 min	34	17
More than 1hr	148	74
Knowledge of the adult women about the Benefits of vitamin D		
Bone health	52	26
Immune health	24	12
Prevents Rickets	38	19
Vision	30	15
Pregnancy & breast feeding	44	22
Hair growth	26	13
Diabetes mellitus	16	8
Cardiovascular health	12	6
Cognitive health	14	7
Cancer prevention	30	15
Cancer prevention	58	29
Calcium absorption	22	11
Prevent osteoporosis	16	8
Knowledge on factors affecting vitamin D level among adult women		

Skin pigment	38	19
Cloud & shade	22	11
Time of the day	38	19
Latitude	6	3
Season	24	12
Age	40	20
Pregnancy/ lactation	38	19
Fatty diet	12	6
Sunscreen use	22	11
Vegetarian diet	16	8
Diary allergy	18	9
Pollution	24	12
Wind	18	9
Smoking	20	10
BMI	36	18

Table 3 shows Knowledge on time needed in sun to get adequate vitamin D the most of participant More than 1hr were (74.0%), regarding Knowledge of the adult women about the Benefits of vitamin D answer Cancer prevention were (29.0%) ,but Knowledge on factors affecting vitamin D level among adult women the most of participant Pregnancy/ lactation were (20.0%),

Table 4: Distribution of the Knowledge regarding Vitamin D deficiency among female adult Makkah Al-Mokarramah, 2021

Knowledge		
	N	%
Weak	98	49
Average	66	33
High	36	18
Total	200	100
Chi-square	X²	28.84
	P-value	<0.001*

Table 4 shown, the results of the Vitamin D deficiency Knowledge a significantWeak relation were (98.0%) while p=0.001 and X²28.84.

Figure 1 Distribution of the Knowledge regarding Vitamin D deficiency among female adult Makkah Al-Mokarramah, 2021





Table (5) distribution of the relation of the Knowledge regarding Vitamin D deficiency among female adult and demographic data

			Weak		Average		High		Chi-square	
			98		66		36			
		N	N	%	N	%	N	%	X ²	P-value
Age	<25	56	39	39.80	12	18.18	5	13.89	28.124	0.0001
	25-30	74	19	19.39	35	53.03	20	55.56		
	30-35	38	22	22.45	11	16.67	5	13.89		
	>35	32	18	18.37	8	12.12	6	16.67		
Level of education	Primary	24	13	13.27	8	12.12	3	8.33	16.037	0.0136
	Intermediate	56	35	35.71	17	25.76	4	11.11		
	Secondary	66	28	28.57	21	31.82	17	47.22		
	High education	54	13	13.27	20	30.30	12	33.33		
Marital status	Married	170	67	68.37	30	45.45	28	77.78	140.896	<0.001*
	Unmarried	30	22	22.45	36	54.55	8	22.22		

Table 5 show the relation of the Knowledge regarding Vitamin D deficiency among female adult and demographic data regarding age a significant relation were P-value=0.0001 and X²5.920 in average and high, regarding Level of education no significant relation were P-value=0.0136 and X²16.037 in average . regarding Marital status a significant relation were P-value=0.001 and X²140.896 in average

Figure (2) distribution of the relation of the Knowledge regarding Vitamin D deficiency among female adult and demographic data



M

Conclusion

Health education interventions that increase awareness about vitamin D sources, especially food sources, are made. Also, educational interventions should focus on increasing awareness about the sufficient time of the day and duration for sun exposure to improve vitamin D status and the importance of the intake of vitamin D supplements as an affordable source to improve vitamin D status. Increasing males' awareness of the benefits of vitamin D is important to encourage them to adopt behaviors to improve vitamin D.

Discussion

In this study, the level and knowledge on vitamin D and adherence to supplementation of the female adult attending the the Primary Health clinic at Makkah, Saudi Arabia, 2021. Results revealed that majority of the mothers under study thought that vitamin D is important for the health of their the female adult, however large percentage of them had limited information about vitamin deficiency. Similarly, the work of Hoel and colleagues (25), Babelghaith and co-workers (26), Alamoudi and others (27) in Saudi Arabia showed that the participants under their studies exhibited limited knowledge when it comes to Vitamin D and its deficiency. Al-Saleh and others (28) mentioned that possible reasons for the said deficiency can be the insufficient knowledge about vitamin D, suggesting an increase in public awareness, especially to mothers, to be promoted to manage the increasing prevalence of Vitamin D deficiency and avoid its further complications. Contrary to the result of current study, a similar cross-sectional study in

Majmaah City, Saudi Arabia reported that majority (80.5%) of the participants were aware about vitamin D (29)

In this study, family is the major reported source (60.0%) of information concerning Vitamin D, followed by doctor (20.0%) and media (20.0%). Other studies in Saudi Arabia indicated the primary source of the Vitamin D knowledge of respondents as media (29), Doctor (20) and health care professionals (21). However, this value is two-fold lower than the reported overall knowledge mean score of 5.9 ± 1.2 of the participants in the study of Alamoudi and colleagues (17) in Jeddah, Saudi Arabia.

Regarding adherence to vitamin D supplementation, majority of the respondents was found to not adhere to vitamin D supplementation. In other Arab country, Alotaibi and co-workers (19) also reported that majority (74.8%) of the participants in their study did not take vitamin D supplements. Comparable results were also seen to other published non-Arab works wherein a percentage ranging from 2 to 19% of breastfeeding infants were reported to be under vitamin D supplementation (22, 23). Contrary to the result of current study, 90% have received vitamin D supplementation in the prospective cohort study of pregnant women and their children (24)

The low percentage of adherence to Vitamin D of the mothers under study can possibly be due to their poor knowledge on vitamin D and its deficiency. Doctors and family doctors allied health care providers have significant roles in advising or educating parents to practice supplementation for their female adult.

References:

1. Aljefree, N., Lee, P., & Ahmed, F. (2017, October). Exploring knowledge and attitudes about vitamin D among adults in Saudi Arabia: A qualitative study. In *Healthcare* (Vol. 5, No. 4, p. 76). MDPI.
2. Aljefree, N. M., Lee, P., & Ahmed, F. (2017). Knowledge and attitudes about vitamin D, and behaviors related to vitamin D in adults with and without coronary heart disease in Saudi Arabia. *BMC public health*, *17*(1), 1-12.
3. Babelghaith, S. D., Wajid, S., Al-Zaaqi, M. A., Al-Malki, A. S., Al-Amri, F. D., Alfadly, S., ... & Alarifi, M. N. (2017). Knowledge and practice of vitamin D deficiency among people lives in Riyadh, Saudi Arabia-A cross-sectional study. *Biomedical Research*, *28*(7), 3114-3118.
4. AlFaris, N. A., AlKehayez, N. M., AlMushawah, F. I., AlNaeem, A. N., AlAmri, N. D., & AlMudawah, E. S. (2019). Vitamin D deficiency and associated risk factors in women from Riyadh, Saudi Arabia. *Scientific reports*, *9*(1), 1-8.
5. Alotaibi, A. A., Alsalhi, W. A., Almutiri, A. N., Alzahrani, A. J., Alsaab, A. S., Abdurehman, M., ... & Aleliwi, Y. S. (2019). Knowledge and practice of vitamin D deficiency and risk of hair loss among adult population in Majmaah city, Saudi Arabia. *Int J Med Dev Ctries. Published online*, 173-178.
6. Alotaibi, A. A., Alsalhi, W. A., Almutiri, A. N., Alzahrani, A. J., Alsaab, A. S., Abdurehman, M., ... & Aleliwi, Y. S. (2019). Knowledge and practice of vitamin D deficiency and risk of hair loss among adult population in Majmaah city, Saudi Arabia. *Int J Med Dev Ctries. Published online*, 173-178.

7. Safdar, O., Baajlan, O., Alamri, A., Dahmash, R., Alloush, A., Ateeq, R., ... & Jiffri, H. (2019). Assessment of knowledge and awareness of vitamin D among physicians and students of healthcare in Jeddah, Saudi Arabia. *Australasian Medical Journal (Online)*, *12*(5), 143-153.
8. Roth, D. E., Abrams, S. A., Aloia, J., Bergeron, G., Bourassa, M. W., Brown, K. H., ... & Whiting, S. J. (2018). *Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low-and middle-income countries* (Vol. 1430, No. 1, pp. 44-79).
9. Cardwell, G., Bornman, J. F., James, A. P., & Black, L. J. (2018). A review of mushrooms as a potential source of dietary vitamin D. *Nutrients*, *10*(10), 1498.
10. Pilz, S., März, W., Cashman, K. D., Kiely, M. E., Whiting, S. J., Holick, M. F., ... & Zittermann, A. (2018). Rationale and plan for vitamin D food fortification: a review and guidance paper. *Frontiers in endocrinology*, *9*, 373.
11. Roth, D. E., Abrams, S. A., Aloia, J., Bergeron, G., Bourassa, M. W., Brown, K. H., ... & Whiting, S. J. (2018). *Global prevalence and disease burden of vitamin D deficiency: a roadmap for action in low-and middle-income countries* (Vol. 1430, No. 1, pp. 44-79).
12. Saggese, G., Vierucci, F., Prodam, F., Cardinale, F., Cetin, I., Chiappini, E., ... & Corsello, G. (2018). Vitamin D in pediatric age: consensus of the Italian Pediatric Society and the Italian Society of Preventive and Social Pediatrics, jointly with the Italian Federation of Pediatricians. *Italian journal of pediatrics*, *44*(1), 1-40.
13. Aparna, P., Muthathal, S., Nongkynrih, B., & Gupta, S. K. (2018). Vitamin D deficiency in India. *Journal of family medicine and primary care*, *7*(2), 324.
14. ElSORI, D. H., & Hammoud, M. S. (2018). Vitamin D deficiency in mothers, neonates and children. *The Journal of steroid biochemistry and molecular biology*, *175*, 195-199.
15. Zareef, T. A., Jackson, R. T., & Alkahtani, A. A. (2018). Vitamin D intake among premenopausal women living in Jeddah: food sources and relationship to demographic factors and bone health. *Journal of nutrition and metabolism*, *2018*.
16. Tariq, A., Khan, S. R., & Basharat, A. (2020). Assessment of knowledge, attitudes and practice towards Vitamin D among university students in Pakistan. *BMC Public Health*, *20*(1), 1-10.
17. Holick, M. F. (2018). Photobiology of vitamin D. In *vitamin D* (pp. 45-55). Academic Press.
18. Pludowski, P., Holick, M. F., Grant, W. B., Konstantynowicz, J., Mascarenhas, M. R., Haq, A., ... & Wimalawansa, S. J. (2018). Vitamin D supplementation guidelines. *The Journal of steroid biochemistry and molecular biology*, *175*, 125-135.
19. Syed, F., Latif, M. S. Z., Ahmed, I., Bibi, S., Ullah, S., & Khalid, N. (2019). Vitamin D deficiency in Pakistani population: critical overview from 2008 to 2018. *Nutrition & Food Science*.

20. Hastie, C. E., Mackay, D. F., Ho, F., Celis-Morales, C. A., Katikireddi, S. V., Niedzwiedz, C. L., ... & Pell, J. P. (2020). Vitamin D concentrations and COVID-19 infection in UK Biobank. *Diabetes & Metabolic Syndrome: Clinical Research & Reviews*, *14*(4), 561-565.
21. Hoeller, U., Baur, M., Roos, F. F., Brennan, L., Daniel, H., Fallaize, R., ... & Weber, P. (2016). Application of dried blood spots to determine vitamin D status in a large nutritional study with unsupervised sampling: the Food4Me project. *British Journal of Nutrition*, *115*(2), 202-211.
22. Goodman, S., Morrongiello, B., & Meckling, K. (2016). A randomized, controlled trial evaluating the efficacy of an online intervention targeting vitamin D intake, knowledge and status among young adults. *International Journal of Behavioral Nutrition and Physical Activity*, *13*(1), 1-13.
23. Toher, C., Lindsay, K., McKenna, M., Kilbane, M., Curran, S., Harrington, L., ... & McAuliffe, F. M. (2014). Relationship between vitamin D knowledge and 25-hydroxyvitamin D levels amongst pregnant women. *Journal of Human Nutrition and Dietetics*, *27*(3), 261-269.
24. Kotta, S., Gadhvi, D., Jakeways, N., Saeed, M., Sohanpal, R., Hull, S., ... & Griffiths, C. (2015). "Test me and treat me"—attitudes to vitamin D deficiency and supplementation: a qualitative study. *BMJ open*, *5*(7), e007401.
25. Hoel DG, Berwick M, de Gruijl FR, Holick MF. The risks and benefits of sun exposure 2016. *Dermatoendocrinol*. 2016 Jan-Dec;8(1):e1248325
26. Babelghaith SD, Wajid S, Al-Zaaqi MA, Al-Malki AS, Al-Amri FD, Alfadly S, et al. Knowledge and practice of vitamin D deficiency among people lives in Riyadh, Saudi Arabia - A cross sectional study. *Biomed Res*. 2017;28(7):3114-8.
27. Alamoudi LH, Almuteeri RZ, Al-Otaibi ME, Alshaer DA, Fatani SK, Alghamdi MM, et al. Awareness of Vitamin D Deficiency among the General Population in Jeddah, Saudi Arabia. *J NutrMetab*. 2019;2019:4138187.
28. Al-Saleh Y, Al-Daghri NM, Khan N, Alfawaz H, Al-Othman AM, Alokail MS, et al. Vitamin D status in Saudi school children based on knowledge. *BMC Pediatr*. 2015 May 6;15:53.
29. Alotaibi AA, Alsalhi WA, Almutiri AN, Alzahrani AJ, Alsaab AS, Alhassan MA, et al. Knowledge and practice of vitamin D deficiency and risk of hair loss among adult population in Majmaah city, Saudi Arabia. *IJMDC*. 2019;3(2):001-6