

KINEMATICS AND KINETICS PERFORMANCE DURING MENSTRUAL CYCLE PHASE AMONG ELITE ATHLETE

Ruaibah Yazani Tengah¹; Ashril Yusof², Abd. Halim Mokhtar², Nelfianty Mohd Rasyid¹

Universiti Pendidikan Sultan Idris¹; Universiti Malaya²

ruaibah@fsskj.upsi.edu.my

Abstract

Understanding the effect of menstrual cycle on kinematics and kinetics of exercise performance is important in the sense that menstrual cycle disrupts the hormonal balance of athletes. Changes in hormonal balance might affect exercise performance, and thus might affect longitudinal adaptations which occur from training, and also with biomechanical changes which increase the likelihood of injury. Twelve well trained elite female athletes voluntarily participated in this study. Thirteen kinematic and kinetic variables were measured via force platform during a counter movement jump exercise. General linear model one way repeated measure (ANOVA) and paired sample t test were used to meet the objective of this study. Three variables namely peak power ($p = 0.030$), mean power ($p = 0.010$) and peak velocity ($p = 0.016$) were significantly differed ($p < 0.05$) across the three menstrual cycle phases, where a reduction was clearly observed during the menses phase. Based on the results, menses phase was detrimental to exercise performance that requires explosive power component for elite level athletes. Thus, training with aims and goals of developing explosive power adaptation is not recommended to be carried out by elite athletes during their menses, with change of strategy during competition is suggested when a decrease in explosive power output might impair related sports specific performance. As a conclusion, athletes' menstrual cycle process should be monitored closely with training and competition for performance purposes.

Keywords: menses, hormonal responses, power, strength, force, velocity.

1. INTRODUCTION

Sports performance particularly relies a lot on strength and power capabilities of the athletes to sprint faster [1-2], jump higher [3-4], minimize the movement metabolic requirement [5-7] and improve other functional capabilities [8]. It is worth to note that a decrease or an increase of muscle performance depends on many factors which include mechanical [9-10], hormonal [11-12], metabolic [13] and neural [14-15] factors. Menstrual cycle phase experienced by the female athletes caused continuous changes in hormone concentration [16-17] and thus might alter the hormonal stimulus secreted and received during the strength training session, which might affect exercise performance. There have been many studies that have investigated the effect of menstrual cycle on human sports and exercise performance [18-27]. Certain phases of menstrual cycle have been identified to increase risk of anterior cruciate ligament injury [28-29]. This is not surprising as menses is a disruption to hormonal activities [30-32] and changes in hormonal function might affect physical activities performance. Female sex hormones fluctuation during the menstrual cycle is reported to have

significant effects on the neuromuscular system [33-35]. On the other hand, few studies performed on recreationally trained athletes or non-athletic population indicated that there were no significant differences on the neuromuscular performance during menstrual phases [36-38]. At present, little data is available regarding the effect of menstrual cycle on exercise performance in elite athletes. Other findings suggest that hormonal fluctuations due to menstrual cycle phases do not interfere with maximal intensity whole body sprinting and the metabolic responses to such exercise [39-40]. Since the finding remains inconclusive, further investigation in this area is needed. In term of team sports especially in relation to match or game based sports such as hockey matches, the ability to sustain high intensity intermittent loads is believed to be crucial for the final outcome of the match [41-42]. As mentioned earlier, strength and power will improve functional performance during the games which will contribute towards improving the physical performance of the athletes. Vertical jump exercise is one of the typically used exercises for strength and power training and assessment [43-45]. A decrease in vertical jump's kinematics and kinetics during training might decrease overall performance adaptation longitudinally. Therefore, the purpose of this study was to identify the differences in vertical jump kinematics and kinetics that might occur between the menses, follicular phase and luteal phase of the menstrual cycle among elite athletes.

2. METHODS

2.1 Experimental Approach to the Problem

Acute randomized within-subjects cross-over design was used in this study, where subjects were asked to attend one familiarization and three testing sessions. The subjects randomly performed the testing session starting at different phases of the menstrual cycle which includes i) Menses (Day 1 – 5); ii) Follicular Phase (Day 6 -13); and, iii) Luteal Phase (Day 15 – to next cycle). This design allowed the subjects to complete testing sessions within one menstrual cycle, and some were allowed to carry over the testing sessions into their next cycle. Two weeks before their first testing session all subjects were asked to attend a familiarization session at the Exercise Physiology Laboratory, National Institute of Sports, Malaysia. During the familiarization session, the subjects were given briefings on exercise and experimental protocols, procedures and exercise techniques.

2.2 Subjects

Twelve well trained elite female intermittent sport athletes voluntarily participated in this study. All subjects were elite athletes ranging from intermittent sport athletes with the national team of Malaysia at the time of the testing. Subjects' selection criteria were, aged between 18-28 years old with a maximum oxygen uptake ($\dot{V}O_2\text{max}$) of ≥ 50 ml/kg/min, healthy with no allergies, free of metabolic, cardiovascular or respiratory diseases, non-smoker and regular menstrual cycles lasting 28 ± 2 days in duration (3 month monitoring of regularity). All subjects were fully informed of the study's purposes and possible risks before signing a consent form. Subjects were also told that they are free to withdraw at any stage of the study without the obligation to give reasons. Prior to commencing the study, approval was obtained from the ethical committee of University of Malaya and National Sports Council, Malaysia.

2.3 Equipment

Counter movement vertical jump was conducted on a force plate (400 Series Performance Force Plate, Fitness Technology, Australia) interfaced with computer software (Ballistic Measurement System, Fitness Technology, Australia) that allowed direct measurements of displacement, force, velocity and power variables. Data sampling rate was set at 200 Hz with filter cut-off frequency rate of 10 Hz. Jump height was estimated using the following equation [$h = t^2 \cdot g / 8$]. Where g is the gravitational acceleration estimated at 9.81ms^{-2} and is the flight time.

2.4 Procedures

Preliminary Assessments and Familiarization: Subjects' body height and weight were measured after the briefing and their informed consent form were collected. Subjects were then showed the counter movement vertical jump technique, followed by several trials of the jump by each of the subject on the force plate. Improper technique was corrected and the subjects performed several more trials until the researcher was satisfied with the jump technique. Verification of menstrual cycle phase: Menstrual phases of each subject was verified using blood samples to determine the level of estrogen, progesterone, LH and FSH based on method by [46] along with basal body temperature (BBT) charts [47].

Counter movement vertical jump technique: The counter movement vertical jump ready position was standardized with subjects standing with the feet approximately shoulder width apart and hands on their hips. The jumps began with the subjects lowered down their buttock, bend their knees to quarter-squat position, and later immediately straighten the knee with feet pushing the ground for their take-off. The subjects were then considered had completed one repetition of the jump when they landed with their hands still on their hip within the force plate area with the knee again bend during the landing to absorb impact. The subjects made another jump straight-away until desired repetitions were achieved.

Testing procedures: The subjects were tested randomly at different phases of their menstrual cycle. Method of assessment was replicated as previously described [48] with all subjects tested at similar time of the day for each testing occasion. During each testing, the subjects performed 4 series of 15 seconds vertical counter-movement jump (CMJ). All subjects started their testing session with a standardized general warm-up and practiced jumping several times until they were comfortable and consistent with it. The subjects performed 4 trials x 15 seconds maximum effort counter movement vertical jumps. The subjects were instructed to jump as high as possible on each of the single repetition, with 10 seconds rest periods were given between each trial. Each testing session ended with cooling down and stretching exercises.

2.5 Data Analysis

The force plate measures all dependent variables of interest at a sampling frequency of 200 Hz. Variables of interest include peak force (N), mean force (N), peak power (W), mean power (W), jump height (m), peak distance (m), min distance (m), peak velocity ($\text{m}\cdot\text{s}^{-1}$), min Velocity ($\text{m}\cdot\text{s}^{-1}$), jump repetition, impulse at 100 (ms), 200 (ms) and 300 (ms) were calculated for each set of the jumps involved via the BMS software data analysis program. The variables of interest in this study have been proven to be stable within and between sessions (i.e. typical CV < 5% and ICC~0.95).

2.6 Statistical Analysis

All data from variables of interest during the three phases of menstrual cycle were analyzed using a general linear model one way repeated measure (ANOVA). A one sample Kolomogorov Smirnov test was utilized to determine the normal distribution of all variable. In the cases where a Mauchly's test of sphericity assumption was violated, the Greenhouse-Geisser value was utilized to determine significance. Further analyses using paired sample t test were used to identify relative changes between menses versus luteal phase and follicular phase versus menses. The significance level of this study was set as $p < 0.05$.

3. RESULTS

3.1 Counter Movement Jump (CMJ)

Mean values (\pm SD) for thirteen selected kinematic and kinetic components for CMJ performance during different phases of the menstrual cycle are presented in Table 1. Significant differences ($p < 0.05$) were found in peak power, mean power and peak velocity output between the three phases based on one way repeated measure (ANOVA), while paired simple t test indicated that the significant difference were actually originated from the differences between luteal phase and menses phase.

Table 1:
Mean \pm SD of counter movement jump performance during menstrual cycle phases
(Average of 4 trials x 15s counter movement jump).

Variable (n =12)	Luteal Phase Mean \pm SD	Menses Mean \pm SD	Follicular Phase Mean \pm SD	P Value
Peak Force (Newton)	1154.55 \pm 114.79	1130.98 \pm 130.20	1114.26 \pm 102.08	0.4388
Mean Force (Newton)	863.41 \pm 66.27	845.01 \pm 69.12	841.10 \pm 56.35	0.2027
Peak Power (Watts)	2880.88 \pm 375.46*	1952.30 \pm 135.64	2069.25 \pm 216.99	0.0304
Mean Power (Watts)	1087.24 \pm 168.78*	987.05 \pm 91.81	1009.22 \pm 98.37	0.0096
Jump Height (m)	0.23 \pm 0.02	0.22 \pm 0.02	0.22 \pm 0.03	0.0781
Peak Distance (m)	1.28 \pm 1.55	0.50 \pm 0.48	0.91 \pm 0.87	0.1893
Min Distance (m)	0.14 \pm 2.66	-1.38 \pm 0.65	0.38 \pm 0.82	0.2037
Peak Velocity (m.s-1)	2.39 \pm 0.25*	2.19 \pm 0.14	2.31 \pm 0.16	0.0160
Mean Velocity (m.s-1)	-1.33 \pm 0.25	-1.23 \pm 0.37	-1.37 \pm 0.19	0.4292
Jump Repetition	15.66 \pm 0.9	15.62 \pm 0.70	15.52 \pm 1.3	0.7713
Impulses at 100 (ms)	95.92 \pm 10.29	93.12 \pm 9.20	92.08 \pm 7.0	0.2200
Impulses at 200 (ms)	196.14 \pm 21.98	191.02 \pm 22.20	186.80 \pm 21.31	0.2284
Impulses at 300 (ms)	285.96 \pm 29.83	279.05 \pm 29.13	273.62 \pm 32.12	0.2378

*denote significant between Luteal Phase vs Menses: $p < 0.05$

3.2 Peak power (W)

The analysis of variance for the peak power (W) is shown in Table 1. The peak power (W) varied significantly between the phases in the menstrual cycle [$F(2, 11) = 4.112$, $p < 0.05$]. Pairwise comparison showed that relative peak power (W) responses were affected during M -LP than FP -M.

3.3 Mean power (W)

The analysis of variance for the mean power (W) is shown in Table 1. Mean power (W) was significantly differed in three phases of menstrual cycle [F (2, 11) = 5.784, p < 0.05]. Relative change of mean power slightly decreased before menses (M – LP) compared to after menses (FP – M).

3.4 Peak velocity (m.s-1)

The analysis of variance for the peak velocity (m.s-1) is shown in Table 1. Peak velocity (m.s-1) was found to be significantly differed between the phases of the menstrual cycle [F (2, 11) = 5.020, p < 0.05]. Relative change of peak velocity (m.s-1) was significantly higher during LP – M then FP – M.

3.5 Estrogen / Progesterone Ratio

Mean values (\pm SD) of serum estrogen and progesterone levels in the different phases of the menstrual cycle for the 12 subjects are presented in Table 2. Estrogen levels were low in the menses phase and increased as expected in the luteal phase and follicular phase. Progesterone levels increased in the luteal phase compared to other phases.

Table 2

Mean values (\pm SD) of serum estrogen and progesterone levels in the different phases of the menstrual cycle.

Hormone	Luteal Phase	Menses	Follicular Phase
Estrogen (pmol/I)	405.08 \pm 232.27*	158.58 \pm 43.89	255.50 \pm 118.08
Progesterone (nmol/I)	3.86 \pm 2.7 *	1.94 \pm 0.5	2.46 \pm 0.8
E/P Ratio	104.94	81.74	103.86

*denote significant at luteal phase when compare with menses with p < 0.05,

E = Estrogen; P = Progesterone

3.6 Estrogen (pmol/I)

The estrogen level was significantly differed between the phases of menstrual cycle, F (2, 22) = 7.246, p < 0.05].The estrogen level during luteal phase (405.08 \pm 232.27 pmol/I) was higher compared to menses (158.58 \pm 43.89 pmol/I) and follicular phase (255.50 \pm 118.08 pmol/I).

3.7 Progesterone (nmol/I)

The Progesterone level was significantly differed between the phases of menstrual cycle, [F (2, 22) = 4.366, p < 0.05].The progesterone level during luteal phase (3.86 \pm 2.7 nmol/I) was higher compared to menses (1.94 \pm 0.5 nmol/I) and follicular phase (2.46 \pm 0.8 nmol/I).

DISCUSSION

The main aim of this study is to investigate how some selected kinematics and kinetics were affected by menstrual cycle phases, when performing acute explosive anaerobic type activities such as the

counter movement jump among well trained elite athletes. The most important finding is that significant differences in peak power (W), mean power (W) and peak velocity (m.s⁻¹) between the three menstrual cycle phases. Other variables measured such as peak force (N), mean force (N), jump height (m), peak distance (m), min distance (m), min velocity (m.s⁻¹), jump repetition, impulse at 100 (ms), impulse at 200 (ms) and impulse at 300 (ms) did not show any difference during the menstrual cycle phases, which confirmed what have been found by previous studies [49]. Before further discussions are made on the key findings of this study, several factors that might influence the results obtained should also be noted. First limiting factor that may affect the results of this study is the timing of each testing with regards to the day of the menstrual cycle phases. Janse de Jonge [50] recommended that testing should be conducted on the same day (i.e., 3rd day of LP, M and FP). However in our study it was difficult to gather the subjects to be tested on similar timing each of their menstrual cycle phases. Thus, this might have some minor impact on the results obtained, due to fluctuations in hormone levels within each day of the menstrual phases [51]. Although it was difficult to gather elite athletes at one specific menses timing, all subjects were tested at similar time of the day, to ensure minimal differences in the circadian variations.

The second factor is the pulsatile secretion of hormone. In order to minimize inaccuracy that might be caused by higher secretion due to exercise or time of the day, all subjects were tested within the same time of the day as indicated earlier. All testing's were conducted at rest condition. It has been reported that while the level of estrogen might be similar between subjects, the progesterone level might fluctuate and vary between subjects, and thus may influence the physical exercise performance [52, 50, 53]. Based on observation made on both hormones secreted in this study (Table 2), fluctuations on the estrogen and progesterone ratio during each testing period existed. Therefore this result suggested that the changes in female steroid hormones during the menstrual cycle may affect the mechanical properties of human muscle and tendon. A previous study had also found that muscle fiber recruitment / motor unit activations were negatively affected across menstrual cycle phase [51].

The main findings of this study were reduction in peak power, mean power and peak velocity during the menses phase, and these findings are in agreement with a number of previous studies [54, 49, 55]. However, none of these cited studies used well-trained high performance athletes as subjects. Another study explored by [56] conducted on well trained female athletes showed that they felt nastiest before the menstruation phase, performance level remained the same during menstruation and the performance was poorer following menstrual phase in the first 14 days. Thus, the present study has shown that the well trained athletes also reacted similarly as the untrained subjects in this matter. Training status has previously been shown to have an impact on power production ability and utilization of muscle activities during explosive or non-explosive anaerobic type of exercise or movement [57-58] but this might exclude female athletes with menses, in which their training status might not made them 'immune' towards the effect of menstrual cycle on performance. It could be suggested that, during training or competition, athletes with menses should shift their game play from relying more on power based movement into skills or movement that requires less of this, in order to maintain their typical performance. This kind of change of strategy or skills used might minimize the negative effect assaulted with menstrual cycle experienced by the trained athletes. However, this is far from definite and further studies are on the way to elucidate this.

Soares *et.al*, [59] suggested that the behaviour of the muscles in women presents different characteristics during different phases of the menstrual cycle, in particular, at the end of the luteal phase. Findings of their study also supported what have been found previously, that different neuromuscular patterns were utilized when performing jump sequence during different phase of

menses, as luteal phase has higher estrogen level compared to during early follicular phase, which has lower level of estrogen [60]. Our results also suggested that estrogen and progesterone levels decreased during the menses phase in which the reduction in peak power, mean power, and peak velocity were observed.

Another explanation that may be considered for the reduction in peak power, mean power and peak velocity is pre-menstrual syndrome (PMS). PMS is characterized by cyclical physical and mood disturbance during the luteal phase, where severity of the symptoms (i.e., pain) gradually increases during the luteal phase and disappears a few days after onset of menses [49]. Giacomoni *et.al.*, [61] proposed that although there were no significant differences in maximal anaerobic performance during different menstrual cycle phases, results of their study suggested that the presence or absence of premenstrual or menstrual syndrome symptoms may have an effect, possibly through an action on the stretch shortening cycle of tendons and ligaments. Research has provided strong evidence that PMS symptoms are the result of an abnormal response to normal levels of estrogen and progesterone [62].

Results of this study indicate the need of some considerations to be taken when managing strength and power training for female elite athletes especially during their menses. While the findings of this study is far from definite, coaches and athletes alike should take into consideration to properly monitor female elite athletes that need to perform explosive power type of movement during training and competition. Changes of strategies especially in sports with explosive type of movement might be required in order to ensure continuity of desirable peak performance during competition. Training sessions with an emphasis and goal on peak power output should be adjourned to another day to ensure maximum adaptations can take place longitudinally. A proper monitoring system for female athletes menstrual cycle process should also be a part of conditioning training plans, should include both performance related reasons and injury preventions. However, the results obtained in this study still need to be investigated further, especially by performing longitudinal research and taking into accounts other factors involved such as muscle activation, firing rate and many more before firm conclusion can be elucidated.

Acknowledgement

The authors herewith acknowledging the Research Management and Innovation Centre, Sultan Idris Education University, Malaysia for their support and cooperation to publish this article.

REFERENCES

- [1] Cronin, J., Ogden, T., Lawton, T., & Brughelli, M. (2007). Does increasing maximal strength improve sprint running performance? *Journal of Strength and Conditioning Research*. 29:86-95.
- [2] McMaster, D.T., Gil, N. & Cronin, J. (2014). A brief review of strength and ballistic assessment methodologies in sport. *Sports Med*. 44:603–23.

- [3] Cormie, P., McGuigan, M.R., & Newton, R.U. (2010). Influence of strength on magnitude and mechanisms of adaptation to power training. *Medicine Science Sports Exercise*. 42. 1566-1581.
- [4] Thomas, C., Comfort, P., Jones, P., A. & Dos'Santos, T. (2017). A Comparison of Isometric Midthigh - Pull Strength, Vertical Jump, Sprint Speed, and Change-of-Direction Speed in Academy Netball Players. *International Journal of Sports Physiology and Performance*. Vol 12 (7). pp 916–921
- [5] Storen, O., Helgerud, J., Stoa, E.M. & Hoff, J. (2008). Maximal strength training improves running economy in distance runners. *Medicine Science Sports Exercise*. 40:1089-1094.
- [6] Kempton, T., Sirotic, A.,C. Rampinini, E. & Coutts, A., J. (2015). Metabolic Power Demands of Rugby League Match Play. *International Journal of Sports Physiology and Performance*. Vol 10 (1). pp 23–28.
- [7] Hader, K., Mendez-Villanueva, A., Palazzi, D., Ahmaidi, S., & Buchheit, M. (2016) Metabolic Power Requirement of Change of Direction Speed in Young Soccer Players: Not All Is What It Seems. *PLoS ONE*. 11(3)
- [8] Harris, G.R., Stone, M.H., O'Bryant, H.S., Proulx, C.M., & Johnson, R.L. (2000). Short-term performance effects of high power, high force, or combined weight-training methods. *Journal Strength Conditioning Research*. 14:14-20.
- [9] Baker, D., Nance, S. & Moore, M. (2001). The load that maximizes the average mechanical power output during jump squats in power-trained athletes. *Journal Strength Conditioning Research*. 15 (1), 92-97.
- [10] Mizrahi, J. (2015). Mechanical Impedance and Its Relations to Motor Control, Limb Dynamics, and Motion Biomechanics. *Journal Of Medical And Biological Engineering*. 35, 1–20.
- [11] Bamman, M.M, Shipp, J.R., Jiang, J., Gower, B.A., Hunter, G,R., Goodman, A., McLafferty Jr., C.,L. & Urban, R.,J. (2001). Mechanical load increases muscle IGF-I and androgen receptor mRNA concentrations in humans. *American Journal of Physiology-Endocrinology and Metabolism*. Vol. 280 (3). E383–E390.
- [12] Tieland, M., Trouwborst, I & Clark, B., C. (2018) Skeletal muscle performance and ageing. *Journal of Cachexia, Sarcopenia and Muscle*. 9:3–19
- [13] de Freitas M.,C, Gerosa-Neto, J., Zanchi, N.,E. Lira, F.,S. & Rossi, F.,E. (2017). Role of metabolic stress for enhancing muscle adaptations: Practical applications. *World Journal Methodology*. 7(2):46-54.
- [14] Latash, M., L. (2016). Towards physics of neural processes and behaviour. *Neuroscience & Biobehavioral Reviews*. Vol 69. pp. 136-146.
- [15] Brennecke, A., Guimarães, T.M., Leone, R., Cadarci, M., Mochizuki, L., Simão, R., Amadio, A.C., & Serrão, J. (2009). Neuromuscular activity during bench press exercise performed with and without the preexhaustion method. *Journal Strength Conditioning Research*. 23(7): 1933-1940.
- [16] Lebrun, C.M., & Rumball, J.S. (2001). Relationship between athletic performance and menstrual cycle. *Current Women's Health Reports*. 1(3):232-240.

- [17] Davis, H.C., & Hackney, A.C. (2017). *The Hypothalamic–Pituitary–Ovarian Axis and Oral Contraceptives: Regulation and Function*. Sex Hormones, Exercise and Women. pp1-17.
- [18] Bambaeichi, E., Reilly, T., Cable, N.T., & Giacomoni M. (2004). The isolated and combined effects of menstrual cycle phase and time-of-day on muscle strength of eumenorrheic females. *Chronobiol International*. 21:645-660.
- [19] Montgomery, M.M. & Shultz, S.J. (2010) Isometric Knee-Extension and Knee-Flexion Torque Production During Early Follicular and Postovulatory Phases in Recreationally Active Women. *Journal of Athletic Training*. Vol. 45 (6). pp. 586-593.
- [20] Sung, E., Han, A., Hinrichs, T., Vorgerd, M., Machado, C. & Platen, P. (2014). Effects of follicular versus luteal phase-based strength training in young women. *Springer Plus*. 3(1):668.
- [21] Czajkowska, M., Drosdzol-Cop, A., Gałazka, I., Naworska, B. & Skrzypulec-Plinta, V. (2015). Menstrual Cycle and the Prevalence of Premenstrual Syndrome/Premenstrual Dysphoric Disorder in Adolescent Athletes. *Journal Paediatric Adolescence Gynecol*. 28:492–498
- [22] Tenan, M., Hackney, A.C. & Griffin, L. (2016). Maximal Force And Tremor Changes Across The Menstrual Cycle. *European Journal Applied Physiology*. 116:153–60
- [23] Ana L. S. Gil, Gabriel R. Neto , Maria S. C. Sousa, Ingrid Dias, Jeferson Vianna, Rodolfo A. M. Nunes & Jefferson S. Novaes. (2015). Effect of strength training with blood flow restriction on muscle power and submaximal strength in eumenorrheic women. *Clinical Physiology and Functional Imaging*. Vol 37 (2). pp 221-228.
- [24] Pallavi, L. C., D Souza, U. J., & Shivaprakash, G. (2017). Assessment of Musculoskeletal Strength and Levels of Fatigue during Different Phases of Menstrual Cycle in Young Adults. *Journal of clinical and diagnostic research*. 11(2), CC11–CC13.
- [25] Wikström-Frisén, L., Boraxbekk, C.J., & Henriksson, L. (2017). Increasing training load without risking the female athlete triad: menstrual cycle based periodized training may be an answer? *Journal Sports Med Physical Fitness*. 57(11):1519-1525
- [27] Köse, B. (2018). Analysis of the Effect of Menstrual Cycle Phases on Aerobic-Anaerobic Capacity and Muscle Strength. *Journal of Education and Training Studies*. Vol 6, No 8. pg 21-28.
- [28] Romero-Moraleda, B., Coso, J. D., Gutiérrez-Hellín, J., Ruiz-Moreno, C., Grgic, J. & Lara, B. (2019). The Influence of the Menstrual Cycle on Muscle Strength and Power Performance. *Journal of human kinetics*. 68, 123–133.
- [29] Shultz, S.J., Schmitz, R. J. & Beynon, B.,D. (2011). Variations in varus/valgus and internal/external rotational knee laxity and stiffness across the menstrual cycle. *Journal of Orthopedic Research*. 29 (3). 318–325.
- [30] Balachandar, V., Marciniak, J. L., Wall, O. & Balachandar, C. (2017). Effects of the menstrual cycle on lower-limb biomechanics, neuromuscular control, and anterior cruciate ligament injury risk: a systematic review. *Muscles, Ligaments And Tendons Journal*, 7(1), 136–146.

- [31] Ahrens, K.A., Vladutiu, C.J., Mumford, S.L., Schliep, K.C., Perkins, N.J., Wactawski-Wende, J & Schisterman, E.F. (2014). The effect of physical activity across the menstrual cycle on reproductive function. *Annals of Epidemiology*. 24(2). 127–134.
- [32] Caldwell, A.E., Hooper, P.L. (2017). Sex Hormones and Physical Activity in Women: An Evolutionary Framework. In: Hackney A. (eds) *Sex Hormones, Exercise and Women*. Springer, Cham
- [33] Ekenros, L., Papoutsis, Z., Fridén, C., Dahlman, W. K., & Lindén, H. A. (2017). Expression of sex steroid hormone receptors in human skeletal muscle during the menstrual cycle. *Acta Physiology*. 219 (2). 486-493.
- [34] Tornberg, A. B., Melin, A, Koivula, F.M., Johansson, A., Skouby, S., Faber, J. & Sjödin, A. (2017). Reduced Neuromuscular Performance in Amenorrheic Elite Endurance Athletes. *Medicine Science Sports Exercise*. Vol. 49 (12). pp. 2478–2485.
- [35] Ansdell, P, Brownstein, C.G., Škarabot, J., Hicks, K.M., Simoes, D.C.M., Thomas, K., Howatson, G, Hunter, S.K., & Goodall, S. (2019). Menstrual cycle-associated modulations in neuromuscular function and fatigability of the knee extensors in eumenorrheic women. *Journal of Applied Physiology*. 126.6, 1701-1712
- [36] Lebrun, C.M, McKenzie, D.C, Prior, J.C., & Taunton, J.E. (1995). Effects of menstrual cycle phase on athletic performance. *Medicine Science Sports Exercise*. 27:437- 444.
- [37] Bennal, A.S., Chavan, V., Taklikar, R.H, & Takalkar, A. (2016). Muscular performance during different phases of menstrual cycle. *Indian Journal of Clinical Anatomy and Physiology*. 3(1):1-3 1
- [38] Ronaldo K, Irineu, L. Renato B, Saulo G, Rogério C, Carlos U, Hamilton R. & Valmor, T. (2017). Effects of Different Combinations of Strength, Power, and Plyometric Training on the Physical Performance of Elite Young Soccer Players. *The Journal of Strength & Conditioning Research*. Vol 31 (6). pp 1468-1476
- [39] Tsampoukos, A., Peckham, E., James, R., & Nevill, M. (2010). Effect of menstrual cycle phase on sprinting performance. *European Journal of Applied Physiology*. 109:659-667.
- [40] Julian, R., Hecksteden, A., Fullagar, H. H., & Meyer, T. (2017). The effects of menstrual cycle phase on physical performance in female soccer players. *PloS one*, 12(3).
- [41] Reilly, T., & Borrie, A. (1992). Physiology applied to field hockey. *Sports Medicine*. 14:10-26.
- [42] Woznak, C., & Sadowski, C. (2018) Anterior Cruciate Ligament Injuries. (2018). The Evidence That Oral Contraceptives Reduce the Risk in Female Athletes. *Journal of Orthopaedics for Physician Assistants*. Vol 6(3). pp e23.
- [43] Hanson, E.D., Leigh, S., & Mynark, R.G. (2007). Acute effects of heavy-and light-load squat exercise on the kinetic measures of vertical jumping. *Journal Strength Conditioning Research*. 21:1012-1017
- [44] Hester, G.M., Pope, Z.K., Sellers, J.H., Thiele, R.M., & De Freitas, J.M. (2017). Potentiation: Effect of ballistic and heavy exercise on vertical jump performance. *Journal of Strength & Conditioning Research*. Vol 31(3). 660–666

- [45] Grgic, J., Trexler, E.T., & Lazinica, B. (2018). Effects of caffeine intake on muscle strength and power: a systematic review and meta-analysis. *Journal of the International Society of Sports Nutrition*. 15, 11
- [46] Kleiblova, P., Springer, D., & Haluzik, M. (2006). The influence of hormonal changes during menstrual cycle on serum adiponectin concentrations in healthy women. *Physiological Research*. 55:661-666.
- [47] Guermandi, E., Vegetti, W., Bianchi, M.M., Uglietti, A., Ragni, G., & Crosignani, P. (2001). Reliability of ovulation tests in infertile women. *Journal of Obstetrics and Gynaecology*. 97:92- 96.
- [48] Friden, C., Hirschberg, A.L., & Saartok, T. (2003). Muscle strength and endurance do not significantly vary across 3 phases of the menstrual cycle in moderately active premenopausal women. *Clinical Journal Sport Medicine*. 13:238-41.
- [49] Hespanhol, J.E., da Silve Neto, L.G., & de Arruda, M. (2006). Reliability of the four series 15second vertical jumping test. *Revista Brasileira de Medicina do Esporte*. 12:95-98.
- [50] Janse de Jonge, XAK. (2003). Effects of the menstrual cycle on exercise performance. *Sports Medicine*. 33:833-851.
- [51] Bambaiechi, E., Reilly, T., Cable, N.T., & Giacomoni M. (2004). The isolated and combined effects of menstrual cycle phase and time-of-day on muscle strength of eumenorrhic females. *Chronobiol International*. 21:645-660.
- [52] Bunt, J.C. (1990). Metabolic actions of estradiol: significance for acute and chronic exercise responses. *Medicine Science Sports Exercise*. 22:286-290.
- [53] Dawson, E., A. & Reilly T. (2009) Menstrual cycle, exercise and health. *Biological Rhythm Research*, 40:1. 99-119.
- [54] Wojtys, E.M., Huston, L.J., Lindenfeld, T.N., Hewett, T.E., & Greenfieldn M.L.V.H. (1998). Association between the menstrual cycle and anterior cruciate ligament injuries in female athletes. *American Journal of Sports Medicine*. 26:614-619
- [55] Nicolay, C.W., Kenney, J.L., & Lucki, N.C. (2007). Grip strength and endurance throughout the menstrual cycle in eumenorrhic and women using oral contraceptives. *International Journal of Industrial Ergonomics*. 37:291-301.
- [56] Sung, E., Han,A., Hinrichs, T., Vorgerd, M., Manchado,C., & Platen, P. (2014). Effects of follicular versus luteal phase-based strength training in young women. *Springerplus*. 3: 668.
- [57] Pick, J., & Becque, M, D. (2000). The relationship between training status and intensity on muscle activation and relative submaximal lifting capacity during the back squat. *Journal Strength Conditioning Research*.14:175-181.
- [58] Baker, D., Nance, S. & Moore, M. (2001). The load that maximizes the average mechanical power output during jump squats in power-trained athletes. *Journal Strength Conditioning Research*, 15 (1), 92-97.
- [59] Soares, F.A., Salomoni, S.E., Veneziano, W.H., de Carvalho J.L.A., de Oliveira Nascimento, F.A., Pires, K., & da Rocha, A.F. (2011). On the behavior of surface electromyographic variables during the menstrual cycle. *Physiological Measurement*. 32:543-57.

- [60] Dedrick, G.S., Sizer, P.S., Merkle, J.N., Hounshell, T.R., Robert - McComb, J.J., Sawyer, S.F., Brismee, J-M. & Roger, J.C. (2008). Effect of sex hormones on neuromuscular control patterns during landing. *Journal of Electromyography and Kinesiology*. 18:68-78.
- [61] Giacomoni, M., Bernard, T., Gavarry, O., Altare, S., & Falgairette, G. (2000). Influence of the menstrual cycle phase and menstrual symptoms on maximal anaerobic performance. *Medicine Science Sports Exercise*. 32:486-92.
- [62] Schmidt, P.J., Nieman, L.K., Danaceau, M.A., Adams, L.F., & Rubinow, D.,R. (1998). Differential behavioral effects of gonadal steroids in women with and in those without premenstrual syndrome. *Obstetrical & Gynecological Survey*. 53:291-292.