

Reasons For 1-17-Year-Old Children To Visit A Dentist During The Covid-19 Pandemic

M.K. Kamalova¹, I.V. Fomenko², D.S. Dmitrienko², N.V. Matvienko², E.N. Arjenovskaya²,
A.G. Gevorkyan², K.V. Nikitina², E.E. Maslak²

¹*Doctor of Philosophy of Medical Sciences, Head of the Department of Surgical Dentistry, Bukhara State Medical Institute of the Ministry of Health of Uzbekistan.*

²*Head of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia. Volgograd, Russian Federation*

Abstract—COVID-19 pandemic significantly affected dentistry due to high coronavirus infectivity and dangerous consequences. Special recommendations were developed for dentists to limit the infection of staff and patients. to study the reasons for children visiting dental clinics during COVID-19 pandemic and give the characteristics of the patients and provided dental treatment. The cross-sectional study was conducted in four dental clinics, which worked for dental emergency in the cities of Volgograd and Volzhky during the peak of COVID-19 pandemic. The data about 166 paediatric dental patients aged 1-17 years were collected. The reasons for children's dental visits, age and behavior of the children, the tooth types, and the types of provided treatment were analyzed. The children's behavior was assessed according to Frankl's Behavior Rating Scale. Proportions (%), 95% Confidence Interval (CI), mean-value with standard error ($M \pm m$) were calculated. The differences were assessed by Pierson's Chi-square test at p -value 0.05. Microsoft-office Excel-2019 programs were used for the statistic analysis. The majority (98.8%, 95% CI 95.7-99.7%) of the children visited dental clinics due to acute toothache. The primary teeth needed urgent treatment significantly more often than permanent teeth: 83.7% (95% CI 77.8-88.3%) and 16.3% (95% CI 11.7-22.2%) respectively, $\chi^2 = 172.171$, $p < 0.0001$. Among the primary teeth the most affected ones were the first and second molars, among the permanent teeth – the first molar. The common reasons for urgent dental treatment in children were caries consequences: apical periodontitis (51.6%, 95% CI 44.5-58.6) or pulpitis (35.3%, 95% CI 28.8-42.3). The other reasons were dental trauma, caries, periostitis, physiological primary tooth mobility. Acute pain in the primary teeth was caused mainly by apical periodontitis, in the permanent teeth by pulpitis. The applied treatment methods were aimed at limiting the use of aerosol generating procedures, reducing the duration of the treatment and preventing the patients' hospitalization. Almost all the primary teeth with apical periodontitis were extracted; the number of extracted and endodontically treated permanent teeth was approximately equal. The current methods of pulpitis treatment (pulp capping, pulpotomy and pulpectomy) were applied along with the pulp necrotizing agent use. Many (39.7%, 95% CI 32.6-47.4%) children demonstrated negative behavior. Behavior management techniques were used, including tell-show-do (54.5%, 95% CI 42.6-66.0%), sedation (3.0%, 95% CI – 0.8-10.4%), general anesthesia (3.0%, 95% CI – 0.8-10.4%), passive restraint (39.4%, 95% CI 28.5-51.4%). In the limitation of this study we can conclude that COVID-19 pandemic changed the profile of paediatric dental patients and dentist's treatment options. In 1-17-year-old children the main reason for an emergency dental visit was acute dental pain due to caries consequences. The primary teeth needed urgent dental treatment significantly more often than the permanent teeth. The children often demonstrated negative behavior and required behavior management. The applied treatment methods were aimed at limiting the use of

aerosol generating procedures, decreasing the duration of the treatment and avoiding the patients' hospitalization.

Key words: COVID-19, dentistry, children, emergency, dental treatment.

1. INTRODUCTION

World Health Organization (WHO) confirmed new coronavirus infection as pandemic COVID-19 on 11 March, 2020 [1]. By the date of 12.06.2020 there were more than 7.5 million people infected and more than 420 thousand died [2]. This infection is very contagious and is easily transmitted with droplets and aerosols [3]. The specific feature of dentistry is working in the area of patients' breath, often in contact with blood and saliva. Thereby, dentistry was recognized as the professional activity with dangerous environment regarding virus transmission from patients to dental staff, from staff to patients or cross-infection among the patients [4]. A lot of dentists concern about themselves and their families' infection but do not follow current protective recommendations completely [5]. In many countries during the peak of COVID-19 pandemic dental offices were closed except emergency. The rationale for this decision was the attempt to prevent unnecessary coronavirus infection of dental staff and dental patients and to avoid the overloading of the healthcare system [6,7].

The concept of dental emergency includes the following conditions. Actually dental emergency has been designed for the immediate treatment of dental diseases which are life threatening (uncontrolled bleeding, aggravated bacterial inflammation, severe soft tissue and bone trauma, etc.). Urgent dental care aimed at relieving severe dental pain, alleviating the risk of inflammation increase which could lead to emergency hospitalization [8]. Acute pulpitis, periapical periodontitis, dental trauma, oral and maxillofacial infections have been recognized as emergency situations which force parents to take children to dental clinics [9].

The recommendations have been developed for dental offices which continue to work as dental emergency clinics during COVID-19 pandemic [10]. Main organizational aspects have been described: patients and staff examination for timely revealing coronavirus infection, protective equipment, patients' management, decision-making process, staff educating and training [11]. According to dental treatment methods it has been recommended: to restrict the use of high speed burs, ultrasonic scalers, intraoral radiographs; to prefer simple restorative materials (i.e. glass ionomer cements), tooth extraction rather than endodontic treatment; to use hand instruments and non-invasive methods where it is possible, to limit the operative time and to finish treatment in one appointment [12,13,14].

Self-isolation mode for citizens and the information through mass-media about new rules in dental clinics for patients (admission only for emergency including acute pain) have led to the decrease of dental patients' number. However, every day many children visit a dentist with emergency and urgent needs. Acute dental pain per se in children forces parents to take them to emergency dental clinics [15].

The aim of our study was to analyze the reasons for children visiting dental clinics during COVID-19 pandemic and to give the characteristics of applied dental treatment methods.

2. MATERIALS AND METHODS

The cross-sectional study was conducted in four dental clinics from 20.03.2020 to 24.04.2020 (one week in each clinic), which worked for dental emergency in the cities of Volgograd and Volzhky during the peak of COVID-19 pandemic. All the clinics organized

their work according to the current recommendations for the prevention of coronavirus spreading among the population. According to the Russian law the parents of the children under 15 years of age and adolescents older than 15 years gave written informed consents for dental examination and treatment.

The data about 166 paediatric dental patients were collected. The reasons for children's dental visits, age and behavior of the children, the tooth types, and the types of provided treatment were analyzed. Children's behavior was assessed according to Frankl's Behavior Rating Scale [16]. The proportions (%), 95% Confidence Interval (CI), mean-value with standard error ($M \pm m$) were calculated. The differences were assessed by Pierson's Chi-square test at p-value 0.05. Microsoft-office Excel-2019 programs were used for the statistic analysis.

3. RESULTS

In the circumstances of COVID-19 pandemic restrictions the majority of the children (164 of 166) visited dental clinics due to acute toothache and only 2 children due to acute oral mucosa disease (herpetic gingivostomatitis): 98.8% (95% CI 95.7-99.7%) and 1.2% (95% CI 0.3-4.3%) respectively. There were no cases of dental emergency due to periodontal disease.

An emergency visit to a dentist was the first dental visit in the current year for the majority of the children: 116 of 166 cases – 69.9% (95% CI 62.5-76.3%). However, approximately one of three children visited a dentist 1-3 months earlier due to toothache or routine caries treatment.

The number of boys and girls were similar: 85 and 81 cases – 51.2% и 48.8% respectively, $p > 0.05$. The age of the children was from 1 to 17 years, mean-value 7.6 ± 0.3 years (table 1). The children with acute pain in primary teeth were aged 1-12 years ($n=133$), in permanent teeth – 7-17 years ($n=31$).

The children with acute pain in primary teeth were aged 4-7 years (82 cases, 61.6%, 95% CI 53.2-69.5%) more often, than 8-10 years (28 cases, 21.0%, 95% CI 15.0-28.7%), 1-3 years (19 cases, 14.3%, 95% CI 9.3-21.2%), or 11-12 years (5 cases, 3.8%, 95% CI 1.6-8.5%).

The majority of the children with permanent teeth problems were older than 11 years: the same number (13 cases) in 11-14 and 15-17 year olds – 41.9% (95% CI 26.4-59.2%). Some children (4 cases) were in the age of 7-10 years: 12.9% (95% CI 5.1-28.8%).

One hundred and ninety teeth which needed therapeutic treatment or extraction were revealed in the children (table 2). The number of primary teeth ($n=159$) in need of urgent dental treatment was significantly higher than the number of permanent teeth ($n=31$): 83.7% (95% CI 77.8-88.3%) and 16.3% (95% CI 11.7-22.2%) respectively, $\chi^2 = 172.171$, $p < 0.0001$.

The first ($n=71$) and the second ($n=64$) primary molars were the most frequent teeth with acute pain among all primary teeth: 44.6% (95% CI 37.1-52.4%) and 40.2% (95% CI 32.9-48.0%) respectively. The numbers of incisors ($n=22$) and canines ($n=2$) were lower: 3.8% (95% CI 9.3-20.1%) и 1.3% (95% CI 0.3-4.5%). In total, the children complained of pain in upper primary teeth more often than in lower teeth: 92 and 67 cases, 57.9% (95% CI 50.1-65.3%) and 42.1% (95% CI 34.7-49.9%) respectively, $\chi^2 = 7.914$, $p = 0.0049$. However, the differences in every type of teeth were not significant statistically.

Acute toothache in the first permanent molars occurred more often than in other types of permanent teeth: in the first molars – 19 cases, 61.3% (95% CI 43.8-76.3%); the incisors – 5 cases, 16.1% (95% CI 7.1-32.6%); the second molars – 4 cases, 12.9% (95% CI 5.1-28.8%); premolars – 3 cases, 9.7% (95% CI 3.3-25.9%). The differences between the

frequencies of children's complaints about pain in upper and lower permanent teeth were not significant statistically.

The most common reasons for urgent dental treatment in children were caries consequences (table 3). The majority of the cases were caused by apical periodontitis or pulpitis: 98 and 67 cases, 51.6% (95% CI 44.5-58.6%) and 35.3% (95% CI 28.8-42.3%) respectively. Less often acute pain was caused by trauma – 12 cases, 6.3% (95% CI 3.6-10.7%); periostitis – 3 cases, 1.6% (95% CI 0.5-4.5%), primary tooth mobility due to natural root resorption – 9 cases, 4.7% (95% CI 2.5-8.8%) or deep caries – 1 case, 0.5% (95% CI 0.1-2.9%).

All (n=9) the primary teeth with pain due to their mobility were extracted.

Dental trauma was the cause of acute pain in 10 primary teeth and 2 permanent teeth. All the permanent teeth and 5 primary teeth required only conservative treatment. Every second primary tooth was extracted.

Intensive pain which accompanied pulpitis was revealed among permanent teeth significantly more often than among primary teeth: 17 of 31 teeth, 54.8% (95% CI 37.8-70.8%), and 50 of 159 teeth, 31.4% (95% CI 24.7-39.0%) respectively, $\chi^2 = 6.067$, $p = 0.0138$.

In contrast, apical periodontitis was more common among primary teeth than among permanent teeth: 89 of 159 teeth 56.0% (95% CI 48.2-63.5%), and 9 of 31 teeth, 29.0% (95% CI 16.1-46.6%) respectively, $\chi^2 = 7.695$, $p = 0.0055$.

In the outpatient clinic pulpitis treatment was impossible to provide in 2 of 67 (3.0%, 95% CI 0.8-10.2%) primary teeth due to lack of cooperation of the young patients. These patients were referred on the treatment under general anesthesia. In 2 cases (3.0%, 95% CI 0.8-10.2%) pulpitis treatment was performed with conscious sedation use. The applied pulpitis treatment methods were the following: 1) one-appointment vital therapy – direct pulp capping, pulpotomy, pulpectomy; 2) non-vital therapy with the application of a pulp necrotizing agent at the first appointment. Vital methods of pulpitis treatment were used more often than the methods with the use of a pulp necrotizing agent: 38 and 27 cases, 58.5% (95% CI 46.3-69.6%) and 41.5% (95% CI 30.4-53.7%) respectively. However, the difference was not significant statistically, $\chi^2 = 3.728$, $p = 0.0535$ (table 4). Local anesthesia was used in all cases of vital pulp therapy and in the most cases of non-vital pulp therapy (22 of 27 cases, 81.5%, 95% CI 63.3-91.8%).

Pulpotomy was applied in the primary teeth significantly more often than in the permanent teeth: 29 of 48 cases, 60.4% (95% CI 46.3-73.0%), and 2 of 17 cases, 11.8% (95% CI 3.3-34.3%) respectively, $\chi^2 = 11.703$, $p = 0.0006$. Pulpectomy in the primary teeth was significantly less common than in the permanent teeth: 1 of 48 cases, 2.1% (95% CI 0.4-10.9%), and 4 of 17 cases, 23.5% (95% CI 9.6-47.3%), respectively, $\chi^2 = 7.968$, $p = 0.0048$.

Almost all the primary teeth with apical periodontitis were extracted – 88 of 89 cases, 98.9% (95% CI 93.9-99.8%), (table 5). Apical periodontitis in the permanent teeth developed to periostitis in 3 of 12 cases (25.0%, 95% CI 8.9-53.2%). The endodontic treatment was performed in 5 permanent teeth (41.7%, 95% CI 19.3-68.0%), extraction – in 4 teeth (58.3%, 95% CI 31.9-80.7%) which were non-restorable.

Many children, who visited a dentist during COVID-19 pandemic, had negative behavior – 66 of 166 cases, 39.7% (95% CI 32.6-47.4%). The children with acute pain in the primary teeth had non-cooperative behavior significantly more often than the children with acute pain in the permanent teeth: 61 of 135 cases and 5 of 31 cases, 45.2% (95% CI 37.0-53.6%) и 16.1% (95% CI 7.1-32.6%) respectively, $\chi^2 = 10.603$, $p = 0.0011$. Different behavior management techniques were used for these 66 children: tell-show-do (54.5%, 95% CI 42.6-66.0%), sedation (3.0%, 95% CI – 0.8-10.4%), general anesthesia (3.0%, 95% CI – 0.8-10.4%), passive restraint (39.4%, 95% CI 28.5-51.4%). Treatment with passive

restriction, sedation or general anesthesia was provided only for children aged from 2 to 7 years.

4. DISCUSSION

COVID-19 pandemic due to high rate of mortality and transmission has changed the mode of dental service [17]. The dental procedures have been provided in the close contact with patients, can produce airborne particles and aerosols and place dental care workers in the high risk to coronavirus exposure [18,19]. In many countries routine dental care was suspended and only emergency and urgent dental care were allowed during the period of coronavirus pandemic peak [20,21,22]. High priority was given to relieving dental pain and to eliminating acute infection [23]. Many articles described the problems and new rules for dentistry which aimed at preventing infection spreading [24,25,26]. Special attention was paid to organization of children's appointments in dental setting and the implementation of minimally invasive concept in dental treatment [27,28]. However, there is lack of data about patients' characteristics and provided dental treatment in current circumstances of dental offices' work.

Before arising COVID-19 pandemic, dental pain and decayed teeth were the common reasons for the first children's visit to a dental clinic [29,30,31]. Rogers et al. examined 177 children from deprived families aged 4-18 years and revealed that 45% of children experienced dental pain. However, despite 95% caries prevalence, dmft=3.88 and DMFT=3.44, the children had low "Care index" (mft/dmft=0.1), which reflected a low level of oral care [32]. Slade GD showed that the rate of reported dental pain prevalence in children was connected with caries prevalence and varied from 5% to 33% in different countries [33]. Among paediatric patients of dental emergency clinics the main complaint (82-96.7%) was dental pain [34,35].

In our research we also revealed that the main reason for children to visit a dentist during coronavirus pandemic was dental pain (98.8%). It was found that the number of the children with acute dental pain in the primary teeth were significantly higher than in the permanent teeth. The most affected among the primary teeth were molars, among the permanent teeth – the first molar. These findings corresponded to high caries prevalence in primary dentition in Volgograd children [36], confirmed insufficient level of prevention and treatment of early childhood caries [37], low parents' care of their children's primary teeth [38]. On the other hand, these findings could be connected with asymptomatic caries development, rapid primary teeth destruction, quick spread of infection from dentine to pulp, and from pulp to periodontium. The lower number of urgent situations which was associated with caries complications in the permanent teeth could be explained by well-established school dentistry in the Volgograd region [39].

Martens et al. revealed that dental pain in emergency paediatric patients was caused by caries and its consequences (50.2%) or dental trauma (26.7%) [34]. Wong et al. reported about 74-75% of paediatric dental emergency which developed due to caries problems [40]. In our research the majority (88.4%) of the patients felt pain due to caries consequences and only 6.3% due to trauma. The decreased proportion of dental trauma could be connected with the prohibition of outdoor games, sport trainings and competitions for children during the self-isolation period caused by COVID-19 pandemic.

Among paediatric dental emergency patients 8-11% was preschoolers, which were referred for treatment under general anesthesia [40]. We revealed that among all the patients the number of children aged 1-3 years was 11.6%, 4-7 years – 50.6%. The majority of the children received dental treatment in outpatient clinics; only 2 children (1.2%) were referred to hospital for treatment under general anesthesia and the same number of the children was treated under conscious sedation with nitrous oxide. We suppose that the recommendations to

reduce aerosol dental procedures and to prevent children's hospitalization during coronavirus pandemic could increase the number of children treated without sedation or general anesthesia.

The choice of the method of pulpitis or apical periodontitis treatment is left to the judgment of a clinician because of many factors affecting the decision-making process. All current pulpitis treatment methods (pulp-capping, pulpotomy, pulpectomy) have sufficient clinical and radiographical results [41,42]. Our study revealed that dentists applied not only these methods but also the method with pulp-necrotizing agent use. The reason for this fact might be connected with the recommendations to limit the aerosol generating procedures and the duration of the treatment which could lower the probability of coronavirus infection. This reason could also explain the high level of primary teeth extraction due to apical periodontitis which was found out in our study. Meanwhile, apical periodontitis per se is the main reason for primary teeth extraction [43,44].

Dental fear and anxiety prevalence in children varies from 10% to 29% in different countries [45]. Wogelius et al. revealed high level (37.2%) of the history of behavior management problems in Danish 6-8 year old children who had dental treatment, although the anxiety prevalence in this population was only 5.7% [46]. In our study negative behavior at dental appointments was demonstrated by 39.7% children. This fact confirmed that the majority of pediatric patients during COVID-19 pandemic may be categorized as the patients with dental fear and anxiety who typically visit a dentist only when acute dental pain arises.

In anxious children different techniques of behavior problem management are usually applied: tell-show-do, nitrous-oxide sedation, passive restraint, oral premedication, handover-mouth, voice control, active restraint, and general anesthesia [47]. Luis de León et al. considered tell-show-do technique as the most acceptable by the parents and handover-mouth technique as the least acceptable one [47]. However, Al Zoubi et al. defined nitrous-oxide sedation as the most permissible technique (68.2% parents' answers) and passive restraint as the least acceptable technique (37.8% parents' answers) in emergency situations [48]. We found that in coronavirus pandemic circumstances the dentists used tell-show-do technique for behavior management in paediatric patients most often (54.5%), followed by passive restraint (39.4%).

5. CONCLUSIONS

In the limitation of this study we can conclude that COVID-19 pandemic changed the profile of paediatric dental patients and dentist's treatment options. In 1-17-year-old children the main reason for an emergency dental visit was acute dental pain due to caries consequences, pulpitis (35.3%) or apical periodontitis (51.6%). Every second patient was at the age from 4 to 7 years. The primary teeth needed urgent dental treatment significantly more often than the permanent teeth (83.7% and 16.3%, $p < 0.0001$). The children often demonstrated negative behavior and required behavior management. The applied treatment methods were aimed at limiting the use of aerosol generating procedures, decreasing the duration of the treatment and avoiding the patients' hospitalization.

6. REFERENCES

- [1] WHO Director-General's opening remarks at the media briefing on COVID-19 - 11 March 2020. <https://www.who.int/dg/speeches/detail/who-director-general-s-opening-remarks-at-the-media-briefing-on-covid-19---11-march-2020>
- [2] COVID-19 coronavirus pandemic <https://www.worldometers.info/coronavirus/>

- [3] He F, Deng Y, Li W. Coronavirus disease 2019: what we know? *J Med Virol.* 2020 Jul;92(7):719-725. PMID: 32170865 PMCID: [PMC7228340](#) DOI: [10.1002/jmv.25766](#)
- [4] Zimmermann M, Nkenke E. Approaches to the management of patients in oral and maxillofacial surgery during COVID-19 pandemic. *J Craniomaxillofac Surg.* 2020 May;48(5):521-526. PMID: 32303420 PMCID: [PMC7128256](#) DOI: [10.1016/j.jcms.2020.03.011](#)
- [5] Duruk G, Gümüşboğa ZŞ, Çolak C. Investigation of Turkish dentists' clinical attitudes and behaviors towards the COVID-19 pandemic: a survey study. *Braz Oral Res.* 2020;34:e054. PMID: 32490887 DOI: [10.1590/1807-3107bor-2020.vol34.0054](#)
- [6] Ge ZY, Yang LM, Xia JJ, FuXH, Zhang YZ. Possible aerosol transmission of COVID-19 and special precautions in dentistry. *J Zhejiang Univ Sci B.* 2020 May;21(5):361-368. PMID: 32425001 PMCID: [PMC7089481](#) DOI: [10.1631/jzus.B2010010](#)
- [7] Coulthard P. The oral surgery response to coronavirus disease (COVID-19). Keep calm and carry on? *Oral Surg.* 2020 doi: 10.1111/ors.12489. published online March 20. [[CrossRef](#)] [[Google Scholar](#)]
- [8] What constitutes a dental emergency? ADA. Updated 3/31/20. https://success.ada.org/~media/CPS/Files/Open%20Files/ADA_COVID19_Dental_Emergency_DDS.pdf?_ga=2.253879752.110187285.1584496315-1622146531.1565271894 Accessed June 14, 2020
- [9] Wang Y, Zhou CC, Shu R, Zou J. [Oral Health Management of Children During the Epidemic Period of Coronavirus Disease 2019]. *Sichuan Da Xue Xue Bao Yi Xue Ban.* 2020 Mar;51(2):151-154. [Article in Chinese] PMID: 32220180 DOI: [10.12182/20200360101](#)
- [10] Guidance for Dental Settings. CDC. 24/7. 19.05.2020. <https://www.cdc.gov/coronavirus/2019-ncov/hcp/dental-settings.html#Management> Accessed June 14, 2020
- [11] Panesar K, Dodson T, Lynch J, Bryson-Cahn C, Chew L, Dillon J. Evolution of COVID-19 guidelines for University of Washington Oral and Maxillofacial Surgery Patient Care. *J Oral Maxillofac Surg.* 2020 Apr 28;S0278-2391(20)30440-7. PMID: 32389541 PMCID: [PMC7194827](#) DOI: [10.1016/j.joms.2020.04.034](#)
- [12] Dave M, Seoudi N, Coulthard P. Urgent dental care for patients during the COVID-19 pandemic. *Lancet.* 2020 18-24 April; 395(10232): 1257. Published online 2020 Apr 3. PMID: [32251619](#) PMCID: [PMC7270877](#) doi: [10.1016/S0140-6736\(20\)30806-0](#)
- [13] Meng L, Hua F. Coronavirus disease 2019 (COVID-19): emerging and future challenges for dental and oral medicine. *J Dent Res.* 2020;12 [[PMC free article](#)] [[PubMed](#)] [[Google Scholar](#)]
- [14] Bali RK, Chaudhry K. Maxillofacial surgery and COVID-19, The Pandemic !! *J Maxillofac Oral Surg.* 2020 Jun; 19(2): 159–161. PMID: [32292253](#) PMCID: [PMC7148429](#) doi: [10.1007/s12663-020-01361-8](#)
- [15] Vanden Abbeele A, Vanhée T. [Emergencies in Paediatric Dentistry]. *Rev Med Brux.* 2018;39(4):325. [Article in French] PMID: 30320996
- [16] Frankl SN, Shiere FR, Fogels HR. Should the parent remain with the child in the dental operator? *J Dent Child.* 1962. 29:150-63. [[Google Scholar](#)]
- [17] Odeh ND, Babkair H, Abu-Hammad S, Borzangy S, Abu-Hammad A, Abu-Hammad O. COVID-19: Present and Future Challenges for Dental Practice. *Int J Environ Res Public Health.* 2020 Apr 30;17(9):3151. PMID: 32366034 PMCID: [PMC7246705](#) DOI: [10.3390/ijerph17093151](#)
- [18] Sabino-Silva R, Jardim ACG, Siqueira WL. Coronavirus COVID-19 impacts to dentistry and potential salivary diagnosis. *Clin Oral Investig.* 2020 Apr; 24 (4):1619-21. PMID: 32078048 PMCID: [PMC7088419](#) DOI: [10.1007/s00784-020-03248-x](#)

- [19] Alharbi A, Alharbi S, Alqaidi S. Guidelines for dental care provision during the COVID-19 pandemic. *Saudi Dent J.* 2020 Apr 7;32(4):181-186. PMID: 32292260 PMCID: [PMC7141449](#) DOI: [10.1016/j.sdentj.2020.04.001](#)
- [20] Coulthard P. Dentistry and Coronavirus (COVID-19) - Moral Decision-Making. *Br Dent J.* 2020 Apr;228(7):503-505. PMID: 32277203 DOI: [10.1038/s41415-020-1482-1](#)
- [21] Ahmed MA, Jouhar R, Ahmed N, Adnan S, Aftab M, Zafar MS, Khurshid Z Fear and Practice Modifications Among Dentists to Combat Novel Coronavirus Disease (COVID-19) Outbreak. *Int J Environ Res Public Health.* 2020 Apr 19;17(8):2821. PMID: 32325888 PMCID: [PMC7216192](#) DOI: [10.3390/ijerph17082821](#)
- [22] Pereira LJ, Pereira CV, Murata RM, Pardi V, Pereira-Dourado SM. Biological and social aspects of coronavirus disease 2019 (COVID-19) related to oral health. *Braz Oral Res.* 2020 May 8;34:e041. PMID: 32401931 DOI: [10.1590/1807-3107bor-2020.vol34.0041](#)
- [23] Berlin-Broner Y, Levin L. 'Dental hierarchy of needs' in the COVID-19 era - or why treat when it doesn't hurt? *Oral Health Prev Dent.* 2020;18(2):95. PMID: 32238979 https://ohpd.quintessenz.de/ohpd_2020_02_s0095.pdf
- [24] Barabari P, Moharamzadeh K. Novel coronavirus (COVID-19) and dentistry-a comprehensive review of literature. *Dent J (Basel).* 2020 May 21;8(2):E53. PMID: 32455612 DOI: [10.3390/dj8020053](#)
- [25] Spagnuolo G, De Vito D, Rengo S, Tatullo M. COVID-19 Outbreak: An Overview on Dentistry. *Int J Environ Res Public Health.* 2020 Mar; 17(6): 2094. PMID: 32235685 PMCID: [PMC7143628](#) DOI: [10.3390/ijerph17062094](#)
- [26] Fallahi HR, Keyhan SO, Zandian D, Kim SG, Cheshmi B. Being a front-line dentist during the Covid-19 pandemic: a literature review. *Maxillofac Plast Reconstr Surg.* 2020 Apr 24;42(1):12. PMID: 32341913 PMCID: [PMC7180678](#) DOI: [10.1186/s40902-020-00256-5](#)
- [27] American Academy of Pediatric Dentistry. <https://www.aapd.org/about/about-aapd/news-room/covid-19/>. Accessed June 14, 2020.
- [28] Mallineni SK, Innes NP, Raggio DP, Araujo MP, Robertson MD, Jayaraman J. Coronavirus disease (COVID-19): characteristics in children and considerations for dentists providing their care. *Int J Paediatr Dent.* 2020 May;30(3):245-250. PMID: 32250505 PMCID: [PMC7228382](#) DOI: [10.1111/ipd.12653](#)
- [29] Olatosi OO, Onyejaka NK, Oyapero A, Ashaolu JF, Abe A. Age and reasons for first dental visit among children in Lagos, Nigeria. *Niger Postgrad Med J.* Jul-Sep 2019;26(3):158-163. PMID: 31441453 DOI: [10.4103/NPMJ.NPMJ_60_19](#)
- [30] Daou MH, Eden E, El Osta N. Age and reasons of the first dental visit of children in Lebanon. *J Med Liban.* Jan-Mar 2016;64(1):18-22. PMID: 2716 9161 DOI: [10.12816/0023827](#)
- [31] Mika A, Mitus-Kenig M, Zeglen A, Drapella-Gasior D, Rutkowska K, Josko-Ochojska J. The Child's first dental visit. age, reasons, oral health status and dental treatment needs among children in Southern Poland. *Eur J Paediatr Dent.* 2018 Dec;19(4):265-270. PMID: 30567441 DOI: [10.23804/ejpd.2018.19.04.3](#)
- [32] Rogers HJ, Tariq U, Olsson L, Riaz SA, Miah MR. Caries prevalence, clinical consequences and self-reported pain experienced by children living in the West Bank. *Eur Arch Paediatr Dent.* 2019 Aug;20(4):333-338. PMID: 30911989 DOI: [10.1007/s40368-018-00412-6](#)
- [33] Slade GD. Epidemiology of dental pain and dental caries among children and adolescents. *Community Dent Health.* 2001 Dec;18(4):219-27. PMID: 11789699
- [34] Martens LC, Rajasekharan S, Jacquet W, Vandenbulcke JD, Van Acker JWG, Cauwels RGEC. Paediatric dental emergencies: a retrospective study and a proposal for

- definition and guidelines including pain management. *Eur Arch Paediatr Dent*. 2018 Aug;19(4):245-253. PMID: 29949083 DOI: [10.1007/s40368-018-0353-9](https://doi.org/10.1007/s40368-018-0353-9)
- [35] Shqair AQ, GomesGB, Oliveira A, Goettems ML, Romano AR, Schardozim LR, Bonow MLM, Torriani DD. Dental emergencies in a university pediatric dentistry clinic: a retrospective study. *Braz Oral Res*. Jan-Feb 2012;26(1):50-6. PMID: 22344338 DOI: [10.1590/s1806-83242012000100009](https://doi.org/10.1590/s1806-83242012000100009)
- [36] Osokina AS, Maslak EE. Early childhood caries prevalence and tooth surface distribution in Volgograd children. *Community Dental Health*. 2017. 34:S8-S9. doi:10.1922/CDH_EAPDHAAbstracts2017
- [37] Panchenko ML, Maslak EE, Dyachenko TS. Problems associated with providing dental care to children in the Volgograd region. *Journal of VolgSMU*. 2017. 3:130-132. DOI: [10.19163/1994-9480-2017-3\(63\)-130-132](https://doi.org/10.19163/1994-9480-2017-3(63)-130-132)
- [38] Onishchenko LF, Maslak EE, Kurkina ON. The results of parents' questioning on oral health in preschool children. *Dental Forum*. 2017. 4:64-65. [In Russ]. eLIBRARY ID: 30504310 <https://elibrary.ru/item.asp?id=30504310>
- [39] Maslak EE, Shkarin VV, Yanovskaya ML, Romanchuk EV, Luneva NA. Need for school dentistry and results of schoolchildrens dental care in Volgograd. *Annali di stomatologia*. 2011. T. 2. № S1-2. C. 12. <https://www.annalidistomatologia.com/common/php/portiere.php?ID=fef866e89c701497c262f9397c6a79b3>
- [40] Wong NHY, Tran C, Pukallus M, Holcombe T, Seow WK. A three-year retrospective study of emergency visits at an oral health clinic in South-East Queensland. *Aust Dent J*. 2012 Jun;57(2):132-7. PMID: 22624751 DOI: [10.1111/j.1834-7819.2012.01688.x](https://doi.org/10.1111/j.1834-7819.2012.01688.x)
- [41] Smaïl-Faugeron V, Glenny AM, Courson F, Durieux P, Muller-Bolla M, Fron Chabouis H. Pulp treatment for extensive decay in primary teeth. *Cochrane Database Syst Rev*. 2018 May 31;5(5):CD003220. PMID: 29852056 PMCID: [PMC6494507](https://pubmed.ncbi.nlm.nih.gov/PMC6494507/) DOI: [10.1002/14651858.CD003220.pub3](https://doi.org/10.1002/14651858.CD003220.pub3)
- [42] Gadallah L, Hamdy M, El Bardissy A, Abou El Yazeed M. Pulpotomy versus pulpectomy in the treatment of vital pulp exposure in primary incisors. A systematic review and meta-analysis. *F1000Res*. 2018 Sep 26;7:1560. PMID: 31249668 PMCID: [PMC6584970](https://pubmed.ncbi.nlm.nih.gov/PMC6584970/) DOI: [10.12688/f1000research.16142.3](https://doi.org/10.12688/f1000research.16142.3)
- [43] Bansal M, Gupta N, Gupta P., Arora V., Thakar S. Reasons for extraction in primary teeth among 5-12 years school children in Haryana, India - A cross-sectional study. *J Clin Exp Dent*. 2017. 9(4):e545-e549. DOI: [10.4317/jced.53076](https://doi.org/10.4317/jced.53076)
- [44] Ockell NM, Bågesund M. Reasons for extractions, and treatment preceding caries-related extractions in 3-8 year-old children. *Eur Arch Paediatr Dent*. 2010. 11(3):122-30. DOI: [10.1007/bf03262727](https://doi.org/10.1007/bf03262727)
- [45] Cianetti S, Lombardo G, Lupatelli E, Pagano S, Abraha I, Montedori A, Caruso S, Gatto R, De Giorgio S, Salvato R. Dental Fear/Anxiety Among Children and Adolescents. A Systematic Review. *Eur J Paediatr Dent*. 2017 Jun;18(2):121-130. PMID: 28598183 DOI: [10.23804/ejpd.2017.18.02.07](https://doi.org/10.23804/ejpd.2017.18.02.07)
- [46] Wogelius P, Poulsen S, Sørensen HT. Prevalence of dental anxiety and behavior management problems among six to eight years old Danish children. *Acta Odontol Scand*. 2003 Jun;61(3):178-83. PMID: 12868693 DOI: [10.1080/00016350310003468](https://doi.org/10.1080/00016350310003468)
- [47] Luis de León J, Guinot Jimeno F, Bellet Dalmau LJ. Acceptance by Spanish parents of behaviour-management techniques used in paediatric dentistry. *Eur Arch Paediatr Dent*. 2010 Aug;11(4):175-8. PMID: 20840827 DOI: [10.1007/BF03262739](https://doi.org/10.1007/BF03262739)
- [48] Al Zoubi L, Schmoeckel J, Mustafa Ali M, Alkilzy M, Splieth CH. Parental acceptance of advanced behaviour management techniques in normal treatment and in emergency situations used in paediatric dentistry. *Eur Arch Paediatr Dent*. 2019 Aug;20(4):319-323. PMID: 30600475 DOI: [10.1007/s40368-018-0408-y](https://doi.org/10.1007/s40368-018-0408-y)

Table 1. Age distribution of the children who visited dental clinics with acute toothache in primary or permanent teeth

Age, years	The number of visits due to acute pain in:					
	primary teeth		permanent teeth		total	
	n	% (95% CI)	n	% (95% CI)	n	% (95% CI)
1	1	0.7 (0.1-4.1)			1	0.6 (0.1-3.4)
2	5	3.8 (1.6-8.5)			5	3.0 (1.3-6.9)
3	13	9.8 (5.8-16.0)			13	7.9 (4.7-13.1)
1-3	19	14.3 (9.3-21.2)			19	11.6 (7.5-17.4)
4	18	13.5(8.7-20.4)			18	11.0 (7.1-16.7)
5	22	16.5 (11.2-23.8)			22	13.4 (9.0-19.5)
6	23	17.3 (11.8-24.6)			23	14.0 (9.5-20.2)
7	19	14.3 (9.3-21.2)	1	3.2 (0.6-16.2)	20	12.2 (8.0-18.1)
4-7	82	61.6 (53.2-69.5)	1	3.2 (0.6-16.2)	83	50.6 (43.0-58.2)
8	9	6.8 (3.6-12.4)	2	6.4 (1.8-20.7)	11	6.7 (3.8-11.6)
9	11	8.3 (4.7-14.2)	1	3.2 (0.6-16.2)	12	7.3 (4.2-12.4)
10	8	6.0 (3.1-11.4)	0	0.0 (0.0-11.0)	8	4.9 (2.5-9.3)
8-10	28	21.0 (15.0-28.7)	3	9.7 (3.3-24.9)	31	18.9 (13.6-25.5)
11	3	2.3 (0.8-6.4)	4	12.9 (5.1-28.8)	7	4.3 (2.1-8.5)
12	2	1.5 (0.4-5.3)	2	6.4 (1.8-20.7)	4	2.4 (0.9-6.1)
13			3	9.7 (3.3-24.9)	3	1.8 (0.6-5.2)
14			4	12.9 (5.1-28.8)	4	2.4 (0.9-6.1)
11-14	5	3.8 (1.6-8.5)	13	41.9 (26.4-59.2)	18	11.0 (7.1-16.7)
15			2	6.4 (1.8-20.7)	2	1.2 (0.3-4.3)
16			6	19.3 (9.2-36.3)	6	3.6 (1.7-7.7)
17			5	16.1 (7.1-32.6)	5	3.0 (1.3-6.9)
15-17			13	41.9 (26.4-59.2)	13	7.9 (4.7-13.1)
Total	133	100	31	100	164	100
	133	81.1 (74.4-86.3)	31	18.9 (13.6-25.6)	164	100

Table 2. Types of teeth with acute toothache

Types of teeth	The number and percentage of the cases					
	Upper teeth		Lower teeth		Total	
	n	% (95% CI)	n	% (95% CI)	n	% (95% CI)
Primary teeth	92	57.9 (50.1-65.3)	67	42.1 (34.7-49.9)	159	100
incisors	20	12.6 (8.3-18.6)	2	1.3 (0.3-4.5)	22	13.8 (9.3-20.1)
canines	2	1.3 (0.3-4.5)	0	0.0 (0.0-2.4)	2	1.3 (0.3-4.5)
first molars	43	27.0 (20.7-34.4)	28	17.6 (12.5-24.3)	71	44.6 (37.1-52.4)
second molars	27	17.0 (11.9-23.6)	37	23.3 (17.4-30.4)	64	40.2 (32.9-48.0)
Permanent teeth	17	54.8 (37.8-70.8)	14	45.2 (29.2-62.2)	31	100
incisors	4	12.9 (5.1-28.8)	1	3.2 (0.6-16.2)	5	16.1 (7.1-32.6)
canines	0	0.0 (0.0-11.0)	0	0.0 (0.0-11.0)	0	0.0 (0.0-11.0)
premolars	2	6.4 (1.8-20.7)	1	3.2 (0.6-16.2)	3	9.7 (3.3-25.9)
first molars	7	22.6 (11.4-39.8)	12	38.7 (23.7-56.2)	19	61.3 (43.8-

second molars	4	12.9 (5.1-28.8)	0	0.0 (0.0-11.0)	4	76.3 12.9 (5.1-28.8)
---------------	---	-----------------	---	----------------	---	-------------------------

Table 3. The reasons for urgent dental treatment in children

Reasons for treatment	The number and percentage of the cases							
	Primary teeth		Permanent teeth		Chi-square test		Total	
	n	% (95% CI)	n	% (95% CI)	x ²	p-value	n	% (95% CI)
Caries	1	0.6 (0.1-3.6)	0	0.0 (0.0-11.0)	NA*		1	0.5 (0.1-2.9)
Pulpitis	50	31.4 (24.7-39.0)	17	54.8 (37.8-70.8)	6.193	0.0128	67	35.3 (28.8-42.3)
Apical periodontitis	89	56.0 (48.2-63.5)	9	29.0 (16.1-46.6)	7.533	0.0061	98	51.6 (44.5-58.6)
Acute periostitis	0	0.0 (0.0-2.4)	3	9.7 (3.3-24.9)	NA*		3	1.6 (0.5-4.5)
Acute dental trauma	10	6.3 (3.4-11.2)	2	6.4 (1.8-20.7)	NA*		12	6.3 (3.6-10.7)
Primary teeth mobility	9	5.7 (3.0-10.4)	x	x	x	x	9	4.7 (2.5-8.8)
Total	159	100	31	100	-	-	190	100
	159	83.7 (77.8-88.3)	31	16.3 (11.7-22.2)	172.171	<0.0001	190	100

*NA – not applicable

Table 4. The methods of pulpitis treatment

Methods	The number and percentage of teeth				Chi-square test		Total	
	primary		permanent		x ²	p-value	n	% (95% CI)
	n	% (95% CI)	n	% (95% CI)				
Vital therapy	31	64.6 (50.4-76.6)	7	41.2 (21.6-64.0)	2.787	0.0950	38	58.5 (46.3-69.6)
pulp capping	1	2.1 (0.4-10.9)	1	5.9 (1.0-27.0)	NA*		2	3.1 (0.8-10.5)
pulpotomy	29	60.4 (46.3-73.0)	2	11.8 (3.3-34.3)	11.703	0.0006	31	47.7 (36.0-59.6)
pulpectomy	1	2.1 (0.4-10.9)	4	23.5 (9.6-47.3)	7.968	0.0048	5	7.7 (3.3-16.8)
Pulp-necrotizing agent application	17	35.4 (23.4-49.6)	10	58.8 (36.0-78.4)	2.787	0.0950	27	41.5 (30.4-53.7)
Total	48	100	17	100			65	100
	48	73.8 (62.0-	17	26.1 (17.0-	29.351	<0.0001	65	100

	83.0)	37.9)	1
--	-------	-------	---

*NA – not applicable

Table 5. The methods of apical periodontitis treatment

Methods	The number and percentage of teeth				Chi-square test		Total		
	primary			permanent		x ²	p-value	n	% (95% CI)
	n	% (95% CI)	n	% (95% CI)					
Endodontic treatment	1	1.1 (0.2-6.1)	5	41.7 (19.3-68.0)	30.967	< 0.0001	6	5.9 (2.7-12.4)	
Extraction	88	98.9 (93.9-99.8)	7	58.3 (31.9-80.7)	30.967	< 0.0001	95	94.1 (87.6-97.2)	
Total	89	100	12	100			101	100	
	89	88.1 (80.4-93.1)	12	11.9 (6.9-19.6)	116.709	< 0.0001	101	100	

Kamalova Mekhriniso Kilichevna., Doctor of Philosophy of Medical Sciences, Head of the Department of Surgical Dentistry, Bukhara State Medical Institute of the Ministry of Health of Uzbekistan.

Address: 200118, Bukhara, st. A. Navoi. Tel 8 (365) -223-00-50.

E-mail mexriniso.stomatolog@mail.ru

ORCID: [0000-0002-8540-376X](https://orcid.org/0000-0002-8540-376X)

Fomenko Irina V., Doctor of Medicine, Head of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 902 387 34 48.

E-mail fomenira@mail.ru

ORCID: [0000-0002-5279-6106](https://orcid.org/0000-0002-5279-6106)

Dmitrienko Dmitriy S. Doctor of Medicine, Professor of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 905 336 89 54.

E-mail vsp79@mail.ru

ORCID: [0000-0002-9555-6612](https://orcid.org/0000-0002-9555-6612)

Matvienko Natalya V. Candidate of Medical Sciences, Associate Professor of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

E-mail kuyuma@yandex.ru

ORCID: [0000-0002-3913-5725](https://orcid.org/0000-0002-3913-5725)

Arjenovskaya Evgenia N. PhD student of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia. Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 905 336 89 54.

E-mail acontario@rambler.ru

ORCID: [0000-0001-5300-6498](https://orcid.org/0000-0001-5300-6498)

Gevorkyan Areknazan G. Clinical resident of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 902 387 34 48.

E-mail areknazan_gevorkyan@mail.ru

ORCID: [0000-0001-8153-8686](https://orcid.org/0000-0001-8153-8686)

Nikitina Kseniya V. Clinical resident of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 902 387 34 48.

E-mail kseniya-bonya@mail.ru

ORCID: [0000-0003-3439-7440](https://orcid.org/0000-0003-3439-7440)

Maslak Elena E., Doctor of Medicine, Professor of Pediatric Dentistry Department, Volgograd State Medical University, Ministry of Health of Russia.

Address: 400131, Volgograd, Pl. Pavshikh Bortsov 1. Tel. +7 905 336 89 54.

E-mail emaslak@yandex.ru

ORCID: [0000-0003-2011-9714](https://orcid.org/0000-0003-2011-9714)