

Original research article

A retrospective analysis to estimate the prevalence of anemia in exclusively breastfed babies between 3-6 months of age

Dr. Pankaj Kumar Kashyap¹, Dr. A.K. Jaiswal²

¹ Senior Resident, Upgraded Department of Paediatrics, Patna Medical College & Hospital, Patna, Bihar, India.

² Professor & HOD, Upgraded Department of Paediatrics, Patna Medical College & Hospital, Patna, Bihar, India

Corresponding Author: Dr. Pankaj Kumar Kashyap

Abstract

Aim: to estimate the prevalence of anemia in exclusively breastfed babies between 3-6 months of age so that we can identify, prevent and treat anemia and its consequences at an early age.

Material and methods: This retrospective observational study was done the Upgraded Department of Paediatrics, Patna Medical College & Hospital, Patna, Bihar, India, from feb 2017 to July 2018. Exclusive breastfed term infants aged between 3-6 months of age (90 days to 180 days) who undergone complete blood count (CBC) examination for some minor ailments was included in this study. All the infants were looked for anemia and its severity.

Results: Total 550 infants aged 3-6 months attended OPD during study period. Out of them, 150 infants had their CBC done for various reasons and for pallor. After excluding 40 babies as per our exclusion criteria, 110 infants matched the selection criteria and remained as study group. Among 110 infants, 65 were males and 45 were females. According to age 35, 47 and 28 babies were belonging to 3-4, 4-5 and 5-6 months respectively. Mean age was 4.5 month and Hb was 9.9 mg/dl. Median Hb was 9.6 mg/dl. A total of 92 children had anemia as per WHO criteria of <11 gm% of Hb, giving a prevalence of anemia of 83.64%. However if we take 10.5 mg% as cut off value 85 (77.27%) had anemia and for 10 mg%, 69 (62.73%) had anemia. Out of 92, 31 (88.57%) babies with age of 3-4 months, 40 (85.11%) aged 4-5 months and 21 (75%) aged 5-6 months had anemia. out of 92, 62 males and 30 females were anemic.

Conclusion: we concluded that the exclusively breastfed infants between 3-6 months are at increased risk of anemia. Therefore infants after 3 months, should be evaluated for anemia and iron deficiency which is the commonest cause of anemia. Such infants should be supplemented with oral iron in addition to exclusive breast feeding for 6 months, to prevent adverse effects of IDA on infants' growth and development.

Keywords: Anemia, Exclusive breast feeding, Full-term baby, Prevalence

Introduction

The World Health Organization (WHO) and American Academy of Pediatrics unequivocally recommend that exclusive breastfeeding is the ideal nutrition for infants and is sufficient to support optimal growth for the first six months of life.^{1,2} However, there is controversy about the adequacy of breast milk in maintaining optimum iron status of exclusively breastfed babies. A WHO committee expressed concern that some exclusively breastfed infants may become iron deficient.¹ Glader recommended that infants exclusively breastfed should receive iron supplementation from four months of age.³ Calvo et al. evaluated the iron and nutritional status of exclusively breastfed infants for a prolonged period in relation to their growth rate and dietary changes and recommended that breastfed infants should be given supplemental iron from the fourth month of life.⁴ Dewey et al. also evaluated the effect of introducing complementary foods before six months of age in exclusively breastfed infants in Honduras.⁵

They recommended iron drops for breastfed infants with birth weights between 2500 g and 3000 g. But McMillan et al. reported that term breastfed infants did not need supplemental iron until the birth weight tripled, which occurred at about 12 months of age.⁶ Similarly Owen et al. found that infants breastfed until 20 weeks of life, had sufficient iron stores at 6 months of age.⁷ Zavaleta et al. reported an interesting observation that maternal anemia did not affect breast milk iron or lactoferrin concentration at birth and during early lactation.⁸ The transfer of iron from the mother to the fetus is supported by a substantial increase in maternal iron absorption and is regulated by the placenta when the receptors for iron located on the apical surface of placental syncytiotrophoblast gradually increases near term. The capacity of this system may be inadequate to maintain iron transfer to the fetus when the mother is iron deficient. Maternal iron deficiency during pregnancy may hamper the development of fetal iron stores prior to birth, and perhaps well into the first year of life also and therefore increasing the risk of anemia during infancy. This has adverse consequences on the neurological development of these infants. Iron deficiency anemia in pregnancy leads to decreased placental weight and significantly reduced number of placental cotyledons and thus is a risk factor for preterm delivery, intra uterine growth retardation and low birth weight and neonatal mortality. However mounting evidences suggest that even though infants of anemic mothers are born with adequate birth weight, they have low iron stores and are more likely to develop anemia.⁹⁻¹¹ In fact, the infants risk of having low hemoglobin when their birth weight was normal (>2500g) but mother was anemic (<11gm/dl) was greater than the infants who had a low birth weight (<2500g) and were born to non-anemic mother(>11g/dl). So the present study was conducted to estimate the prevalence of anemia in exclusively breastfed babies between 3-6 months of age so that we can identify, prevent and treat anemia and its consequences at an early age.

Material and methods

This retrospective observational study was done the Upgraded Department of Paediatrics, Patna Medical College & Hospital, Patna, Bihar, India, from feb 2017 to July 2018, after taking the approval of the protocol review committee and institutional ethics committee.

Inclusion criteria

Exclusive breastfed term infants aged between 3-6 months of age (90 days to 180 days) who attended OPD of the hospital and had undergone complete blood count (CBC) examination for some minor ailments. According to AAP, CBC is not advised routinely in all infants unless there is clinical suspicion of anemia and risk factors for anemia. So CBC was not done for all infants but done for those who presented with pallor and some illnesses.¹²

Exclusion criteria

Children born prematurely, intrauterine growth retarded babies, those with repeated infections, severe infections, babies with supplements, bottle feeds and haemoglobinopathies were excluded from the study.

Methodology

The patients' data and Hb levels were collected from the hospital software. All the infants were looked for anemia and its severity. Since, there is no established cut off available for hemoglobin concentration to diagnose anemia in infants <6 months, the WHO definition of anemia for infants above 6 months, i.e. hemoglobin concentration <11 gm/dl was used to diagnose anemia for babies between 3-6 months and it was also used to assess for severity; severe <7 mg/dl, moderate 7.00-8.9 mg/dl and mild 9.00-10.9 mg/dl.^{13,14} In addition, the cut-off values suggested by Domelloff et al (<10.5 mg/dl) and Lonnerdal et al (<10 mg/dl) were

also used.^{15,16} So the infants were assessed for all 3 cut-off values i.e. <10 mg/dl, <10.5 mg/dl and <11 mg/dl.

Results

Totally 550 infants aged 3-6 months attended OPD during study period. Out of them, 150 infants had their CBC done for various reasons and for pallor. After excluding 40 babies as per our exclusion criteria, 110 infants matched the selection criteria and remained as study group. Out of 110 infants in the study group, 17 had URI with pallor, 45 had ALRI and another 15 had gastroenteritis with pallor. Three each presented with UTI, Febrile convulsions, Rickettsial fever, septicemia and laryngomalacia. Remaining 18 children had different minor illnesses but none had severe life threatening illness.

Among 110 infants, 65 were males and 45 were females. According to age 35, 47 and 28 babies were belonging to 3-4, 4-5 and 5-6 months respectively. Mean age was 4.5 month and Hb was 9.9 mg/dl. Median Hb was 9.6 mg/dl.

Table 1: Demographic profile of patients

Gender	Number of patients	Percentage
Male	65	59.10
Female	45	40.90
Age in months		
3-4 months	35	31.82
4-5 months	47	42.73
5-6 months	28	25.45

Table 2: Prevalence of anemia based on Hb level

Hb level	Number of patients	Percentage
<11 gm%	92	83.64
10.5 mg%	85	77.27
10 mg%,	69	62.73

A total of 92 children had anemia as per WHO criteria of <11 gm% of Hb, giving a prevalence of anemia of 83.64%. However if we take 10.5 mg% as cut off value 85 (77.27%) had anemia and for 10 mg%, 69 (62.73%) had anemia.

Table 3: Age wise distribution of patients

Age in months	Total no. of patients	Anemic patients	Percentage
3-4 months	35	31	88.57
4-5 months	47	40	85.11
5-6 months	28	21	75

Out of 92, 31 (88.57%) babies with age of 3-4 months, 40 (85.11%) aged 4-5 months and 21 (75%) aged 5-6 months had anemia (table 3).

Table 4: Gender wise distribution of patients

Gender	Number of patients	Anemic patients	Percentage
Male	65	62	95.38
Female	45	30	66.67

Segregating by gender, 62 males and 30 females were anemic (table 4).

Table 5: Severity of anemia

Severity of anemia	Number of patients	Percentage
Mild anemia	66	71.74
Moderate anemia	22	23.91
Severe anemia	4	4.35

On classifying anemia based on severity as per WHO criteria for assessing severity of anemia in infants above 6 months, it was found that 66 had mild, 22 had moderate and 4 had severe anemia (table 5).

Discussion

In our study, the prevalence of anemia was 83.64%. A study conducted by Hemachitra et al to assess the prevalence of anemia in 3-6 months old babies with a similar cut off value of Hb showed a prevalence of 65.8%.¹⁷ In Asia, the prevalence of anemia in children under 2 years of age may exceed 90%.¹⁸ The prevalence of anemia among full-term infants may be as high as 80% at 3-6 months and 90% at 6-9 months of age.¹⁹

The extremely high prevalence of anemia in absolutely breast fed infants between 3-6 months of age is similar to the prevalence of anemia observed in children above 6 months. According to NHFS-III, 79% children under 3 years were anemic and in 6-23 months, 81.8% were anemic.¹⁸ In another study conducted in India, 93.5% prevalence of anemia in children aged between 6 to 23 months.²⁰ A study conducted in Odisha yielded 94% prevalence of anemia in under 5 year children.²¹ In a study conducted in Burma, the prevalence of anemia in children between 6-36 months was 72.65%.²²

We tried to find out the cause of anemia in the study group. Babies had no evidence of hookworm infestation, malaria, recurrent diarrhea, recurrent infections and malnutrition which can cause anemia in this age group. Alpha thalassemia trait is uncommon in this area.

A number of studies, conducted largely among infants aged 6-12 months, found that infants born to anemic mothers had a lower Hb concentration.^{23,24} Eighty percent of the iron present in a newborn term infant is accumulated during the third trimester of pregnancy.²⁵ Hemachitra et al, in their study confirmed that low haemoglobin and poor weight gain in term 3-6 months old babies were significantly associated with maternal anemia in last trimester.¹⁷ In support of this, various studies conducted to know the prevalence of anemia in mothers around this region showed high prevalence of anemia in pregnant mothers. Prevalence of anemia in antenatal mothers was 73% in Belgaum and 96.5% in Koppal, Karnataka India, whereas it was 98% in Sonapat Haryana India.²⁶⁻²⁸

Recently it has also been proved by different studies that time of umbilical cord clamping at the time of delivery also affects Hb levels in infants. These studies have reported that late-timing of cord clamping, might be associated with better hemoglobin values, higher stores of iron at 6 months of age and lower incidence of anemia.²⁹⁻³¹ Delayed umbilical cord clamping (approximately 120 to 180 seconds after delivery) is associated with improved iron status (ferritin levels) at two to six months of age.^{32,33}

As per previous literature available, it was understood that the breastfeeding protected children from ID/IDA until the 4th month of age.^{25,34} So surveillance is required by the 4th month after birth in order to identify children in need of iron supplementation in fully breastfed babies when they reach 4 months instead of 6 months of age.²⁹ Therefore it is recommended that exclusive breastfed term infants receive an iron supplementation of 1 mg/kg per day, starting at 4 months of age and continued until appropriate iron-containing complementary foods have been introduced.²⁵ However in our study, Out of 92, 31 (88.57%) babies with age of 3-4 months, 40 (85.11%) aged 4-5 months and 21 (75%) aged 5-6 months had anemia. This suggests that anemia is common even at 3 months of age necessitating iron supplementation at 3 months. Even though the study population in the present study is biased towards sick children, high prevalence value is unexpected and alarming. One of the reasons for higher prevalence of

anemia in the present study could be the high prevalence of maternal anemia in this region. The timing of cord clamping could not be assessed in our study. The other factors could be, majority of the children were poor, rural and from low socioeconomic status.^{35,36} Low socioeconomic status can play a role in different ways, including poorer nutritional status of the mother, maternal anemia and hence poorer fetal nutrition.^{13,16} One more reason is delayed initiation of iron supplementation.

Conclusion

We concluded that the exclusively breastfed infants between 3-6 months are at increased risk of anemia. Therefore infants after 3 months, should be evaluated for anemia and iron deficiency which is the commonest cause of anemia. Such infants should be supplemented with oral iron in addition to exclusive breast feeding for 6 months, to prevent adverse effects of IDA on infants' growth and development.

Reference

1. World Health Organization: The Optimal Duration of Exclusive Breastfeeding. 2001 [<http://www.who.int/inf-pr-2001/en/note2001-07.html>].
2. American Academy of Pediatrics Work Group on Breastfeeding: Breastfeeding and the use of human milk. *Pediatrics* 1997, 100:1035-1039.
3. Glader B: Anemias of inadequate production. In Nelson Textbook of Pediatrics 17th edition. Edited by: Behrman, Kliegman, Jenson. Philadelphia: Saunders; 2004:1606-1617.
4. Calvo EB, Galindo AC, Aspres NB: Iron status in exclusively breast-fed infants. *Pediatrics* 1992, 90:375-379.
5. Dewey KG, Cohen RJ, Rivera LL, Brown KH: Effects of age of introduction of complementary foods on iron status of breast-fed infants in Honduras. *Am J Clin Nutr* 1998, 67:878-884.
6. McMillan JA, Landaw SA, Oski FA: Iron sufficiency in breastfed infants and availability of iron from human milk. *Pediatrics* 1976, 58:686-691.
7. Owen GM, Garry PJ, Hooper EM, Gilbert BA, Pathak D: Iron nutriture of infants exclusively breastfed the first five months. *J Pediatr* 1981, 99:237-240.
8. Zavaleta N, Nombera J, Rojas R, Hambraeus L, Gislason J, Lonnerdal B: Iron and lactoferrin in milk of anemic mothers given iron supplements. *Nutr Res* 1995, 15:681-690
9. Ahamad S, Amir M, Ansari Z, Ahmed KN. Influence of maternal iron deficiency anemia on the fetal total body iron. *Indian Pediatr*. 1983;20:643-6.
10. Pasricha S. Determinants of Anemia among Young children in Rural India. *Am Acad Pediatr*. 2010;126:e140-9.
11. Dallman PR, Siimes MA, Stekel A. Iron deficiency in infancy and childhood. *Am J Clin Nutrition* 1980;33:86-118.
12. Wang M. Iron deficiency and other types of anemia in infants and children. *Am Fam Phys*. 2016;93(4):270-8.
13. De Pee S, Bloem MW, Sari M, Kiess L, Yip R, Kosen S. The high prevalence of low hemoglobin concentration among Indonesian infants aged 3-5 months is related to maternal anemia. *J Nutr*. 2002;132(8):2215-21.
14. World Health Organization. Haemoglobin concentrations for the diagnosis of anaemia and assessment of severity. 2011. Available at: <https://apps.who.int/iris/handle/10665/85839>. Accessed on 12 January 2021.

15. Domellöf M, Cohen RJ, Dewey KG, Hernell O, Rivera LL, Lönnerdal B. Iron supplementation of breast-fed Honduran and Swedish infants from 4 to 9 months of age. *J Pediatr*. 2001;138(5):679-87.
16. Dhanasekaran R, Sumitha A, Suguna. Impact of maternal anaemia on cord blood haemoglobin. *Int J Contemp Pediatr* 2019;6:1235-8.
17. Hemachitra J, Monish A. Risk of infant anemia in 3-6 months old babies and its association with maternal anemia. *Int J Contemp Pediatr*. 2018;5(3):938.
18. Kotecha PV. Nutritional anemia in young children with focus on Asia and India. *Indian J Community Med*. 2011;36(1):8-16.
19. Lutter CK. Iron deficiency in young children in low- income countries and new approaches for its prevention. *J Nutr*. 2008;138:2523-8.
20. Marol RN, Marol RR, Marol RR. Prevalence of anemia in children with acute lower respiratory tract infection: a case control study in children between 6 months to 23 months. *Int J Contemp Pediatr*. 2020;7:1573-7.
21. Sahu T, Sahani NC, Patnaik L. Childhood anemia- a study in tribal area of Mohana block in Orissa. *Indian J Community Med*. 2007;32(1):43-5
22. Black MM, Quigg AM, Hurley KM, Pepper MR. Iron deficiency and iron-deficiency anemia in the first two years of life: strategies to prevent loss of developmental potential. *Nutr Rev*. 2011;69(1):S64- 70.
23. Strauss MB. Anemia of infancy from maternal iron deficiency in pregnancy. *J Clin Invest*. 1933;12(2):345-53.
24. Colomer J, Colomer C, Gutierrez D, Jubert A, Nolasco A, Donat J, et al. (1990) Anaemia during pregnancy as a risk factor for infant iron deficiency: report from the Valencia Infant Anaemia Cohort (VIAC) study. *Paediatr Perinat Epidemiol*. 1990;4:196-204.
25. Baker RD, Greer FR, Committee on Nutrition American Academy of Pediatrics. Diagnosis and prevention of iron deficiency and iron-deficiency anemia in infants and young children (0-3 years of age). *Pediatrics*. 2010;126(5):1040-50.
26. Prashant D. Prevalence of anemia among pregnant women attending antenatal clinics in rural field practice area of Jawaharlal Nehru Medical College, Belagavi, Karnataka, India. *Int J Community Med Public Health*. 2017;4(2):537.
27. Seema BN. Prevalence of anemia among pregnant women in rural Koppal: a study from teaching hospital, Koppal, India. *Int J Reprod Contracept Obstet Gynecol*. 2017;6(9):3792-5.
28. Mangla M, Singla D. Prevalence of anaemia among pregnant women in rural India: a longitudinal observational study. *Int J Reprod Contracept Obstet Gynecol*. 2016 Dec 15;5(10):3500-5
29. Marques RF, Taddei JA, Lopez FA, Braga JA. Breastfeeding exclusively and iron deficiency anemia during the first 6 months of age. *Revista da Associação Médica Brasileira*. 2014;60(1):18-22.
30. Chaparro CM, Neufeld LM, Alavez GT, Cedillo REL, Dewey KG. Effect of timing of umbilical cord clamping on iron status in Mexican infants: a randomized controlled trial. *Lancet*. 2006;367:1997- 2004.
31. Van Rheenen PF, Brabin BJ. A practical approach to timing cord clamping in resource poor settings. *BMJ*. 2006;333:954-8.
32. Hutton EK, Hassan ES. Late versus early clamping of the umbilical cord in full-term neonates: systematic review and meta-analysis of controlled trials. *JAMA*. 2007;297(11):1241-52.
33. Andersson O, Hellström-Westas L, Andersson D, Domellöf M. Effect of delayed versus early umbilical cord clamping on neonatal outcomes and iron status at 4 months: a randomised controlled trial. *BMJ*. 2011;343:d7157

34. Dube K, Schwartz J, Mueller MJ, Kalhoff H, Kersting M. Iron intake and iron status in breastfed infants during the first year of life. *Clin Nutr.* 2010;29:773-8
35. WHO. The global prevalence of anemia in 2011. World Health Organization. Geneva: Switzerland; 2015
36. Census Data 2001- Census of India. Available at: https://censusindia.gov.in/2011-common/census_data_2001.html. Accessed on 10 November 2020.