

Original research article

Hyperhomocysteinemia in pregnancy induced hypertension: a case-control study

Dr. Anupam Kumari¹, Dr. Jayanti Sinha², Dr. Sweety Sinha³

Senior Resident, Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India.

²Department of Obstetrics & Gynaecology, AIIMS, Patna, Bihar, India.

³Assistant. Professor, Department of Obstetrics & Gynaecology, PMCH, Patna, Bihar, India

Corresponding Author: Dr. Anupam Kumari

Abstract

Aim: To study the Hyperhomocysteinemia levels in pregnancy induced hypertension.

Materials and Methods: A Case control study was conducted in the Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India, for 1 year. Total 240 pregnant female subjects in the age group of 16-55 yrs was selected. Out of them 120 were normotensive pregnant women (NPW) in their third trimester and were chosen as control Group 1 and 120 pregnancy induced hypertensive (PIH) patients in their third trimester were chosen as study Group 2.

Results: The hypertensive subjects had similar age distributions and values of BMI when compared to normotensive controls. The percentage of subjects who were smokers was not different between the two groups. Although most of the hypertensive patients were receiving antihypertensive treatment, their systolic and diastolic blood pressure were significantly higher than those of the normotensive individuals. The mean and standard deviation of Homocysteine, uric acid and magnesium levels of NPW and PIH groups. The Hcy level was significantly increased with a mean and standard deviation (SD) value of $20.77 \pm 5.95 \mu\text{mol/l}$ (p-value=0.001) and the uric acid level was also increased with a mean and SD of $6.15 \pm 0.77 \text{mg/dl}$ (p-value= 0.002) in the PIH. The Hcy level was $9.12 \pm 2.33 \mu\text{mol/l}$ and the Uric acid level was $3.17 \pm 0.82 \text{mg/dl}$ in NPW group. . The magnesium level was lower in the PIH compared to NPW (p - value=0.001).

Conclusion: The Hcy and uric acid levels were increased and Magnesium level was decreased in PIH women than the NPW. So these parameters should be part of the evaluation of the pregnant women presenting with hypertension. Thereby, we can reduce the maternal and fetal mortality rate.

Keywords: Homocysteine, Hyperhomocysteinemia, Uric acid, Magnesium and Pregnancy Induced Hypertension (PIH).

Introduction

Preeclampsia is one of the most common pregnancy complications causing high morbidity and mortality for both mother and fetus especially in developing countries. Although the exact cause of preeclampsia is still unknown, the basic pathology lies in the endothelial dysfunction and intense vasospasm.¹ Homocysteine is an amino acid formed during demethylation of dietary methionine. Elevated blood levels of homocysteine are now recognized as an important risk marker for both preeclampsia and eclampsia.² The vascular effects of hyper homocysteinemia includes endothelial cell injury and thrombus formation. Elevated plasma homocysteine is an independent risk factor for peripheral vascular disease

and coronary artery disease.³ Homocysteine is an essential amino acid required for growth of cells and tissues in the human body.⁴ Homocysteine is eliminated from the body via conversion to cystathionine by a reaction catalysed by vitamin B6 and to methionine catalysed by vitamin B12 and folic acid. Homocysteine is found in low concentrations in all tissues under normal conditions whereas it accumulates in tissues and plasma if these catalytic vitamins were depleted. Both retrospective and prospective studies have demonstrated that hyperhomocysteinemia is a risk factor for cardiovascular disease that is independent of classic risk factors, such as smoking, hypercholesterolemia, and hypertension.⁵⁻⁷ Hypertensive disorders are common in pregnancy and they cause serious complications like eclampsia, hemorrhage and infection leading to increased maternal and fetal mortality. Pregnancy induced hypertension is defined as a condition that results in persistent elevation of blood pressure of $\geq 140/90$ mm Hg or more (confirmed by two measurements in sitting posture at least six hours apart) or 30 mm Hg systolic or ≥ 15 mm Hg diastolic over base values arising *denovo* in pregnancy. Homocysteine, an essential amino acid is found in many animals and plant foods, it's formed from methionine, a sulphur containing amino acid. Elevated levels of homocysteine play an independent role for atherosclerosis and vascular thrombosis.⁸ A derangement in the homocysteine-methionine metabolism leads to vascular damage causing hypertension and further to the classical clinical manifestations of preeclampsia. Elevated homocysteine is a risk factor for endothelial dysfunction and vascular disease such as atherosclerosis and occlusive vascular disorders.⁹ The mean homocysteine levels normally decrease with gestation either due to physiological response to the pregnancy, increase in estrogen, hemodilution from increased plasma volume or increased demand for methionine by both the mother and fetus.¹⁰ Homocysteine is a naturally occurring amino acid derivative in the body. Increase in hormones such as estrogen and cortisol during pregnancy may also mediate, specific decrease in Hcy concentration. The mechanism behind the endothelial dysfunction had been demonstrated in experiments. Hcy decreases the expression of a wide range of antioxidant enzymes. This impairs endothelial nitric oxide (NO) bioavailability by decreasing Glutathione peroxidase activity which raises the possibility that Hcy sensitizes cells for reactive oxygen species (ROS).¹¹ During early pregnancy serum uric acid levels fall, often to 3 mg/dl or below, related to the uricosuric effects from estrogen and from the increase in renal blood flow. Uric acid levels then increase during the third trimester. However, it is known that subjects destined to develop preeclampsia show slightly higher serum uric acid levels during the first trimester in association with a relative reduction in urinary urate excretion.¹² Increasing evidence suggests that an elevated serum uric acid in pregnancy may not only be a valuable biomarker for preeclampsia but may also have a contributory role in the pathogenesis of the maternal and fetal manifestations. Uric acid is a potent inhibitor of endothelial function, induces systemic and glomerular hypertension in animals, and passes freely into the fetal circulation.¹³ Uric acid has been found to block vascular endothelial growth factor (VEGF)-induced endothelial proliferation and thus may have a direct role in blocking fetal angiogenesis resulting in small for gestational age infants.¹⁴ Uric acid can also block trophoblast invasion *in vitro*.¹⁵ These studies suggest that measurement of serum uric acid is clinically useful and Serum calcium and magnesium are very important for metabolism at the cellular level and are vital for muscle contraction and cell death and neuronal activity making it very essential in pregnancy.¹⁶ Magnesium plays an important role in peripheral vasodilatation. Homocysteine level causes injury to the vascular system of both maternal and fetal organs and increased uric acid level and decreased magnesium levels affects vascular and renal systems, thereby aggravating the process leading to eclampsia, resulting in increased maternal-fetal mortality and morbidity. So this study is taken up to assay the levels of Homocysteine, uric acid and magnesium in PIH and NPW group.

Material and Methods

A Case control study was conducted in the Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India, for 1 year.

Inclusion Criteria:

Control: Normotensive pregnant women in their third trimester with no complications.

Cases: Pregnant women with Pregnancy Induced Hypertension in their third trimester with persistent elevation of blood pressure 140/90 mmHg and more confirmed by two measurement (In the sitting posture, at least six hours apart) or increase of at least 30mm of Hg systolic or 15mmHg diastolic over baseline value and both groups in age matched in the range 16 to 55yrs were included for the study.

Exclusion Criteria:

Women with previous history of Hypertension, Diabetes Mellitus, Renal or heart disease and other complications of pregnancy were excluded.

Methodology

Total 240 pregnant female subjects in the age group of 16-55 yrs was selected. Out of them 120 were normotensive pregnant women in their third trimester and were control. 120 pregnancy induced hypertensive (PIH) women in their third trimester were chosen as cases. Five ml blood sample was collected by venepuncture of the cubital vein after an overnight fast. Homocysteine was estimated by Axis Homocysteine enzyme Immunoassay [ELISA] method and uric acid was estimated by uricase method and magnesium estimated by colorimetric method.

Results

The hypertensive subjects had similar age distributions and values of BMI when compared to normotensive controls. The percentage of subjects who were smokers was not different between the two groups (Table 1). Although most of the hypertensive patients were receiving antihypertensive treatment, their systolic and diastolic blood pressure were significantly higher than those of the normotensive individuals.(Table 2).

Table 3 shows the mean and standard deviation of Homocysteine , uric acid and magnesium levels of NPW and PIH groups. The Hcy level was significantly increased with a mean and standard deviation (SD) value of $20.77 \pm 5.95 \mu\text{mol/l}$ (p- value=0.001) and the uric acid level was also increased with a mean and SD of $6.15 \pm 0.77 \text{mg/dl}$ (p-value= 0.002) in the PIH. The Hcy level was $9.12 \pm 2.33 \mu\text{mol/l}$ and the Uric acid level was $3.17 \pm 0.82 \text{mg/dl}$ in NPW group. . The magnesium level was lower in the PIH compared to NPW (p - value=0.001)

Table 1: demographic profile of patients

| Variables | PIH(mean ± sem) | NPW(mean± sem) | P-value |
|--------------------------|-----------------|----------------|---------|
| Age | 49.5± 1 | 48.7±1 | 0.71 |
| BMI (kg/m ²) | 25.9 ± 0.39 | 24.5±0.21 | 0.61 |
| Smoking status | 27.5% | 32.5% | 0.288 |

Table 2: Blood pressure (mm Hg)

| | PIH | NPW | P-value |
|------------------------|---------|--------|---------|
| Blood pressure (mm Hg) | | | |
| Systolic | 140 ±12 | 117±23 | .001 |
| Diastolic | 83±22 | 73±23 | .001 |

Table 3: Homocysteine, Uric acid and Magnesium levels in PIH & NPW

| Parameter | Patients | Mean± Std. Deviation | P value |
|-----------------------|----------|----------------------|---------|
| Homocysteine (µmol/l) | PIH | 20.77±5.95 | .001 |
| | NPW | 9.12 ± 2.33 | |
| Uric acid level mg/dl | PIH | 6.15±0.77 | .002 |
| | NPW | 3.17 ± 0.82 | |
| Magnesium | PIH | 1.45 | .001 |
| | NPW | 1.49 | |

Discussion

In the present study, there was an increased Hcy level with a p value=0.001 and an increased uric acid level with a p value=0.002 in the PIH women when compared with NPW group.

In this study, we found homocysteine levels were higher in hypertensive patients as compared to a group of age, gender, and BMI-matched normotensive individuals. It has been shown that plasma homocysteine concentrations are higher in elderly isolated systolic hypertensive subjects than in normotensive controls.¹⁷ Increased Hcy level had been proposed to explain endothelial cell dysfunction including direct cell injury in maternal circulation which causes a chronic inflammatory and endothelial damage and impairs synthesis of nitric oxide which causes uteroplacental insufficiency causing maternal vascular damage and increased reactive oxygen species (ROS), leading to hypertension.¹⁸ Similar findings are reported by Sanchez et al.¹⁹, Harma et al.²⁰ and Maruotti et al.²¹ The interplay of various biological mechanism and effects of HHcy activates multiple processes leading to disorders. It enhances the production of several pro-inflammatory cytokines factors like Interlukin-8(IL-8).²² It enhances the intracellular production of superoxide anions. Lopez et al. found an association between hyperhomocysteinemia and preeclampsia. In their study, the concentration of plasma homocysteine levels in patients with preeclampsia was higher than the NPW.²³ Power et al. found high levels of homocysteine in preeclamptic patients as compared to the NPW. The present study showed that serum magnesium level was significantly reduced in the PIH women than the NPW. These findings confirmed that hypomagnesemia may be one of the etiologies of preeclampsia. These results were consistent with earlier study by Zhao F and Frankel.Y et al, also showed mean serum magnesium was slightly lower in PIH women as compared to NPW group.²⁴ Lowered plasma or serum magnesium concentrations in preeclampsia may contribute to the development of hypertension in pregnancy.

Decreased renal excretion due to hypertension causes increased uric acid level in women with PIH initiating maternal and fetal complications. Similarly, in the present study, the Uric acid levels were increased in PIH women. In the study by Bellomo et al the results were remarkable, Uric acid conferred an 8-9 fold risk for preeclampsia and a 1.6 - 1.7 fold risk for small for gestational age infants.. These studies suggest that measurement of serum uric acid is clinically useful and should be part of the evaluation of the pregnant patient presenting with hypertension. Increasing evidence suggests that an elevated serum uric acid in pregnancy may not only be a valuable biomarker for preeclampsia but may also have a contributory role in the pathogenesis of the maternal and fetal manifestation.²⁵

Conclusion

The Hcy and uric acid levels were increased and Magnesium level was decreased in PIH women than the NPW. So, early screening and diagnosis of PIH can be done by adding these parameters with other routine antenatal work-up in pregnant women.

Reference

1. Roberts J, Taylor R, Musci T, Rodgers G, Hubel C, McLaughlin M. Preeclampsia: An

- endothelial cell disorder. *Am J Obstet Gynecol.* 1989;161:1200-4.
2. Rajkovic A, Catalano PM, Manilow MR. Elevated homocysteine levels with preeclampsia. *Obstet Gynecol.* 1997;90:168-71.
 3. Refsum H, Ueland PM, Nygard O, Vollset SE. Homocysteine and cardiovascular disease. *Annu. Rev. Med.* 1998;49:3-62.
 4. Refsum H, Ueland PM, Nygard O, Brattstrom L, Vollset SE. Total homocysteine and cardiovascular disease. *J Int Med.* 1999;246:425-54
 5. Boushey CJ, Beresford SA, Omenn GS, Motulsky AG: A quantitative assessment of plasma homocysteine as a risk factor for vascular disease: probable benefits of increasing folic acid intakes. *JAMA* 1995;274:1049–1057
 6. Graham IM, Daly LE, Refsum HM, Robinson K, Brattstrom LE, Ueland PM, Palma-Reis RJ, Boers GH, Sheahan RG, Israelsson B, Uiterwaal CS, Meleady R, McMaster D, Verhoef P, Witteman J, Rubba P, Bellet H, Wautrecht JC, de-Valk HW, Sales-Luis AC, Parrot-Rouland FM, Tan KS, Higgins I, Garcon D, Andria G, Medrano MJ, Candito M, Evans A, Andria G: Plasma homocysteine as a risk factor for vascular disease. *JAMA* 1997;277:1775–1781.
 7. Nygard O, Nordrehaug JE, Refsum H, Ueland PM, Farstad M, Vollset SE: Plasma homocysteine levels and mortality in patients with coronary artery disease. *N Engl J Med* 1997;337:230–236.
 8. Antoniadis C, Antonopoulos AS, Tousoulis D, Marinou K, Stefanadis C. Homocysteine and coronary atherosclerosis for folate fortification to the recent clinical trials. *EurHeart J.* 2009;30: 6-15.
 9. Allain.C.C, Poon LS, Chan CS, Richmond W, Fu PC. Enzymatic determination of total serum cholesterol. *ClinChem.* 1974;20: 470- 475.
 10. Also-Rallo E, Lopez-Quesada E, Urreiziti R, Vilaseca MA, Laila JM, Balcells S, et al. Polymorphisms of genes involved in homocysteine metabolism in preeclampsia and in uncomplicated pregnancies. *Eur J Obstet Gynecol Reprod Biol.* 2005; 120: 45-52.
 11. Weiss N, Heydrick S, Xhangyy, Bierl C. Cellular redox state and endothelial dysfunction in mildly hyper homocysteinemic cystathionine beta- synthase-deficient-mice. *Arterioscler thrombvascoil.* 2002; 22:34-41.
 12. Laughon SK, Catov J, Powers RW, Roberts JM, Gandley RE. First trimester uric acid and adverse pregnancy outcomes. *Am J Hypertension.* 2011; 24:489–495.
 13. Kang DH, Finch J, Nakagawa T, Karumanchi SA, Kanellis J, Granger J, et al. Uric acid, endothelial dysfunction and pre-eclampsia: searching for a pathogenetic link. *Journal of hypertension.* 2004;22:229–235
 14. Feig DI, Nakagawa T, Karumanchi SA, Oliver WJ, Kang DH, Finch J et al. Hypothesis: Uric acid, nephron number, and the pathogenesis of essential hypertension. *Kidney international.* 2004;66:281–287.
 15. Shannon A, Bainbridge, James M. Roberts, Frauke von Versen-Höynck, Jessa Koch, Lia Edmunds, and Carl A. Hubel. Uric acid attenuates trophoblast invasion and integration into endothelial cell monolayers. *Am J Physiol Cell Physiol.* 2009;297:C440–C450.
 16. Sandip S, Asha K, Paulin G, Hiren S, Gagandeip S, Amit V. A comparative study of serum uric acid, calcium and magnesium in preeclampsia and normal pregnancy. *J Advanc Res Biol Sci.* 2013 ;5(1):55-58.
 17. Sutton-Tyrell K, Bostom A, Selhub J, Zeigler-Johnson C: High homocysteine levels are independently related to isolated systolic hypertension in older adults. *Circulation* 1997;96:1745–1749.
 18. Anderson A, Hullberg.B, Brattstrom L. Isaksson A. Decreased serum homocysteine in pregnancy. *Eur J Clin Chem. ClinBiochem.* 1992;30:377-9.
 19. Sanchez SE, Zhang C, Rene Malinow M, Ware-Jauregui S, Larrabure G, Williams MA.

- Plasma folate, vitamin B(12) , and homocysteine concentrations in preeclamptic and normotensive Peruvian women. *Am J Epidemiol.* 2001;53: 474-480.
20. Harma M, Harma M, Kocyigit A. Correlation between maternal plasma homocysteine and zinc levels in preeclamptic women. *Biol Trace Elem Res.* 2005;104: 97-105.
 21. Maruotti G, Del Bianco A, Amato AN, Lombardi L, FulgeriAM, Pietropolo F. Preeclampsia and high serum levels of homocysteine. *Minerva Ginecol.* 2005;57: 165-170.
 22. Wang G,Siow YL,Homocysteine induced monocyte chemoattractant Protein 1 expression by activating NF-Kappa B in THP-1 macrophages. *Am J Physiol Heart Circ Physiol.*2001;280:H2: 840-7.
 23. Lopez- Qyesada E, Vilaseca MA, Laila JM. Plasma total homocysteine in uncomplicated pregnancy and in preeclampsia. *Eur J Obstet Gynecol.* 2007;275: 121-124.
 24. Zhao F. Ca, Mg, Cu and Zn contents of the maternal and umbilical cord serum in pregnancy induced hypertension. *Chinese J of obstet and Gynecol.*1989;24(4):212-14.
 25. Bainbridge SA, Roberts JM. Uric Acid as a Pathogenic Factor in Preeclampsia. *Placenta.* 2008;29S:67–72.

Received: 01-08-2020 || Revised: 16-08-2020 || Accepted: 28-08-2020