

Original research article

Observational cross-sectional study to determine the menstrual pattern and various associated menstrual problems among 10-19 years old females

Dr. Anupam Kumari¹, Dr. Jayanti Sinha², Dr. Sweety Sinha³

¹Senior Resident, Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India.

²Department of Obstetrics & Gynaecology, AIIMS, Patna, Bihar, India.

³Assistant. Professor, Department of Obstetrics & Gynaecology, PMCH, Patna, Bihar, India

Corresponding Author: Dr. Anupam Kumari

Abstract

Background: Adolescence is a transitional stage extending from 10-19 years characterized by rapid physical, mental and sexual changes. It is also characterized by hormonal changes. As girls attain puberty at this age, they have various problems associated with menstruation.

Aim: The aim of the present study was to determine the menstrual pattern and various menstrual problems among urban adolescent girls.

Material and methods: This was a cross sectional study conducted in the Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India, for 18 months. 110 adolescent girls randomly who presented with menstrual problems in gynaecology OPD were included in this study. Total of 110 girls were asked to fill up a questionnaire which included questions like age of menarche, pattern of menstruation, duration and amount of flow, number of pads used.

Result: Majority of girls (63.63%) attained their menarche between 10-13 years of age and remaining 36.37% were in the age group of 14-16 years. Majority of girls had problem of oligomenorrhea (60.91%). 25.45% of girls had normal cycle length of 28-35 days and only 8.18% had frequent bleeding with cycle length less than 28 days. But duration (3-5 days) and quantity of flow was normal in maximum girls 80% and 62.73% respectively. Excessive flow with passage of clots or duration of flow for more than 5 days was noted in 20% and 13.64% of girls respectively.

Conclusion: Menstrual problems are present in majority of girls in the study group which has some effect on the academic performance and other extracurricular activities in school.

Keywords: Adolescent girls, Menstrual disorders, Dysmenorrhoea, Premenstrual symptoms

Introduction

Adolescence is a transitional period from childhood to adulthood characterized by rapid physical, mental, and sexual development. Menstruation is a natural phenomenon which is an important indicator of women's health, reflecting their endocrine function. However data on experiences of menstruation and its impact on health status, quality of life, and social integration among women in developing countries is scanty. Dysmenorrhoea, for example is a common problem, yet it remain poorly understood and is not taken into consideration when assessing adolescent health problems. Dysmenorrhoea is defined as difficult menstrual flow or painful menstruation. About 50-75% of young girls complain of this. This is one of the leading cause of loss school days.¹ Premenstrual syndrome is characterized by mood changes, anxiety and somatic symptoms like backache, breast tenderness. They appear about one week

before the onset of menstrual periods and disappear after the appearance of menstrual periods.²

Adolescence is a period between childhood and adulthood, a transition phase characterised by discernible reproductive maturity. As per definition of World Health Organization, the period of adolescence ranges from 10- 19 years of age.¹ India has largest population of adolescents and of the total female population in India 10.7% and 9.7% are girls in the vulnerable age groups of 10-14 years and 15-19 years respectively.² Adolescent girls face substantial morbidities related to menstruation which are generally unrecognized and uncared due to unfamiliarity of girls or, difficulty in understanding normal characteristics of their menstrual cycles and further their reluctance to discuss this issue.³ Attainment of menarche at appropriate age signifies and assures normal development and functioning of the female reproductive system. However, it's timing and subsequent progression are individual-specific and vary within a broad range of normality depending on maturation of the complex interactions of hypothalamus, pituitary gland and ovary. Menarche typically occurs within 2-3 years after thelarche at Tanner stage IV breast development usually between 12 and 13 years.^{3,4} Among Indian females the mean age at menarche is 13.76 years with a secular decline of nearly one month per decade.⁵ Disturbances of menstrual bleeding manifest in a wide range of presentations and abnormal uterine bleeding (AUB) is the overarching term used to describe any departure from normal menstruation or from a normal menstrual cycle pattern. The key characteristics are regularity, frequency, heaviness of flow, and duration of flow, but each of these may exhibit considerable variability.⁶ As evident from previous literature, usually after third year of menarche the interval between bleeding periods is in the range of 21-34 days, with a flow lasting from 3 to 7 days and a mean menstrual blood loss of 35 ml (range 5-80 ml).³⁻⁸ Significant variations may be somewhat physiologic first few years after menarche or, may be attributable to significant underlying medical issues like polycystic ovary syndrome, thyroid problems, mental stress, hypothalamic dysfunction, primary pituitary disease, primary ovarian insufficiency, coagulopathies, uterine lesions and rarely malignancy.³ And therefore it is important to have an understanding of the menstrual pattern and its characteristics among adolescent girls. Hence the present investigation was conducted to study the pattern and features of menstruation among adolescent girls.

Material and methods

This was a cross sectional study conducted in the Department of Obstetrics & Gynaecology, NSMCH, Bihta, Patna, Bihar, India, for 18 months.

Methodology

110 adolescent girls randomly who presented with menstrual problems in gynaecology OPD were included in this study. Total of 110 girls were asked to fill up a questionnaire which included questions like age of menarche, pattern of menstruation, duration and amount of flow, number of pads used. Data collected compiled and analyzed.

Result

This study shows Majority of girls (63.63%) attained their menarche between 10-13 years of age and remaining 36.37% were in the age group of 14-16 years (Table 1). Majority of girls had problem of oligomenorrea (60.91%). 25.45% of girls had normal cycle length of 28-35 days and only 8.18% had frequent bleeding with cycle length less than 28 days. But duration (3-5 days) and quantity of flow was normal in maximum girls 80% and 62.73% respectively. Excessive flow with passage of clots or duration of flow for more than 5 days was noted in 20% and 13.64% of girls respectively. Only 6.36% had flow for less than 2 days (Table 2).

Menstruation is associated with multiple morbidities, in our study majority of girls had problem of oligomenorrhea (60.91%) followed by dysmenorrhea (51.82%) and menorrhagea (22.73%). Dysmenorrhea was the main reason of absenteeism from school (with 43.33%). Menorrhagia along with dysmenorrhea was second most common reason (13.33%) for absenteeism from school. Third main reason was menorrhagia, reported by 23.33%.

Table 1: Distribution of girls according to age at menarche

Age at menarche	Number of patients	Percentage
10-13	70	63.63
14-16	40	36.37
17-19	0	0

Table 2: Pattern of menstrual cycle

Duration of blood flow (days)	Number of patients	Percentage
<2	7	6.36
3-5	88	80
>5	15	13.64
Length of cycle (days)		
<28	9	8.18
28-35	35	25.45
>35	66	60
Quantity of blood flow		
Normal	69	62.73
Excessive	22	20
Scanty	19	17.27

Table 3: Common menstrual problems

Menstrual problems	Number of patients	Percentage
Oligomenorrhea	67	60.91
Dysmenorrhea	57	51.82
Menorrhagia	25	22.73
Premenstrual symptoms	17	15.45
Leucorrhea	2	1.82

Table 4: Reasons for absenteeism from school related to menstrual problem (N-30)

Menstrual Problem	Numbers	Percentage (%)
Dysmenorrhea	13	43.33
Menorrhagea + dysmenorrea	4	13.33
Menorrhagia	7	23.33
Premenstrual syndrome	4	13.33
Fear of leakage	2	6.67

Discussion

In our study the majority of girls (63.63%) attained their menarche between 10-13 years of age and remaining 36.37% were in the age group of 14-16 years which collaborates with study conducted by S Rokade⁹, who reported mean age of menarche as 12.6±1.05years in Maharashtrian girls, Varuna Pathak¹⁰ reported age of menarche as 13 years. It is also collaborates with various studies conducted in developed countries like studies report the mean age of menarche as 13.28years in Turkey¹¹, 13 years in Russia¹², 12.3 years in

Thailand¹³, 12.5 years in Japan¹⁴, 13.2 years in Egypt.¹⁵ Our study shows 60% of girls had prolong cycles of more than 35 days with irregular cycles. However studies conducted in Guntur, Andhra Pradesh¹⁶ and Meerut, Utter Pradesh¹⁷ observed 66.54% and 66.9% girls with regular menstruation cycles, and in study conducted by M. Kulkarni 11.16% girls with irregular cycles. This difference may be due to nutritional factors and general health of study subjects. Menstruation is associated with multiple morbidities, in our study majority of girls had problem of oligomenorrhea (60.91%) followed by dysmenorrhea (51.82%) and menorrhagea (22.73%) of the population where as various studies showed dysmenorrhea as a commonest menstrual problem^{10,18,19}, and oligomenorrhoea was observed in 16.08% girls by Mohite RV¹⁹ and 3.2% by Varuna Pathak.¹⁰ In our study dysmenorrhea was the main reason of absenteeism from school (with 43.33%). Menorrhagia along with dysmenorrhea was second most common reason (13.33%) for absenteeism from school. Third main reason was menorrhagia, reported by 23.33%, whereas various studies showed premenstrual symptoms ranged from 41.5% to 75.4%. Menorrhagia was seen in 17.82% population by Mohite RV, 16.07% by M Kulkarni which is comparable with our study.^{18,19} Our study shows oligomenorrhea is the most common menstrual problem and second one is dysmenorrhea. As peri pubertal phase is phase of hormonal imbalance, further investigations and evaluations are needed for the reason of oligomenorrhea and menorrhagea. Our study also reveals that the dysmenorrhea is the most common reason of absentees from school/college, similar findings were found in the studies conducted in North India South India and Mumbai.²⁰⁻²² Level of education, awareness, cultural taboos and practices affects the percentage of girls presenting with their menstrual complaints to healthcare facility. Other factors like nutrition, anemia, personal hygiene, geographic conditions, socio-economic factors also influence menstrual problems.

Through our study we gained knowledge on various menstrual problems and their contribution in physical as well as psychological disturbance among adolescent girls.

Conclusion

Menstrual problems are present in majority of girls in the study group which has some effect on the academic performance and other extracurricular activities in school.

References

1. Salkhan S. Adolescent gynaecological issue. Text book of Gynaecology. 1st ed. Chapter 13. Jaypee brothers Medical publishers (p) Ltd; 2011: 70-72.
2. Mashankar VA. Menstrual Problems in adolescents. In: Bhave SY, editor. Bhave's Textbook of Adolescent Medicine. New Delhi: Jaypee Brothers Medical Publishers (P) Ltd; 2006: 95-104.
3. Menstruation in girls and adolescents: using the menstrual cycle as a vital sign. Committee Opinion No. 651. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2015; 126:e143-6.
4. Padubidri VG, Daftary SN. Chapter 4: Puberty, paediatric and adolescent gynaecology. Shaw's textbook of Gynaecology. 16th ed. New Delhi: RELX India Private Limited; 2016:51-63.
5. Pathak PK, Tripathi N, Subramanian SV. Secular trends in menarcheal age in India-evidence from the Indian human development survey. *PLoS ONE*. 2014;9(11):1-13.
6. Fraser IS, Critchley HD, Broder M, Munro MG. The FIGO Recommendations on Terminologies and Definitions for Normal and Abnormal Uterine Bleeding. *Seminars in Reproductive Medicine*. 2011; 29(5):383-90.

7. Rigon F, De Sanctis V, Bernasconi S, Bianchin L, Bona G, Bozzola M, et al. Menstrual pattern and menstrual disorders among adolescents: an update of the Italian data. *Italian J Pediatr.* 2012;38(1):38.
8. Van Hooff MH, Voorhorst FJ, Kaptein MB, Hirasig RA, Koppelaar C, Schoemaker J. Relationship of the menstrual cycle pattern in 14-17 year old adolescents with gynaecological age, body mass index and historical parameters. *Human Reproduction (Oxford, England).* 1998;13(8):2252-60.
9. Rokade S, Mane A. A Study Of Age At Menarche, The Secular Trend And Factors Associated With It. *The Internet Journal of Biological Anthropology*, 2008, 3(2).
10. Varuna Pathak, Madhuri Chandra, Mamta Paraste. Study of Menstrual Patterns among School Going Adolescent Girls of Urban India. *Journal of Evolution of Medical and Dental Sciences.* 2015; 4(70):12174-12178, DOI: 10.14260/jemds/2015/175,
11. Vicdan K, Kükner S, Dabakolu T, Ergin T, Kele° G, Gokmen O. Demographic and epidemiologic features of female adolescents in Turkey. *J Adolesc Health.* 1996; 18:54-8.
12. Iampol skaia I. Dynamics of puberty levels in girls of Moscow. *Gig Sanit.* 1997; 3:29-30.
13. Piya-Anant M, Suvanichchati S, Bharscharil M, Jirochkul V, Worapitaksanond S. Sexual maturation in Thai girls. *J Med Assoc Thai.* 1997; 80:557-64.
14. Nakamura I, Shimura M, Nonaka K, Miura T. Changes of recollected menarcheal age and month among women in Tokyo over a period of 90 years. *Ann Hum Biol.* 1986; 13:547-54.
15. Attallah N, Matta W, El-Mankoushi M. Age at menarche of schoolgirls in Khartoum. *Ann Hum Biol.* 1983; 10:185-8.
16. Keerti Jogdand, Pravin Yerpude. A community based study on menstrual hygiene among adolescent girls. *Indian Journal of Maternal and Child Health.* 2011; 13(3):1-6.
17. Dixit RR, Sabane HH, Durge PM. Reproductive Health of Adolescent Girls in an urban population of Meerut, Uttar Pradesh. *Health and Population: Perspectives and Issues.* 2009; 32(4):204-209.
18. 8. Kulkarni M, Durge PM. Reproductive Health morbidities among adolescent girls: Breaking the silence. *Ethno Med.* 2011; 5(3):165-168.
19. Mohite RV, Mohite VR, Kumbhar SM, Ganganahali P. Common Menstrual Problems among Slum Adolescent Girls of Western Maharashtra, India JKIMSU, 2013, 2(1).
20. Singh AJ. The place of menstruation in the reproductive lives of women of rural north India. *Indian J Comm Med.* 2006; 31(1):10-14.
21. Ali TS, Rizvi SN. Menstrual knowledge and practices of female adolescents in urban Karachi, Pakistan. *J Adolesc.* 2010; 33(4):531.
22. Narayana KA, Srinivasa DK, Petlo PJ. Puberty rituals, reproductive knowledge and health of adolescent school girls in south India. *Asia Pacific Population Journal.* 2001; 16(2):225-38

Received: 01-07-2020 || Revised: 09-07-2020 || Accepted: 24-07-2020