

Original research article

Platelet Parameters in Neonatal Bacterial Sepsis with Different Isolated Organisms: prospective observational study**Dr. Amresh Kumar Sahu¹, Dr. Shyam Bahadur Prasad², Dr. Arvind Kumar³****¹Senior Resident, Department of Paediatrics, Government Medical College and Hospital, Bettiah, West Champaran, Bihar, India.****²Senior Resident, Department of Paediatrics, Government Medical College and Hospital, Bettiah, West Champaran, Bihar, India.****³Professor and HOD, Department of Paediatrics, Government Medical College, Bettiah, West Champaran, Bihar, India.****Corresponding Author: Dr. Shyam Bahadur Prasad****Abstract**

Aim: The aim of the present study is to determine the Platelet Parameters with Different Isolated Organism in Neonatal Bacterial Sepsis.

Methods: This cross sectional prospective observational study was done in the Department of Paediatrics, Government Medical College, Bettiah, West Champaran, Bihar, India from December 2018 to July 2019. Neonate with presence of more than 2 clinical features suggestive of sepsis with positive C - reactive protein (CRP), were enrolled for the study. Baseline neonatal characteristics like gestational age, birth weight, gender, mode of delivery, APGAR score was noted for all enrolled babies. Neonates were examined clinically for primary illness according to standard format. CBC was done by automated cell counter. Blood culture was done by BACTEC method. Platelet parameters studied were platelet count at baseline & at onset of sepsis; degree of thrombocytopenia; duration of thrombocytopenia; mean platelet volume (MPV) and platelet nadir. The platelet count at the onset of sepsis was considered as platelet count coinciding with blood sample showing positive culture report.

Results: Total 100 babies with culture positive sepsis were analyzed. Mean gestational age of study group was 34.98 ± 3.55 weeks; mean birth weight was 1.899 ± 0.72 kg; male: female ratio was 1.22: 1. Mode of delivery was normal vaginal delivery (NVD) in 69% neonates. Mean APGAR score of study group was 7.81 ± 1.33 . Our study observed early onset sepsis in 72% babies and Gram negative sepsis in 75% babies. Overall thrombocytopenia was observed in 90% babies with 57% having moderate to severe degree of thrombocytopenia. The proportion of severe degree of thrombocytopenia (17.33% Vs 4%), higher MPV (59.33% Vs 56%) and longer duration of thrombocytopenia (3.78 ± 0.55 Vs 3.02 ± 0.61) was observed more with Gram negative sepsis than with Gram positive sepsis, but statistical significance was not found. We observed statistically significant platelet nadir with Gram negative sepsis. Out of 100 neonates with culture positive sepsis, Gm negative sepsis was observed in (75) 75% and Gm positive sepsis was observed in (25) 25%. The proportion of isolated organisms was: 30% klebsiella, 25% pseudomonas, 19% Coagulase- negative Staphylococci (CONS), 14% Citrobacter, 6% S. Aureus & 6% Acinetobacter.

Conclusion: Thrombocytopenia is a frequent occurrence in neonates with sepsis especially with Gram negative organism.

Keywords: thrombocytopenia, sepsis, neonates

Introduction

Sepsis is the commonest cause of neonatal mortality; it is responsible for about 30-50% of the total neonatal deaths in developing countries.¹ Neonatal mortality in India is 27 per 1000 live births.² Neonatal sepsis is defined as a clinical syndrome characterized by signs of systemic infection and documented by a positive blood culture in the first four weeks of life.³ Incidence of neonatal sepsis in Asia is 7.1 to 38 per 1000 live births,⁴ in.⁵ The pattern of infection by various organisms varies from one institution to another and even from year to year in the same institution.⁶ The common organisms responsible for neonatal sepsis are Klebsiella pneumoniae, Acinetobacter, Escherichia coli (E.Coli), Pseudomonas aeruginosa, Salmonella, Haemophilus influenzae, Proteus, Coagulase negative staphylococci, Staphylococcus aureus, Streptococcus, Pneumococcus, Flavobacterium freundii, Candida etc.⁷ Blood culture is the gold standard test to diagnosis neonatal sepsis. But it could be positive only in 30-40% cases.⁸ Obtaining blood cultures from neonates can be difficult, sample volumes are small, and a substantial number are negative or contaminated.⁹ Estimation of cytokines and C Reactive Protein (CRP) levels are potentially useful in this respect.¹⁰ Complete blood count including platelet count is a good predictor of sepsis in newborns.¹¹ Thrombocytopenia is a major indicator of sepsis in neonates.¹¹ It has been shown that 75% of culture positive neonates had thrombocytopenia.¹² Majority of newborn developed thrombocytopenia by 36-48 hours after developing neonatal sepsis and average duration of thrombocytopenia persisted around 6 days.¹³ There is a positive association of Gram negative infection and thrombocytopenia.¹⁴ Fungal infection is also associated with a greater degree of thrombocytopenia.¹⁵ Congenital viral infections like cytomegalovirus (CMV), toxoplasma, rubella and herpes simplex are important causes of thrombocytopenia in neonatal period and early infancy.

Material and Methods

This cross sectional prospective observational study was done the Department of Paediatrics, Government Medical College, Bettiah, West Champaran, Bihar, India from December 2018 to July 2019 after taking the approval of the protocol review committee and institutional ethics committee.

Inclusion criteria

- Neonatal sepsis was considered if: presence of more than 2 clinical features suggestive of sepsis with positive C - reactive protein (CRP) and positive blood culture.
- The clinical features suggestive of sepsis considered were: hypothermia/fever; poor cry/poor feeding/refusal to suck; lethargy; fast breathing/respiratory distress/apnoea; abdominal distension/Diarrhoea/vomiting; sclerema; prolonged capillary refill time; bulging anterior fontanel/convulsion.

Exclusion criteria

- Neonates with major congenital malformation, chromosomal anomalies.
- Neonates with maternal thrombocytopenia.
- Maternal history of pregnancy induced hypertension (PIH)

Procedure

Baseline neonatal characteristics like gestational age, birth weight, gender, mode of delivery, APGAR score was noted for all enrolled babies. Neonates were examined clinically for primary illness according to standard format. As per our neonatal unit policy baseline investigation of complete blood count (CBC), CRP and Blood culture were sent at the time of admission and subsequently on clinical suspicion of sepsis. Platelet count was repeated twice a week or at time of sampling for other investigations in thrombocytopenic

neonates until platelet count was normalized. Blood for investigation was collected by vene-puncture under strict asepsis. CBC was done by automated cell counter. Blood culture was done by BACTEC method.

Platelet parameters studied were platelet count at baseline & at onset of sepsis; degree of thrombocytopenia; duration of thrombocytopenia; mean platelet volume (MPV) and platelet nadir. The platelet count at the onset of sepsis was considered as platelet count coinciding with blood sample showing positive culture report. Thrombocytopenia was defined as platelet count less than 150,000 / cmm and graded as mild if platelet count between 50,000 to 150,000 / cmm, moderate if counts were between 20,000 to 50,000 / cmm and severe if count was less than 20,000/ cmm. The normal range for MPV was 7.4 to 11.4 fentolitre (fl) (as per lab reference value). Duration of thrombocytopenia was the number of continuous days that the platelet count remained below 150,000/cmm. Platelet nadir means lowest platelet count noted during the period of thrombocytopenia.

Identification of isolated organism was done based on colony characteristic after overnight incubation on sheep blood agar.

Statistical analysis

Descriptive statistical analysis was done to analyze data. Qualitative data were represented by percentage and quantitative data were represented by mean with standard deviation. Chi-square test for qualitative data and Independent 't' test was applied for quantitative data. Statistical analysis was done using Open- Epi software.

Results

Total 100 babies with culture positive sepsis were analyzed. Mean gestational age of study group was 34.98 ± 3.55 weeks; mean birth weight was 1.899 ± 0.72 kg; male: female ratio was 1.22: 1. Mode of delivery was normal vaginal delivery (NVD) in 69% neonates. Mean APGAR score of study group was 7.81 ± 1.33 . Our study observed early onset sepsis in 72% babies and Gram negative sepsis in 75% babies.

Table 3. Overall thrombocytopenia was observed in 90% babies with 57% having moderate to severe degree of thrombocytopenia. The proportion of severe degree of thrombocytopenia (17.33% Vs 4%), higher MPV (59.33% Vs 56%) and longer duration of thrombocytopenia (3.78 ± 0.55 Vs 3.02 ± 0.61) was observed more with Gram negative sepsis than with Gram positive sepsis, but statistical significance was not found. We observed statistically significant platelet nadir with Gram negative sepsis.

Table 4. Out of 100 neonates with culture positive sepsis, Gm negative sepsis was observed in (75) 75% and Gm positive sepsis was observed in (25) 25%. The proportion of isolated organisms was: 30% klebsiella, 25% pseudomonas, 19% Coagulase- negative Staphylococci (CONS), 14% Citrobacter, 6% S. Aureus & 6% Acinetobacter.

Thrombocytopenia was observed in 96.67% babies with Klebsiella and S. Aureus sepsis followed by 96% pseudomonas and 83.33% Acinetobacter sepsis. Severe thrombocytopenia was observed in 66.67% babies with Acinetobacter sepsis, 17.24% Klebsiella sepsis, 16.67% Pseudomonas sepsis and 6.67% CONS sepsis. In S. Aureus sepsis none of the babies had severe thrombocytopenia. This observation reflects that proportion of thrombocytopenia and severity of thrombocytopenia was more with Gm negative organism.

Acinetobacter sepsis had highest proportion (66.67%) of babies with severe thrombocytopenia, 50% had high MPV value with mean MPV 10.87 ± 0.69 which is higher

than other isolates, had longer duration of thrombocytopenia (4.44 ± 0.49 days) and maximum platelet nadir.

Klebsiella sepsis had 2nd highest proportion (17.24%) of severe thrombocytopenia, had greater fall in platelet count after 48 hour of onset of sepsis, 79.31% babies having higher MPV with mean MPV 10.77 ± 0.88 . The duration of thrombocytopenia (4.51 ± 0.42 days) was longer (2nd in order) and platelet nadir was 3rd in order.

Pseudomonas sepsis had 16.67% babies having severe thrombocytopenia, 58.33% had high MPV with mean MPV value 10.77 ± 0.77 . Platelet nadir and duration of thrombocytopenia was less as compared to Acinetobacter & Klebsiella sepsis.

Among Gram positive organism S. Aureus sepsis observed thrombocytopenia in 5 (83.33%) babies. The other platelet parameter like severity of thrombocytopenia, duration of thrombocytopenia, platelet nadir was less affected as compared to other organism.

Table 1: Baseline characteristics: Demographic profile

Neonatal characteristics	n (%) N = 100 Gestational age
< 28 week	5 (5)
28– 32 week	25 (25)
33– 37 week	25 (25)
> 37 week	45 (45)
Mean gestational age = 34.98 ± 3.55 Birth weight	
< 1 Kg	5 (5)
1 – 2 Kg	50 (50)
2 – 3 Kg	40 (40)
3 Kg	5 (5)
Mean Birth weight = 1.899 ± 0.72 Gender	
Male	55 (55)
Female	45 (45)
APGAR score <5 at 5 minute	10 (10)
Mean APGAR score at 5 minute	7.81 ± 1.33
Mode of Delivery	
NVD	69 (69)
LSCS	31 (31)

Table 2: Baseline characteristics: Type of sepsis

Onset of sepsis	N = 100, n (%)
Early onset sepsis	72 (72)
Late onset sepsis	28 (28)
Gram stain based sepsis	
Gram positive	25 (25)
Gram negative	75 (75)

Table 3: Incidence & degree of thrombocytopenia

Platelet parameters	Overall (N=100)	Gram positive (n=25)	Gram negative (n=75)	P value
Thrombocytopenia	90 (90)	20 (80)	70 (93.33)	0.11
Degree of thrombocytopenia				
Mild	33 (33)	8 (32)	32 (42.67)	0.18
Moderate	39 (39)	11 (44)	25 (33.33)	
Severe	18 (18)	1 (4)	13 (17.33)	
High MPV	63 (63)	14 (56)	52 (69.33)	0.43
Mean duration of thrombocytopenia (in days)	3.36 ± 0.67	3.02 ± 0.61	3.78 ± 0.55	0.69
Lowest platelet count (in lacs)	0.25± 0.17	0.28 ± 0.17	0.22 ± 0.11	0.0001

Table 4: Comparison of platelet parameters among different organisms

	Klebsiella	Pseudomonas	Citrobacter	CONS = 19	Staph. Aureus	Acinetobacter
	30(30%)	(25%)	(14%)	(19%)	6(6%)	(6%)
Thrombocytopenia (occurrence)=90	29 (96.67)	24 (96)	12 (85.71)	15 (78.95)	5 (83.33)	5 (83.33)
Degree of thrombocytopenia						
Mild	9 (31.03)	8 (33.33)	10 (83.33)	5(33.33)	3 (60)	1 (16.66)
Moderate	15 (51.72)	12 (50)	2 (16.67)	9 (60)	2 (40)	1 (16.66)
Severe	5 (17.24)	4 (16.67)	0	1 (6.67)	0	4 (66.67)
Platelet count						
Baseline	1.69±0.58	1.95±0.88	2.25±0.74	2.43±1.36	2.81±0.52	1.89±1.12
At onset of sepsis	1.53 ±0.82	1.47±0.82	1.93±0.57	1.53±0.68	1.55±0.55	1.39±1.67
After 48 hour of onset of sepsis	1.21 ±0.49	1.17±0.63	1.21±0.43	1.33±0.66	1.53 ± 0.36	1.11±0.43
MPV						
Normal	6 (20.69)	10 (41.67)	9 (75)	8 (53.33)	3 (60)	3 (50)
High	23 (79.31)	14 (58.33)	3 (25)	7 (46.67)	2 (40)	3 (50)
Mean MPV	10.77±0.88	10.77±0.77	10.02 ± 0.43	10.23±0.88	10.25±1.11	10.87±0.69
Duration of thrombocytopenia (in days)	4.51 ± 0.42	3.17 ± 0.17	2.52 ± 0.93	3.3 ± 0.12	2.8 ± 0.11	4.44 ± 0.49
Platelet Nadir	0.25±0.16	0.28±0.17	0.19±0.13	0.22±0.18	0.27±0.28	0.17±0.2

Discussion

Thrombocytopenia is one of the most common hematological manifestations in neonatal sepsis.¹⁶ The cause of thrombocytopenia in sepsis can be due to increased platelet destruction, decreased platelet production or combination of both. Neonates respond to sepsis by up regulation of thrombopoietin (TPO) production; however the degree of up regulation is only modest. The study has found that Gram negative sepsis didn't have the highest degree of up regulation despite more significant level of thrombocytopenia and more severe illness. It was suggested that during severe illness there is down regulation of thrombopoietic response.^{17,18} The present study showed Gram negative sepsis in 75% and Gram positive sepsis in 25% neonates. The predominance of Gram negative sepsis is consistent with other Indian studies.¹⁹ Among Gram negative organisms *Klebsiella* was the predominant isolates which is also comparable with earlier reports from India.^{20,21} The observed 88% proportion of thrombocytopenia with Gram negative sepsis in our study is consistent with observation by Bhat et al.²² (70%), Sartaj et al. (66%)²³ and Ree IMC et al.²⁴ (69%). The proportion of severe thrombocytopenia was 18% with Gram negative sepsis as compared to 4% with Gram positive sepsis in our study. Similar observation of severe degree of thrombocytopenia with Gram negative sepsis was found by P Ahmed et al.²⁵, R Bhat et al.²² Ree IMC et al.²⁴ in their study found that Gram negative sepsis had severe degree of thrombocytopenia, platelet count seems to fall lower and time for platelet to rise to > 100,000/cmm was longer than Gram positive sepsis. In our study we also observed longer duration of thrombocytopenia and maximum platelet nadir with Gram negative sepsis and among different platelet parameters statistical significance was observed only in platelet nadir between Gram negative and Gram positive sepsis. Manzoni P et al.²⁶ in their study found 17% septic neonates had associated thrombocytopenia and proportion of thrombocytopenia was 19% with fungal sepsis, 16% with bacterial sepsis. In their study they didn't observe significant difference in platelet parameters when clustering for sepsis caused by Gram positive and Gram negative organisms done.²⁶

The organism specific platelet response was studied to some extent by other researchers. Accordingly proportion of thrombocytopenia in *Klebsiella* sepsis was found to be 60% by Charoo et al.²⁷, 73% by S Arif et al.²⁸ and 43% by Sartaj et al.²³ We found 100% neonates having thrombocytopenia with *Klebsiella* sepsis. The effect of *Klebsiella* organism to various platelet parameters is explained by variation in genetic make up of O antigen between *Klebsiella pneumoniae* and other Gram negative organism.²³

The isolation rate of *Acinetobacter* organism in Indian literature ranges from 8.3% to 15.2%.^{17,23,28} We found *Acinetobacter* isolates in 4% of babies with sepsis. Thrombocytopenia with *Acinetobacter* sepsis was seen in 66% neonates by Bhat et al.²² In our study we found 74% rate of thrombocytopenia in case of *Acinetobacter* sepsis. Sartaj et al.²³ found 11% and S Arif²⁸ found 6% rates of thrombocytopenia with *Acinetobacter*. The platelet parameters were maximally altered in neonates with *Acinetobacter* sepsis in our study. In a neonate with clinical features of sepsis along with altered platelet parameters, it can be suggested to consider Gram negative organism especially *Klebsiella* or *Acinetobacter* at our setup. *Acinetobacter* being a nosocomial infection stresses the need for continuous bacteriological surveillance and implementation as well as adherence to strict infection control policy.

The other Gram negative organism *Pseudomonas* had 88% rate of thrombocytopenia in our study which is comparable to rate of 67% seen by Bhat R et al.²²

Among Gram positive organism we found proportion of thrombocytopenia at the rate of 100% with *S. Aureus* and 66% with CONS sepsis. S Arif had 67% rate of thrombocytopenia with *S. Aureus* and 25% with CONS sepsis.²⁷

Conclusion

Thrombocytopenia is a frequent occurrence in neonates with sepsis especially with Gram negative organism. Sepsis with *Acinetobacter*, *Klebsiella* & *Pseudomonas* organism was associated with prolonged duration, higher MPV and lower platelet count as compared to other isolated organisms.

Reference

1. Stoll BJ. The global impact of neonatal infection. *Clin Perinatol* 1997;24:1-21.
2. UNICEF. The state of the World's Children 2012. 87-107.
3. Gotoff SP. Infections of the neonatal infant. In: Behrman RE, Kliegman RM, Jenson HB, editors. *Nelson textbook of pediatrics*. 16th edition. Philadelphia: Saunders, Elsevier 2000:538-52.
4. Vergnano S, Sharland M, Kazembe P, Mwansambo C, Heath P. Neonatal sepsis: an international perspective. *Arch Dis Child Fetal Neonatal Ed* 2005; 90:F220-24.
5. Mannan MA, Shahidullah M, Noor MK, Dey AC, Nasrin N, Marma U. Nosocomial infections in a newborn intensive care unit of a tertiary care health. *Bangladesh J Child Health* 2008; 32(3):92-96.
6. Khatua SP, Dsas AK, Chatterjee BD, Khatua S, Ghose B, Saha A. Neonatal septicaemia. *Indian Pediatr* 1986; 53:509-14.
7. Hossain MM, Afroza S, Shirin M, Chowdhury NA, Saha SK. Bacterial aetiology of neonatal sepsis in a tertiary care hospital in Bangladesh. *Bangladesh J Child Health* 2004; 28(3):81-85.
8. Mokuolu AO, Jiya N, Adesiyun OO. Neonatal septicaemia in Ilorin: bacterial pathogen and antibiotic sensitivity pattern. *Afr J Med Sci* 2002; 31:127-30.
9. Ng PC. Diagnostic markers of infection in neonates. *Arch Dis Child Fetal Neonatal Ed* 2004; 89:F229-35.
10. Chiesa C, Pellegrin G, Panero A, Osborn JF, Signore F, Assumma F, et al. C-reactive protein, interleukin 6 and procalcitonin in immediate postnatal period: influence of illness severity, risk status, antenatal and perinatal complications and infection. *Clin Chem* 2003;49:60-68.
11. Guida JD, Kunig AM, Leef KH, McKenzie SE, Paul DA. Platelet count & sepsis in very low birth weight neonates: Is there an organism- specific Response? *Pediatrics* 2003; 111 (6): 1411-15.
12. Shirin M, Hossain MM, Mamun MAA, Chowdhury NA, Qader A. Sensitivity and specificity of C-reactive protein and thrombocytopenia is the diagnosis of neonatal sepsis. *Bangladesh J Child health* 2005; 29(2):41-45.
13. Sola MC, Vecchio AD, Rimsza L. Evaluation & treatment of thrombocytopenia in the neonatal intensive care unit. *Clin Perinatol* 2000; 27: 655-79.
14. Scheifele DW, Olsen EM, Pendray MR. Endotoxemia & thrombocytopenia during neonatal necrotizing enterocolitis. *Am J clin pathol* 1985; 83: 227-29.
15. Benjamin DK Jr, Ross K, Mckinney RE Jr, Benjamin DK, Auten R, Fisher RG. When to Suspect fungal infection in neonates;clinical comparison of candida albicans & candida parapsilosis fungemia with coagulase-negative staphyococcal bacteremia. *Pediatrics* 2000; 106: 712-18.
16. Storm W. Use of thrombocytopenia for the early identification of sepsis in critically ill newborns. *Acta Pediatr Acad Sci Hung*.1982;23(3):349-35.
17. Sola M, Sallmon H, Brown R. New insights into the mechanism of non immune thrombocytopenia in neonates. *SeminPerinatol*. 2009;33(1):43-51.

18. Bhat R. Platelet indices in neonatal sepsis: A review. *World J Clin Infect Dis.* 2017; 7(1):6-10.
19. Zaidi AK, Huskins WC, Thaver D, Bhutta ZA, Abbas Z, Goldmann DA. Hospital acquired neonatal infections in developing countries. *Lancet.* 2005;365(9465):1175-88.
20. Brown RE, Rimsza LM, Pastos K, Young L, Saxonhouse MA, Bailey M et al. Effects of sepsis on neonatal thrombopoiesis *Pediatr Res.* 2008;64(4):399-404.
21. Muley VA, Ghadage DP, Bhore AV. Bacteriological profile of neonatal septicemia in a tertiary care hospital from western India. *J Glob Infect Dis.* 2015;7(2):75-7.
22. Bhat R, Kousika P, Lewis L, Purkayastha J. Prevalence and severity of thrombocytopenia in blood culture proven neonatal sepsis: A prospective study. *Arch Pediatr Infect Dis.* 2018;6(2):e12471.
23. Bhat S, Naik S, Rafiq W, Tariq S. Incidence of thrombocytopenia and changes in various platelet parameters in neonates with blood culture positive sepsis. *Int J Pediatr.* 2015; 3(4.1):757-766.
24. Ree IMC, Fustolo-Gunnink SF, Bekker V, Fijnvandraat KJ, Steggerda SJ, Lopriore E. Thrombocytopenia in neonatal sepsis: Incidence, severity and risk factors. *PLoS ONE.* 2017;12 (10): e0185581.
25. Ahmed P, Kaith R, Gattoo I, Najar B, Hussain S. Thrombocytopenia as a predictor of neonatal sepsis in very low birth weight babies and its correlation with specific organism involved: A hospital based observational study. *IJNMR.* 2015; 3(3):7-13.
26. Manzoni P, Mostert M, Galletto P et al. Is thrombocytopenia suggestive of organism specific response in neonatal sepsis? *pediatr Int.* 2009;51(2):206-10.
27. Charoo BA, Iqbal JI, Iqbal Q, Mushtaq S, Bhat AW, Nawaz I. Nosocomial sepsis induced late onset thrombocytopenia in a neonatal tertiary care unit: A prospective study. *Hematol oncol stem cell ther.* 2009;2(2):349-53.
28. Arif SH, Ahmed I, Ali SM, Khan HM. Thrombocytopenia and bacterial sepsis in neonates. *Indian J Hematol Blood Transfus.* 2012;28(3):147-151.

Received: 10-07-2020 // Revised: 30-08-2020 // Accepted: 23-09- 2020