

Original research article

The clinic-epidemiological profile of enteric fever cases and trends of antibiotic susceptibility in paediatric age group

Dr. Prem Prakash¹, Dr. Bipin Kumar Verma²

¹Assistant Professor, Department of Pediatrics Jannayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India

²Associate Professor, Department of Pediatrics Jannayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India

Corresponding Author: Dr. Bipin Kumar Verma

Abstract

Aim: This study was conducted to evaluation of clinic-epidemiological profile and patterns of antibiotic sensitivity of enteric fever cases among paediatric patients.

Methods: A prospective observational study was conducted in the Department of Pediatrics Jannayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India for 1 year. Total 220 Children aged below 18 years with history of fever of more than 7-10 days duration were included in this study. In each case, age, sex, presenting complaint, laboratory investigations and antibiotic sensitivity pattern are collected and analysed.

Results: Out of 220 cases, 150 cases (68.18%) were males and 70 cases (31.82%) were females. In our study, the most common symptom was fever (100%), followed by anorexia (63.64%), vomiting (44.09%), pain abdomen (20%), diarrhea (10%), headache (9.09%), and cough (5%). The most common sign we observed was toxic look in 75% of the cases followed by coated tongue in 49.09%, hepatomegaly 40.91%, splenomegaly 20.91%, hepatosplenomegaly in 14.55% of cases and pallor in 5.91% of cases. Blood culture positive for *Salmonella typhi* noted in 44(20%) cases. The most common sensitivity was seen in all the cases ceftriaxone (96%) and followed by cefixime(94%) , ofloxacin (92%), ciprofloxacin (79%), chloramphenicol (78%), Cefotaxime (75%), amoxicillin (62%) and azithromycin in (50%). *S. typhi* was more sensitive to ceftriaxone, cefixime followed by ofloxacin. Least sensitivity was seen with azithromycin

Conclusion: Typhoid fever is a multisystem disease with variable clinical presentations. No sign or symptom is specific for its diagnosis which needs correlation with lab investigations which again are not always helpful. Polymicrobial resistance mandates heightened focus on preventive measures.

Keywords: Children, Clinical profile, Coated tongue, Typhoid fever

Introduction

Enteric fever, a systemic infection by *Salmonella enterica* serotype Typhi (*S. Typhi*) or *Salmonella enterica* serotype Paratyphi A (*S. Paratyphi A*), affects around 11-21 million individuals globally with a high mortality. The term enteric fever includes typhoid fever caused by *S.typhi* and paratyphoid fever caused by *S.paratyphi A, B and C*. Detailed study of enteric fever was presented by Bretonneau (1826) who identified intestinal lesions. The name typhoid fever was given by Louis (1829) to distinguish it from typhus fever. Eberth (1880) described typhoid bacillus.¹ According to WHO Confirmed case of typhoid fever is defined, as a patient with fever (> 38°C) that has lasted for at least three days, with a laboratory confirmed positive culture of *S. typhi*.² Probable case of typhoid fever is a patient with fever (>38°C) that has lasted for > 3 days, with a positive serodiagnosis or antigen detection test but without *S. typhi* isolation.² The world sees approximately 22 million new typhoid cases

occur each year. In India in disease-endemic areas, the annual incidence of enteric fever is about 1%.³ Complete blood counts in enteric fever are found to be unremarkable. The hemoglobin is normal in the initial stages but drops with progressing illness. Severe anemia is unusual and should make one suspect intestinal hemorrhage or hemolysis or an alternative diagnosis like malaria. The WBC count is normal in most cases and leucocytosis makes the diagnosis less probable. Leukopenia is perceived to be an important feature of typhoid fever and has been reported in only 20-25% cases.⁴ The differential count is usually unremarkable except for eosinopenia. Eosinopenia often absolute may be present in 70-80% cases.⁵ Presence of absolute eosinopenia offers a clue to diagnosis but does not differentiate enteric fever from other acute bacterial or viral infections. A normal eosinophil count does make typhoid fever a less likely possibility. Platelet counts are normal to begin with and fall in some cases by the second week of illness. Overall prevalence of thrombocytopenia is around 10-15%.⁴ The emergence of strains of *Salmonella typhi* resistant to multiple antibiotics poses a serious problem. Chloramphenicol was considered the antimicrobial gold standard for the treatment of typhoid fever till 1948.⁶ But in the last two decades there has been increase in the resistance of strains of *S. typhi* to chloramphenicol. It was first reported in Britain, in 1950 and in India in 1972.^{7,8} Gradually, resistance to multiple antibiotics developed.⁹ The first major epidemic of multidrug resistant *S. typhi* was reported in 1972.¹⁰ in Mexico. Since then, an increasing frequency of antibiotic resistance has been reported from all parts of the world, but more so from the developing countries.⁹ The uses of chloramphenicol, ampicillin and co-trimoxazole have become infrequent and quinolones have become the first line of treatment of typhoid fever. It is one of the common causes of fever in children with varied presentation and significant difference in the signs and symptoms compared to adults. The classic Widal agglutination test is one of the most utilized diagnostic tests for typhoid fever, especially in developing countries. This study was conducted to evaluation of clinic-epidemiological profile and patterns of antibiotic sensitivity of enteric fever cases among paediatric patients.

Material and methods

This prospective observational study was carried out in the Department of Pediatrics Jannayak Karpoori Thakur Medical College and Hospital, Madhepura, Bihar, India for 1 year, after taking the approval of the protocol review committee and institutional ethics committee. After taking informed consent detailed history was taken from the patient.

Total 220 Children aged below 18 years who presented to the Paediatric department with history of fever of more than 7-10 days duration were included in this study. Previously antibiotic treated patients and patients with proven localised infection were excluded. These cases were either Widal positive (Widal test TO Titer >1:100 or TH titre >1:200) or blood culture positive for *Salmonella* species. The cases which were discharged against medical advice and cases for which consent was not obtained were excluded from the study. Antibiotics were started in each case after blood was drawn for Widal test and blood culture for *Salmonella* species. Each case was followed up clinically for improvement. For those cases which did not show improvement after 5 days of antibiotics, changes made according to the culture reports. Antibiotic sensitivity pattern was noted for culture positive cases. Cases were followed till discharge. The data collected was analyzed with respect to age, sex and presenting complaints.

Results

In this study, all the cases presented to OPD with a median of 7-10 days duration of fever. 150 cases (68.18%) had received antibiotics for a minimum period of 3-5 days prior to admission. Table 1 shows Out of 220 cases, 150 cases (68.18%) were males and 70 cases

(31.82%) were females. Table 2 shows, most of the cases were aged between 6 and 12 years. 51 cases were below 6 years, representing 23.18%. 54 cases were aged above 12 years, representing 24.55%. 115 cases were aged between 6 and 12 years (52.27%). In all the above age groups male predominance was seen.

Table 1: Sex Distribution of patients

Sex	N=220	%
Male	150	68.18
Female	70	31.82

Table 2: Age wise distribution

Age(years)	N=220	%
0-6 years	51	23.18
6-12 years	115	52.27
12-18 years	54	24.55

Duration of hospital stay varied from up to two week. As shown in Table-3, most of the cases (72.73%) stayed in hospital up to two week after admission. 27.27% cases stayed up to one Week day in hospital. In these cases, fever persisted beyond one Week. No mortality was observed during our study period. Although mild elevated liver enzymes were observed in some cases, no complications were seen in any case.

Table 3: Duration of hospital stay

Duration of hospital stay	No. Of cases	P-value
One Week	60 (27.27%)	0.18
More than one Week	160 (72.73%)	0.00

Typhoid fever presents with a wide range of symptoms. Due to the use of antibiotics prior to diagnosis, children may not present with typical symptoms. However, in our study, the most common symptom was fever (100%), followed by anorexia (63.64%), vomiting (44.09%), pain abdomen (20%), diarrhea (10%), headache (9.09%), and cough (5%)

Table 4: Common presenting symptoms

Presenting symptom	No. of Cases	P-value
Fever	220 (100%)	0.001
Anorexia	140 (63.64%)	0.000
Vomiting	97 (44.09%)	0.001
Pain abdomen	44 (20%)	0.016
Diarrhea	22 (10%)	0.068
Headache	20 (9.09%)	0.89
Cough	11 (5%)	0.154

Table 5: Various physical findings

Signs	No. of Cases	P-value
Toxic look	156 (75%)	0.001
Coated tongue	108(49.09%)	0.001
Hepatomegaly	90 (40.91%)	0.002
Splenomegaly	46 (20.91%)	0.051
Hepatosplenomegaly	31(14.55%)	0.082
Pallor	13(5.91%)	0.213

Coming to physical findings, the most common sign we observed was toxic look in 75% of the cases followed by coated tongue in 49.09%, hepatomegaly 40.91%, splenomegaly 20.91%, hepatosplenomegaly in 14.55% of cases and pallor in 5.91% of cases.

Table 6: Laboratory parameters

Laboratory parameters	Abnormal values	No. of cases	P-value
Hemoglobin	Anemia (Hb<11g%)	44 (20%)	0.033
Total leukocyte count	Leucocytosis (>11000cells/mm ³)	40 (18.18%)	0.027
	Leucopenia (<4000cells/mm ³)	70 (31.82%)	0.00
Polymorphs	Neutropenia	88 (40%)	0.00
	Neutrophilia	69 (31.36%)	0.00
Eosinophils	Eosinophilia	17 (7.73%)	0.18
	Eosinopenia	98 (44.55%)	0.00
Platelets	Thrombocytopenia	32 (14.55%)	0.01
SGOT	Elevated SGOT	27 (12.27%)	0.23
SGPT	Elevated SGPT	32 (14.55%)	0.25
Widal titres	TO >1:100	211 (95.91%)	0.00
	TH >1: 200	168 (76.36%)	0.00
Blood culture positive	Salmonella	44 (20%)	0.01

Table 6 depicts the laboratory parameters. Anemia found in 44 (20%) cases, leucopenia and leucocytosis was observed in 70(31.82%) cases and 40(18.18%) cases respectively. Neutropenia found in 88(40%) cases and neutrophilia was found in 69(31.36%) cases. Eosinopenia was seen in 98(44.55%) cases, eosinophilia in 17(7.73%) cases and thrombocytopenia in 32(14.55%) cases. SGOT levels was elevated (>200IU/ml) in 27(12.27%) cases and SGPT (>200IU/ml) in 32(14.55%) cases. The elevated levels of liver enzymes lasted only few days. There were no complications observed during our study period. *Salmonella typhi* O titres >1:100 was seen in 211(95.91%) cases and TH titres >1:200 in 168(76.36%) cases. Blood culture positive for *Salmonella typhi* noted in 44(20%) cases. Out of 220 cases only 74 cases had been immunized with typhoid vaccine. All of them had taken typhoid polysaccharide vaccine more than 3 years prior to illness.

Table 7: Antibiotic sensitivity pattern

Drug	Sensitivity	P-value
Ceftriaxone	96%	0.000
Cefixime	94%	0.000
Ofloxacin	92%	0.000
Chloramphenicol	78%	0.000
Cefotaxime	75%	0.002
Azithromycin	50%	0.103
Ciprofloxacin	79%	0.001
Amoxicillin	62%	0.046

Significant p<0.01

Table 7 depicts antibiotic sensitivity patterns among culture positive cases. the most common sensitivity was seen in all the cases ceftriaxone (96%) and followed by cefixime(94%) , ofloxacin (92%), ciprofloxacin (79%), chloramphenicol (78%), cefotaxime (75%), amoxicillin (62%) and azithromycin in (50%). *S. typhi* was more sensitive to ceftriaxone, cefixime followed by ofloxacin. Least sensitivity was seen with azithromycin.

Discussion

Typhoid fever is a common infectious disease presenting as acute multisystem febrile illness. Our study has presented detailed study of 220 cases of typhoid fever about symptoms, clinical signs, investigations and in vitro antibiotics sensitivity/resistance pattern and treatment response. The definitive diagnosis of typhoid fever requires a confirmed diagnosis based on the blood or bone marrow culture. However, blood culture has several limitations including amount of blood required due to low levels of bacteremia and prior antibiotic use.¹¹ All the cases presented to OPD with a median of 7-10 days duration of fever.

150 cases (68.18%) had received antibiotics for a minimum period of 3-5 days prior to admission. Out of 220 cases, 150 cases (68.18%) were males and 70 cases (31.82%) were females. Similar results were reported in other studies.¹² most of the cases were aged between 6 and 12 years. 51 cases were below 6 years, representing 23.18%. 54 cases were aged above 12 years, representing 24.55%. 115 cases were aged between 6 and 12 years (52.27%). In all the above age groups male predominance was seen. A study done by R Modi et al also reported maximum incidence of typhoid in the age group 6 to 10 year.¹³ Another study also reported maximum number of cases in the age group above 5 years.¹⁴ The duration of hospital stay varies, with maximum number of cases staying in hospital between 8-10 day. Cases were discharged after 3 consecutive days of afebrile period without antipyretics. These results were in accordance with study done by Hyder et al.¹⁵ We observed high incidence of typhoid fever in lower class, lesser in middle class society and least in higher class. This can be explained by differences in drinking water sources and hygienic practices like hand washing and sanitary latrine facilities. Similar results were reported in other study.¹⁶ in our study, the most common symptom was fever (100%), followed by anorexia (63.64%), vomiting (44.09%), pain abdomen (20%), diarrhea (10%), headache (9.09%), and cough (5%). A study done by Sinha A et al.¹⁷ Kapoor JP et al also reported similar results.¹⁸ Other studies also showed similar clinical picture.¹⁹⁻²¹ Contradictory to this, a study done by Joshi et al. reported headache as the most common symptom next to fever.²² In our study the most common sign we observed was toxic look in 75% of the cases followed by coated tongue in 49.09%, hepatomegaly 40.91%, splenomegaly 20.91%, hepatosplenomegaly in 14.55% of cases and pallor in 5.91% of cases. Study done by Laishram et al reported coated tongue (80%) as the most common sign followed by Hepatomegaly (76%) and splenomegaly (38%).²³ In other study they had reported relative bradycardia and hepatomegaly as the most common sign.²⁴

During our study, all cases were positive for Widal. Blood culture was positive in 20% of cases. Other study also reported 16% culture positive cases.¹⁵ A study done by Banu et al also reported 28% culture positive cases.²⁴ Due to prior use of antibiotics, the culture positive cases are decreasing. Thus, need for relay on other serological tests for diagnosis of typhoid exists. Study done by Modi et al reported 97% Widal positive cases.¹³ Anemia was seen in 22.5% of cases. The other studies reported little higher percentage of anemias. A study done by Raj C et al reported anemia in 41.8% of patients and Lefebvre et al reported anemia in 78% of cases.^{25,26} in our study Leucocytopenia and Eosinopenia found in 70(31.82%) and 98(44.55%) respectively. Similar results reported in Lefebvre et al.²⁶ Although leucocytosis and eosinophilia are rare in typhoid, our study reported leucocytosis in 40(18.18%) of cases

and eosinophilia in 17(7.73%) cases respectively. Thrombocytopenia was found in 32(14.55%) of cases. SGOT levels was elevated (>200IU/ml) in 27(12.27%) cases and SGPT (>200IU/ml) in 32(14.55%) cases. The other study reported elevated liver enzymes in 70% of cases.²⁷

Antibiotic sensitivity was similar to other studies. Most of the culture positive cases showed sensitivity to ceftriaxone, cefixime, ofloxacin, ciprofloxacin. Similar sensitivity pattern reported in other study.²² However sensitivity pattern varies from place to place. Other studies showed return of sensitivity pattern with chloramphenicol, cotrimoxazole, amoxicillin.^{20,28,29} A study done by Mishra et al reported 100% sensitivity to azithromycin.³⁰ In our study the sensitivity to azithromycin was 50%. A Study done by Hyder et al reported 100% sensitivity to ceftriaxone and ciprofloxacin.¹⁵

Conclusion

Typhoid fever is a multisystem disorder with a wide variety of clinical symptoms. There is no clear indication or symptom for diagnosis, which necessitates correlation with lab investigations, which are not necessarily beneficial. Polymicrobial tolerance necessitates a greater emphasis on prevention.

Reference

1. Ananthanarayan R, Panikar CKJ. Text Book Of Microbiology, 9th Edition; Chennai: Orient Longman; 2013.
2. Background document: The diagnosis, treatment and prevention of typhoid fever. Communicable Disease Surveillance and Response Vaccines and Biologicals. World Health Organization [Internet] [cited 2015 April 6]. Available from: <http://www.who.int/rpc/TFGuideWHO.pdf>
3. Darton TC, Blohmke CJ, Pollard AJ. Typhoid epidemiology , diagnostics and the human challenge model. *Curr Opin Gastroenterol* 2014; 30:7-17.
4. Lokhandwala ,Syed Athar,Nicolas P Turin. Role of Absolute Eosinopenia as Marker of Enteric Fever: Experience from a Tertiary Care Hospital in the United Arab Emirates Aliasgar Ibnosina *J Med BS* 2012:249-253
5. Kuvandik C, Karaoglan I, Namiduru M, Baydar I. Predictive value of clinical and laboratory findings in the diagnosis of the enteric fever. *New Microbiol.* 2009 Jan;32(1):25-30.
6. Wood Ward TE, Smadel JE, Ley HL, Green R. Preliminary report on beneficial effects of chloromycetin in treatment of typhoid fever. *Ann Int Med* 1948;**29** ;131-4
7. Calquhoun J, Weetch RS. Resistance to chloramphenicol developing during treatment of typhoid fever. *Lancet* 1950;**2** :621
8. Panicker CK, Vimla KM. Transferable chloramphenicol resistance in salmonella typhi. *Nature* 1973;**239** :109
9. Samantray SK. Typhoid fever resistant to furazolidine, Ampicillin, chloramphenicol and co-trimoxazole. *Indian J Med Sci* 1979;**33** ;1-3
10. Edelman R, Levine MM. Summary of an international workshop on typhoid fever. *Rev Infect Dis* 1986;**8** :329-49
11. World Health Organization, Department of Vaccines and Biologicals. Background document: the diagnosis, treatment and prevention of typhoid fever. 19–23 (2003).
12. Sen S.K, Mahakur A.C. Enteric fever-A comparative study of adult and paediatric cases. *Indian J Pediat.* 1972;39(11):354-60.
13. Gosai MM, Hariyani HB, Purohit PH, et al. A study of clinical profile of multidrug resistant typhoid fever in children. *NJIRM.* 2011;2(3):87-90.
14. R Modi. Clinical profile and treatment outcome of typhoid fever in children at a

- teaching hospital, Ahmedabad, Gujarat, India. *Int J Med Sci Public Health*. 2016;5:212-6.
15. Arora RK, Gupta A, Joshi NM, Kataria VK, Lall P, Anand AC. Multidrug resistant typhoid fever: study of an outbreak at Calcutta. *Indian Pediatr*. 1992;29(1):61-6.
 16. Hyder R, Yasmeen B, Ahmed S. Clinical profile and Outcome of Enteric Fever in hospitalized children aged 6 months to 2 years. *Northern Int Med Coll J*. 2013;5(1):301-5.
 17. Sood SC, Taneja PN. Typhoid fever, clinical picture and diagnosis. *Ind J of Child Health*. 1961;10(2):69- 76.
 18. Sinha A, Sazawal S, Kumar R, Sood S, Singh B, Reddaiah VP et al. Typhoid fever in children aged less than 5 years. *Lancet*. 1999;354:734-7
 19. Parry CM, Hien TT, Dougan G, White NJ. Typhoid fever. *N Eng J Med*. 2002;347:1770-82
 20. Kapoor JP, Mohan M, Talwar V, Daral TS, Bhargava SK. Typhoid fever in young children. *Indian Pediatr*. 1985;22:811-3.
 21. Chowta MN, Chowta NK. Study of Clinical Profile and Antibiotic Response in Typhoid Fever. *Indian J Med Microbiol*. 2005;23:125-7.
 22. Kadhiraivan T, Wig N, Kapil A, Kabra SK, Renuka K, Misra A. Clinical outcomes in typhoid fever: Adverse impact of infection with nalidixic acid resistant *Salmonella typhi*. *BMC Infec Diseas*. 2005;5:37.
 23. Joshi BG, Keyal K, Pandey R, Shrestha BM. Clinical profile and sensitivity pattern of salmonella serotypes in children: a hospital based study. *J Nepal Paediatr Soc*. 2011;31(3):180-3.
 24. Laishram N, Singh PA. Clinical profile of enteric fever in children. *J Evolution Med Dent Sci*. 2016;5(2):114-116.
 25. Banu A, Rahman MJ, Suza-ud-doula A. Clinical Profile of Typhoid Fever in Children in Northern Areas of Bangladesh. *Dinajpur Med Col J*. 2016;9(1):53-8.
 26. Raj C. Clinical profile and antibiotic sensitivity pattern of typhoid fever in patients admitted to paediatric ward in a rural teaching hospital. *Int J Med Res Health Sci*. 2014;3(2):245-8
 27. Lefebvre N, Gning SB, Nabeth P, Ka S, Ba-Fall K, Rique M et al. Clinical and laboratory features of typhoid fever in Senegal: A 70-case study. *Med Trop (Mars)*. 2005; 65(6):543-8.
 28. Malik AS, Malik RH. Typhoid fever in malasian children. *Med J Malaysia*. 2001;56(4):478-90.
 29. Gautam V, Gupta N, Chaudhary U, Arora DR. Sensitivity pattern of Salmonella serotypes in Northern India. *Brazilian J Infect Dis*. 2002;6:281-7.
 30. Mishra SK, Sah JP, Shrestha R, Lakhey M. Emergence of Nalidixic acid resistant Salmonella: a confounding scene in antibiotic armamentarium. *J Nepal Med Lab Sci*. 2008;9(1):61-6

Received: 03-09-2020.

Revised: 08-10-2020.

Accepted: 26-10-2020