

Original research article

A questionnaire-based study of diabetic patients' awareness and understanding of diabetic retinopathy at a tertiary care hospital in Bihar State

Dr. Sanjeev Kumar¹, Dr. Kanhaiya Lal Agrawal²

¹Senior Resident, Department of Ophthalmology, Madhubani Medical College and Hospital, Madhubani, Bihar, India

²Assistant Professor, Department of Ophthalmology, Madhubani Medical College and Hospital, Madhubani, Bihar, India

Corresponding Author: Dr. Kanhaiya Lal Agrawal

Abstract

Aim: The aim of the present study was to assess the awareness and knowledge of Diabetic retinopathy among Diabetic patients in a tertiary care hospital of Bihar state.

Methods: This was a descriptive, cross sectional, questionnaire based study conducted in the Department of Ophthalmology, Madhubani Medical College and Hospital, Madhubani, Bihar, India for 12 months. A knowledge attitude practice questionnaire was prepared and pretested in a sample group of representative population. The response was analyzed as to whether the questions were understood or not. Social workers were trained in administering questionnaire. Diabetic patients were given questionnaires at primary health centre and filled in the presence of social workers.

Results: Out of the 120 patients, 65(54.17%) had no knowledge of diabetic retinopathy compared to 55 (45.83%) who had knowledge. Knowledge was more in age group less than 25 years (69.23%) and least in 35 to 45 age group (45%) which was statistically significant with p value <0.001. Knowledge was found to be high among participants with higher educational status than in those who had college level education (66.67%) which was statistically significant with p value <0.001. About 72.73 % of individuals in knowledge group had right attitude which was significantly higher than non knowledge group (58.46%) with a p value <0.001. Regarding source of information, 48% of patients in knowledge group got information about diabetic retinopathy from physicians, 12.5% from eye specialists, 10% from reading books, 10% from various media and 25% from other sources like family and friends. About 18.18% in knowledge group had practice of visiting ophthalmologist for eye check-up which was significantly higher than non knowledge group with a p value <0.001.

Conclusion: we concluded that the Increasing knowledge about diabetic retinopathy through awareness campaigns can improve attitude & practice. Early detection & timely intervention can help in preventing sightthreatening complications

Keywords: Attitude, Diabetic retinopathy, Knowledge, Practice, Primary health centre

Introduction

Diabetes mellitus (DM) is a metabolic disease characterized by alteration in carbohydrate, lipid and protein metabolism, where the body cannot regulate the amount of glucose in the blood.¹ Based on various studies from different countries, the global prevalence of DM was reported to be 8% in 2011 and is expected to rise to 10% by 2030.² The prevalence of DM is reported to be higher in Jordan than in other countries, with the reported prevalence being 17.1% in 2008 and this rate is expected to double in the next 10 years.³ DM can result in many complications such as nephropathy, cardiovascular, neurologic and ocular complications,⁴ with

diabetic retinopathy (DR) being the most common micro vascular ocular complication of DM.⁵ DR is defined as a disorder of the retinal circulation that compromises the delivery of oxygen and nutrients to the retina, thus being unable to meet the requirements of its high metabolic demands.⁶ Therefore, defects in retinal circulation may affect normal vision, which is considered a leading cause of vision impairment and blindness worldwide.^{1,4,7} Wisconsin Epidemiological study has proved that micro vascular complications such as diabetic retinopathy are linked to duration of diabetes.⁸ Routine dilated fundus examination is recommended at the time of diagnosis of diabetes and then yearly review is required in all patients having type 2 diabetes mellitus. Awareness of importance of routine check-up for the screening of diabetic retinopathy is poor even in developed countries and the situation is much worse in a developing country like India. Previous studies have shown that 63% of the rural diabetic population has not had an eye examination.⁹ Magnitude of blindness caused by diabetic retinopathy is increasing. Lack of awareness about diabetic retinopathy and the preventable complications associated with it worsens the situation. Early detection of diabetic retinopathy and its appropriate management is very important to prevent irreversible visual loss. This can only be achieved with better knowledge and awareness among patients. There have been many studies done in other states in India to assess the knowledge and awareness on diabetes and diabetic retinopathy among patients with diabetic retinopathy, however very few studies assessed knowledge, awareness and practices amongst patients with diabetic retinopathy.¹⁰⁻¹³ The objective of the study was to assess the knowledge, attitudes and practices about diabetes and diabetic retinopathy among patients diagnosed with diabetic retinopathy at a tertiary care hospital in Bihar, India.

Material and Methods

This was a descriptive, cross sectional, questionnaire based study conducted in the Department of Ophthalmology, Madhubani Medical College and Hospital, Madhubani, Bihar, India for 12 months, after taking the approval of the protocol review committee and institutional ethics committee.

Methodology

A detailed search in literature was done to create the knowledge attitude and practice questionnaire which was prepared in English and the local language. Questionnaire was tested in a sample group of representative population. The response was analysed as to whether the questions were understood or not. Social workers were trained in administering the questionnaire. They were given the questionnaire and filled in presence of the social workers after obtaining informed consent. Diabetic patients of age less than 18 years, mentally challenged patients who were not able to give informed consent and patients who were not able to understand and respond to the questions administered were excluded from the study.

Questionnaire consisted of three parts, first part contained the patient profile which included name, gender, occupation, socioeconomic status, educational status and their consent for the study. Second part included details of diabetes mellitus like duration, family history of diabetes and any eye complaints if present. Third part of the questionnaire contained the following questions.

1. Do you know that diabetes can affect many organs in the body?
2. Do you know whether diabetes can affect vision?
3. Do you know whether there is any relation between duration of diabetes mellitus and visual problems?
4. Do you feel eye check-ups are necessary in diabetes?
5. Do you know the frequency of eye check up needed; if yes how frequently is it needed?

6. Do you know about the complications of diabetic eye disease?
 7. Have you ever done an eye check up to know whether diabetes has affected your eye?

Knowledge group included participants who responded “yes” to above questions 1,2,3,4 and those who responded “no” to these questions were grouped under non knowledge group. Attitude was assessed by question number 5 and 6. Practice was assessed by question number 7.

Determinants of knowledge on diabetes and diabetic retinopathy such as age, gender, language, literacy and socioeconomic status were analysed between the groups using univariate analysis. The association of knowledge of diabetic retinopathy with attitude and practices was evaluated between the groups using univariate analysis (chi square test). A two tailed P value <0.05 was considered statistically significant.

Results

The demographic characteristics of the 120 patients recruited into the study are given in Table 1. Out of the 120 patients, 65(54.17%) had no knowledge of diabetic retinopathy compared to 55 (45.83%) who had knowledge. This was statistically significant with p value <0.001.

Table 1: Demographic characteristics of the study population

Age	Number=120	Percentage
Below 25	13	10.83
25-35	31	25.83
35-45	40	33.33
45-55	23	19.17
Above 55	13	10.83
Gender		
Male	34	28.33
Female	86	71.67
Educational status		
Uneducated	31	25.83
class 1-12	74	61.67
college level	15	12.5
Socio economic status		
Lower	64	53.33
Middle	37	30.83
Upper	19	15.83

Table 2: Association of age and knowledge of diabetic retinopathy (DR).

	Knowledge of DR			P-value
	Knowledge group	Non knowledge group	Total	
Below 25	9	4	13	
25-35	10	21	31	
35-45	18	22	40	0.00
45-55	9	14	23	01
Above 55	5	8	13	
Total	55	65	120	

Knowledge was more in age group less than 25 years (69.23%) and least in 35 to 45 age group (45%) which was statistically significant with p value <0.001 (Table 2). Knowledge was more

among females than males which was not significant statistically. Knowledge was found to be high among participants with higher educational status than in those who had college level education (66.67%) which was statistically significant with p value <0.001 (Table 3).

Table 3: Association of educational status and knowledge of DR

Educational status	Knowledge of DR			P-value
	Knowledge group	Non knowledge group	Total	
No education	12	19	31	
Class 1-12	34	40	74	0.00
College	7	6	15	01
Total	55	65	120	

Patients in the upper socioeconomic group had more knowledge about diabetic retinopathy (84.21%) which was statistically significant with a p value of 0.001 (Table 4).

Table 4: Association of socio economic status and knowledge of DR

Income Level (monthly income in rupees)	Knowledge of DR		Total	P-Value
	Knowledge group	Non knowledge group		
Lower (200-500)	21	43	64	
Middle (501-2000)	18	19	37	0.0001
Upper (>2000)	16	3	19	
Total	55	65	120	

There was no significant association between duration of diabetes and knowledge of diabetic retinopathy. About 72.73 % of individuals in knowledge group had right attitude which was significantly higher than non knowledge group (58.46%) with a p value <0.001 . (Table 5).

Table 5: Association of knowledge of DR with attitude towards DR.

Attitude	Knowledge of DR			P - Value
	Knowledge group	Non knowledge group	Total	
Yes	40	38	78	
No	15	27	42	0.0001
Total	55	65	120	

Regarding source of information, 48% of patients in knowledge group got information about diabetic retinopathy from physicians, 12.5% from eye specialists, 10% from reading books, 10% from various media and 25% from other sources like family and friends (table 6)

Table 6: Source of information about diabetic retinopathy

Physicians	48%
Eye Specialists	12.5%

Reading Books	10%
Various Media	10%
Family and Friends	25%

Table 7: Association of knowledge of DR with practice regarding DR

Practice	Knowledge of DR		Total	P value
	Knowledgeable	Non knowledgeable		
Yes	10	11	21	
No	45	54	99	0.0001
Total	55	65	120	

About 18.18% in knowledge group had practice of visiting ophthalmologist for eye checkup which was significantly higher than non knowledge group with a p value <0.001 (Table 7).

Discussion

The lack of awareness about diabetic retinopathy in diabetic patients is considered as a major cause of diabetic blindness in our community. The main objective of this study was to ascertain the awareness level and the practice patterns adopted by diabetic patients in the community.

Diabetic retinopathy is an upcoming cause of visual impairment and prevalence of diabetic retinopathy is more in developing countries. Even though health education statistics and literacy rate in Bihar is superior than national average and close to that of developed countries, diabetes related ocular complications are on the rise. The facilities in primary health centres which are provided free of cost are not utilized properly and this is reflected in the results of our study.

In this study Out of the 120 patients, 65(54.17%) had no knowledge of diabetic retinopathy compared to 55 (45.83%) who had knowledge. This was statistically significant with p value <0.001. Results were similar to study by Rani et al in which knowledge about diabetic retinopathy was noted as 37.1% and Dandonna et al who reported it as 27%.^{14,15} In a population-based awareness study in a sub urban area by Hussain R et al, among diabetic patients only 40.7% had knowledge about diabetic retinopathy.¹⁶

In this study, Patients in the upper socioeconomic group had more knowledge about diabetic retinopathy (84.21%) which was statistically significant with a p value of 0.001. Literacy and its influence on knowledge about diabetes was studied in other studies also.^{17,18} All these studies support the fact that providing education can increase awareness and knowledge about diabetic retinopathy. Dandona et al, also reported increased awareness among subjects older than 30 years or more and those with any level of education and among those belonging to upper and middle socio-economic strata in their study in urban population in India.¹⁵ Al Zarea in Saudi Arabia reported that knowledge regarding ocular complications in diabetes was 75.62% which was an urban study.¹⁹

About 72.73 % of individuals in knowledge group had right attitude which was significantly higher than non knowledge group (58.46%) with a p value <0.001. In the study by Rani et al attitude among knowledge group about diabetic retinopathy was 93.3% and this was 53.8% in the study by Hussain et al.^{14,16} Rani et al, noted that 36.5% with knowledge about diabetic retinopathy thought that there was no need to consult an ophthalmologist if their blood sugar was under control and this was 38.49% in Saudi Arabia study.^{14,19}

In this study, 18.18% in knowledge group had practice of going for eye check-up which was statistically significant. Oveneri-Ogbomo et al, also reported that knowledge of diabetic retinopathy was significantly related to practice of undertaking eye examinations.²⁰ Mwangi et al, reported that 50% of the participants in their study went for eye check-up.²¹ Hussain et al, reported that practice was present in 57.6%.¹⁶ In the study by Al Zarea practice was reported to be 95% which was an urban study.¹⁹ Mahesh G et al and Srinivasan N K et al, also found a statistically significant association between awareness of diabetic retinopathy and good practice patterns regarding retinopathy in their studies.^{22,23}

In primary health centres, physicians can play a major role in creating awareness and imparting knowledge about diabetic retinopathy. Data from our study also reflects this. Regarding source of information, 48% of patients in knowledge group got information about diabetic retinopathy from physicians, 12.5% from eye specialists, 10% from reading books, 10% from various media and 25% from other sources like family and friends. Srinivasan NK et al, also reported that doctors (both physicians and ophthalmologists) constituted the most important source of information in 71.4 % in knowledge group in their study.²³ About 66.4 % obtained their knowledge from general practitioners and nurses in the study by Oveneri-Ogbomo et al.²⁰

Knowledge about diabetes and diabetic retinopathy help patients in developing good practice patterns which can prevent sight threatening complications. Strategies to educate diabetic patients about this potentially blinding disease should be evolved. Health education measures should be implemented at primary, secondary and tertiary levels.

At the primary level, this can be done through regular awareness campaigns, posters, pamphlets, diabetic retinopathy screening camps and through community- based education strategies. Hospital based patient education can be done by involving general practitioners, physicians and endocrinologists in addition to ophthalmologists. Data about source of information in our study also correlates with this.

Conclusion

There is a difference in the level of knowledge about diabetes mellitus among patients, the patients were more familiar with general than specific information about diabetes retinopathy, this leads to gaps between knowing and acting, the current study concludes that more awareness campaigns need to be conducted to target DM complications and the appropriate attitude towards this disease, the information must be more inclusive & easy for the general public to understand and include it in the importance topics that doctors must inform DM patients about, It can also be included in the high school activities, hanging awareness posters with the main information in the waiting areas of all hospitals and health care centres. More research needs to be conducted about the awareness levels & the appropriate ways to deliver this information.

Reference

1. Olokoba AB, Obateru OA, Olokoba LB. Type 2 diabetes mellitus: a review of current trends. *Oman Med J.* 2012;27(4):269–273.
2. Guariguata L, Whiting DR, Hambleton I, Beagley J, Linnenkamp U, Shaw JE. Global estimates of diabetes prevalence for 2013 and projections for 2035. *Diabetes Res Clin Pract.* 2014;103(2):137–149.
3. Ajlouni K, Khader YS, Batieha A, Ajlouni H, El-Khateeb M. An increase in prevalence of diabetes mellitus in Jordan over 10 years. *J Diabetes Complications.* 2008;22(5):317–

- 324.
4. Alberti KG, Zimmet PZ. Definition, diagnosis and classification of diabetes mellitus and its complications. Part 1: diagnosis and classification of diabetes mellitus provisional report of a WHO consultation. *Diabet Med.* 1998;15(7):539–553.
 5. Nsiah-Kumi P, Ortmeier SR, Brown AE. Disparities in diabetic retinopathy screening and disease for racial and ethnic minority populations – a literature review. *J Natl Med Assoc.* 2009;101(5):430–437.
 6. Tajunisah I, Wong P, Tan L, Rokiah P, Reddy S. Awareness of eye complications and prevalence of retinopathy in the first visit to eye clinic among type 2 diabetic patients. *Int J Ophthalmol.* 2011;4(5):519–524.
 7. Wong TY, Cheung N, Tay WT, et al. Prevalence and risk factors for diabetic retinopathy: the Singapore Malay Eye Study. *Ophthalmology.* 2008;115(11):1869–1875
 8. Klein R, Klein B, Moss S. The Wisconsin Epidemiological Study of Diabetic Retinopathy: a review. *Diabetes Meta Review.* 1989;5:5559-5570.
 9. Padmaja KR, Rajiv R, Pradeep GP, Swati A, Kumaramanickavel G, Sharma T. Use of eye care services by people with diabetes-South India experience. *British J Ophtahlmology.* 2005
 10. Srinivasan NK, John D, Rebekah G, Kujur ES, Paul P, John SS. Diabetes and diabetic retinopathy: KAP among diabetes patients in a tertiary care centre. *J Clin Diag Res.* 2017; 11(7):1-7.
 11. Hussain R, Rajesh B, Giridhar A, Gopalakrishnan M, Sadasivan S, James J, et al. Knowledge and attitudes about diabetes and diabetic retinopathy in suburban population of a South Indian state and its practice among patients with diabetes mellitus: A population based study. *Indian J Ophthalmol.* 2016;64(4):272-6.
 12. Rani PK, Raman R, Subramani S, Perumal G, Kumaramanickavel G, Sharma T. Knowledge of diabetes and diabetic retinopathy among rural population in India and its influence on knowledge of diabetic retinopathy on attitude and practice. *Rural and remote health.* 2008;8(3):838.
 13. Koshy J, Varghese DL, Mathew T, Kaur G, Thomas S, Bhatti SM. Study on KAP of ocular complications due to diabetes among type 2 diabetes visiting a tertiary teaching hospital. *Indian J Community health.* 2012;24:27-31.
 14. Rani PK, Raman R, Subramani S, Perumal G, Kumaramanickavel G, Sharma T. Knowledge of diabetes and diabetic retinopathy among rural population in India and the influence of knowledge of diabetic retinopathy on attitude and practice. *Rural Remote Health.* 2008;8(3):838.
 15. Dandona R, Dandona L, John RK, McCarty CA, Rao GN. Awareness of eye diseases in an urban population in Southern India. *Bull World Health Organ.* 2001;79(2):96-102
 16. Hussain R, Rajesh B, Giridhar A, Gopalakrishnan M, Sadasivan S, James J, et al. Knowledge and awareness about diabetes mellitus and diabetic retinopathy in suburban population of a South Indian state and its practice among the patients with diabetes mellitus: A population-based study. *Indian J Ophthalmol.* 2016;64(4):272-6.
 17. Williams MV, Baker DW, Parker RM, Nurss JR. Relationship of functional health literacy to patients' knowledge of their chronic disease. A study of patients with hypertension and diabetes. *Archives Inter Med.* 1998;158:166-172.
 18. Rothman RL, Malone R, Bryant B. The spoken knowledge in low literacy in diabetes scale. *The Diabetes Educator.* 2005;31:215-224.
 19. Al Zarea BK. Knowledge, attitude and practice of diabetic retinopathy amongst the diabetic patients of Aljouf and hail province of Saudi Arabia. *JCDR.* 2016;10(5):NC05-NC08.
 20. Oveneri-Ogbomo GO, Abokyi S, Koffuor GA, Abokyi E. Knowledge of diabetes and

- its associated ocular manifestations by diabetic patients: A study at Korle-Bu Teaching Hospital, Ghana. Niger Med J. 2013;54(4):217-23.
21. Mwangi MW, Githinji GG, Githinji FW. Knowledge and awareness of diabetic retinopathy amongst diabetic patients in Kenyatta national hospital, Kenya. Int J Humanit Soc Sci. 2011;1(21):140-46.
 22. Mahesh G, Elias A, Sandhya N, Giridhar A, Saikumar SJ, Sankaranarayanan, et al. Chengamanad diabetic retinopathy awareness study (CDRAS) Kerala J Ophthalmol. 2006;28:14-21.
 23. Srinivasan NK, John D, Rebekah G, Kujur ES, Paul P, John SS. Diabetes and diabetic retinopathy: knowledge, attitude, practice (KAP) among diabetic patients in a tertiary eye care centre. J Clin Diagn Res. 2017 Jul;11(7):NC01-NC07

Received: 10-09-2020 // Revised: 11-10-2020 // Accepted: 09-11-2020